REPORT OF STREAMS EXAMINATION



SANITARY DISTRICT
OF CHICAGO




A. R. Reynolds. M. D.


June 1903











REPORT

OF

STREAMS EXAMINATION

Chemic and Bacteriologic

OF THE

Waters between Lake Michigan at Chicago and the Mississippi River at St. Louis, for the purpose of determining their condition and quality before and after the opening of the Drainage Channel

MADE UNDER THE DIRECTION OF

ARTHUR R. REYNOLDS, M. D.

Commissioner of Health, City of Chicago

PUBLISHED BY AUTHORITY OF

THE TRUSTEES OF THE SANITARY DISTRICT
OF CHICAGO

December, 1902

PRESS OF
THE BLAKELY PRINTING COMPANY
CHICAGO

WELLCONIC HASTITUTE					
Coll.	wall/!Omec				
Cali					
No.	I E JA				

BOARD OF TRUSTEES OF THE SANITARY DISTRICT OF CHICAGO

WILLIAM H. BAKER.

JOSEPH C. BRADEN.

ZINA R. CARTER.

FRANK X. CLOIDT.

ALEXANDER J. JONES.

WILLIAM LEGNER.

THOMAS A. SMYTH.

THOMAS J. WEBB.

FRANK WENTER.

Digitized by the Internet Archive in 2017 with funding from Wellcome Library

LETTER OF TRANSMITTAL FROM THE DIRECTOR OF THE STREAMS EXAMINATION.

Chicago, December 1, 1902.

To the Honorable, the Board of Trustees, Sanitary District of Chicago.—Gentlemen:—In presenting to you the subjoined reports of the bacteriologic and chemical experts on the effects of the Sanitary Drainage Channel on the waters of the streams between Chicago and St. Louis, I beg to call your attention to the facts, which, in my judgment, seem to be demonstrated in these reports.

For the general reader it should be premised, first, that the immediate object of the Chicago Sanitary District, created under the Act of May 29, 1889, is the diversion from Lake Michigan of the sewage of Chicago and its inoffensive disposal toward the Mexican Gulf. These are the primary and all-important purposes of the expenditure of the \$36,000,000: To prevent the further pollution of the waters along the city front and to relieve the filth-congested river and its branches, the contents of which have been aptly described as "the quintessence of seweral putridity," and to do this without offense to the senses or injury to the health of the communities further down stream—communities which have long suffered from the relatively insignificant sewage discharge through the Illinois and Michigan Canal.

Second, and as a corollary of this first premise, the protection, protentianto, of the Chicago water supply from pollution by this sewage.

Third, the reclamation of the "malaria preserves" along the Illinois river, to the benefit of the public health of those regions.

And, ultimately, the improvement of navigation between Lake Michigan and the Mississippi river.

The inception of this work dates back nearly half a century—to the days of E. S. Chesebrough, one of the best-equipped and most far-seeing sanitary engineers of his period, and during the interval it has received the attention and study of such sanitarians and scientists as Dr. John H. Rauch (for fully thirty years), Prof. John H. Long, Dr. F. W. Reilly, Allen Hazen, Rudolph Hering, Geo. E. Waring, Jr., W. T. Sedgwick, John W. Hill, Lyman E. Cooley, Benezette Williams, S. G. Artingstall and others.

Of the facts demonstrated in the report, your attention is especially invited to that which shows that running streams, adequately diluted, do purify themselves from sewage pollution—a proposition first made by Dr. Reilly as a result of his collation of the reports of Prof. Long's investigations of the water supplies of Illinois, 1888-89, and at a time when water analysts of the old school held that "no stream on earth is long enough to purify itself after once being organically polluted." The demonstration is shown in the complete disappearance of any trace of Chicago sewage in the Illinois river long before it reaches Averyville and in the better quality of the Illinois river water as it empties into the Mississippi at Grafton than that of the Mississippi itself.

All talk of Chicago sewage injuriously affecting the drinking water at St. Louis is thus completely and effectually disposed of by the work of these investigators.

The benefit of the Sanitary waterway on the public water supply has not yet been fully realized, and will not be until the intercepting sewers and other works necessary to the exclusion of all sewage from the lake in the vicinity of Chicago are completed. But what may be confidently anticipated is foreshadowed in the great improvement of the sanitary quality of the water supply, under usual meteorologic conditions, since the Channel was opened in January, 1900.

As to the improvement of the river and the south branch, no one who crosses a bridge from Rush street to Robey, or who lives, works or offices in the vicinity of the main river and south branch, can fail to notice the change in the atmosphere since that date.

Concerning the benefits along the Desplaines and Illinois rivers, the State Fish Commissioner, Col. Bartlett, reports an enormous increase in the fish harvest—a crop more renumerative, acre for acre, than any other in the state. Fish that have been driven away from increasing reaches of the river year by year by the undiluted sewage of the Illinois and Michigan Canal and of the larger towns below Chicago are returning to the purified waters, and the denizens along the banks of the Illinois, erstwhile hostile to the Sanitary District, are now clamoring for the fullest flow of the channel in the interest of improved navigation.

As part of the history of the Sanitary District, I incorporate in this letter of transmittal the correspondence and other data pertaining to the streams examination of which I have had the honor of being the Director. This begins with the following:

Chicago, November 28, 1808.

To the Honorable, the Board of Trustees, Sanitary District of Chicago.

Gentlemen:—I take occasion again to urge—as I have previously done verbally to some of the Trustees—the desirability of an exhaustive series of examinations, chemic and bacteriologic, of the waters between Lake Michigan at Chicago and the Mississippi River at St. Louis, with the object of determining their condition and quality before the completion of the Drainage Channel, for comparison with their condition and quality under the dilution to be afforded by the Channel.

SCHEMATIC REPRESENTATION

of the

Self-Purification of the Waters


between the entrance of the


Main Drainage Channel at Chicago

and the

Mississippi River at Grafton

Based upon Tables of Bacteriologic and Chemic Examinations.
1900-1901





Such examinations have been made from time to time since 1869—the most important, in point of number, being those instituted by the late Dr. John H. Rauch in 1886-89, made by Professor J. H. Long and collated by Dr. F. W. Reilly. While these examinations demonstrated the effective decomposition of sewage matters, and consequent purification of the streams in a flow of less than fifty miles—that is, between Bridgeport and Channahon—they are defective, scientifically, in that they were confined solely to the chemic determinations, and in the present developed state of water analysis and examination, when water bacteriology is deemed essential, they would probably have little weight.

This defect was, in part, remedied by the work done under my direction as late as September, 1894, by Dr. Adolph Gehrmann, of the Department Laboratory, who made a special bacteriologic study of certain organisms in the waters between the South Branch of the Chicago River and a point a few miles below Peoria. But this work was confined to the bacterial examinations, and was not supplemented by simultaneous chemic analyses. As before said, both are essential.

I beg to submit that, in view of the hostile attitude of St. Louis and the threat to appeal to Federal authority to prohibit the use of the Drainage Channel, the Board should fortify itself with the evidence above indicated.

The outlay for this work need not be great, since the Board could undoubtedly arrange with the Illinois University to have the analyses and examinations made in its laboratories, without charge, and the findings would thus have the prestige, as evidence, of a state institution, and the weight of such authorities as Professor Palmer and his colleagues.

To further forestall criticism, I would suggest that duplicate samples should be offered to the St. Louis authorities for examination in some such institution as their Washington University, whose scientific standing is of the highest. It would be well, also, to secure the service of the laboratories of the Chicago University, in which Professor Jordan and others have already done a great amount of work on the general subject.

If this suggestion be acted upon, I should be glad, if desired, to take general charge and arrange with the representatives of the three institutions the details of the collection of samples, the points whence these shall be obtained, the technique to be employed in the examinations, the data to be included—meteorologic, topographic, demographic, etc.

My suggestion is based on the assumption that the opposition to the Drainage Channel has an honest animus; that St. Louis and the lower Illinois valley really believe that the effect of the Channel will be inimical to their health by affecting their water supplies. The way to remove these fears and allay these apprehensions is to ascertain and present the facts truthfully and impartially. I have no doubt as to the result—either as to the certainty of demonstrating a material improvement of the quality of the water by the Channel dilution, or as to the prompt acceptance of the situation by those now opposed when such demonstration has been made. I am, gentlemen,

(Signed) ARTHUR R. REYNOLDS. M, D., Commissioner of Health.


After due consideration the suggestions made in the above letter were adopted, and the following notification of appointment was received January 24, 1899:

SANITARY DISTRICT OF CHICAGO.

Chicago, January 23, 1899.

Arthur R. Reynolds, M. D., Commissioner of Health, City Hall, Chicago.

Dear Sir:—You are hereby notified that the Joint Committee on Federal Relations and Health and Public Order of the Sanitary District of Chicago has selected



MAP OF ILLINOIS RIVER

you, in your official capacity, to take the preliminary steps, under the direction of said Joint Committee, in regard to analyses of the water of the Chicago, the Desplaines, the Illinois and the Mississippi rivers. The method of procedure to be followed is as has been explained by you to Chairman Frank Wenter.

Yours truly,

(Signed) Joseph Haas, Clerk, Sanitary District of Chicago.

April 6 authority was given by the Joint Committee to purchase material and to provide for the payment and expenses of the experts of the Illinois and Chicago Universities and also for like expenditures in the Municipal Laboratory.

January 19, of the following year, in a letter addressed to Hon. Frank Wenter, Chairman of the Joint Committee, the plan, which had already been put into operation for conducting the work was briefly outlined, and copies of the correspondence showing that St. Louis had failed to join in the proposed investigation or to recognize it in any manner were submitted:

Chicago, January 19, 1900.

Hon. Frank Wenter, Chairman,

Joint Committee on Federal Relations, Health and Public Order, Board of Trustees of the Sanitary District of Chicago.

Dear Sir:—Under date of January 23, 1899, I was notified by your Committee that I had been selected to undertake the direction of a complete and comprehensive analysis of the waters flowing between Chicago and the Mississippi River.

On the following day, January 24th, I addressed a letter to Andrew Sloan Draper, LL. D., President, University of Illinois, Champaign, Ill.; Winfield S. Chaplin, LL. D., Chancellor, Washington University, St. Louis, Mo., and William R. Harper, D. D., President, Chicago University, Chicago, Ill. It was proposed to them to collect samples, in triplicate, to be sent to the laboratories of each institution.

Dr. Draper accepted promptly, and appointed Dr. Arthur W. Palmer, Professor of Chemistry, to represent the University. Dr. Harper, in accepting for the University of Chicago, named Edwin O. Jordan, Professor of Bacteriology in that institution, as its representative.

Dr. Chaplin, of Washington University, commended the plan, but informed me that the Washington University, an an institution, could not take part in the conference or the work.

On February 6, 1899, I personally visited St. Louis, called upon the Hon. Henry Ziegenheim, Mayor, and verbally requested that the city of St. Louis take the place of Washington University in the investigation. I was referred by him to Mr. W. L. Holman, Water Commissioner. Mr. Holman was interested in the plan, and deemed it fair and wise.

On February 9th the following letter was addressed to Mayor Ziegenheim:

Hon. Henry Ziegenheim,

Mayor, City of St. Louis.

Chicago, February 9, 1899.

My Dear Sir:—In pursuance of our agreement last Monday, I called at once upon Commissioner Holman, with reference to the chemical and bacterial examination of the streams between Chicago and St. Louis which may be affected by the operation of the Main Channel and Waterway of the Chicago Sanitary District.

While our conference was in every way satisfactory and resulted in a mutual understanding, I think it advisable to state definitely, for your own information, the object and scope of the projected examination.

The project had its direct origin in a communication addressed by me, November 28, 1898, to the Board of Trustees of the Sanitary District of Chicago, in which attention was called to the growing fear of St. Louis and certain towns in the lower Illinois Valley that the operation of the Main Channel of the District would prove injurious to the health of those communities, by increasing the pollution of their water supplies, and that this fear was at the bottom of the opposition to the Channel.

Concerning this I wrote:

"My suggestion is based on the assumption that the opposition to the Drainage Channel has an honest animus; that St. Louis and the lower Illinois Valley really believe that the effect of the Channel will be inimical to their health by affecting their water supplies. The way to remove these fears and allay these apprehensions is to ascertain and present the facts truthfully and impartially. I have no doubt as to the result—either as to the certainty of demonstrating a material improvement of the quality of the water by the Channel dilution, or as to the prompt acceptance of the situation by those now opposed when the demonstration has been made."

To ascertain and present the facts truthfully and impartially, it is proposed to secure a chemic and bacteriologic examination of the waters, collected at various points between Chicago and St. Louis, beginning at the earliest possible moment, and to be continued for a corresponding period after the Channel is opened, in order to compare the sanitary quality of these waters before and after the dilution which the Channel will afford.

That the facts shall be ascertained and presented "truthfully and impartially," it was originally proposed that the examinations should be made by three scientific institutions of high reputation, to-wit, the Washington University of St. Louis, the University of Chicago, and the Illinois State University. Triplicate samples of the waters are to be collected at the various points selected, and one set sent to each institution for examination.

In conference with Mr. Holman, it was suggested that the examinations in St. Louis be made in the laboratory of the Water Commissioner and under his supervision. I see no objection to this modification of the original plan; it will probably result in greater confidence by St. Louis in the results obtained.

The two Illinois institutions have signified their readiness to undertake the work as outlined, and I presume it only needs your approval of Commissioner Holman's suggestion in order that we may have an early meeting of the representatives of the three examining bodies to arrange details.

Awaiting your action, which I trust may not be unduly delayed, and thanking you for the courtesies already extended me, I am, my dear sir," etc.

No response was made to this letter. Mayor Ziegenheim was again addressed on March 9th, and again urged to join in the investigation, to which no answer has yet been received.

The laboratory of this Department was then pressed into service, and the work organized and conducted as indicated in the accompanying report of Dr. Gehrmann, Director of the Municipal Laboratory.

(Signed) ARTHUR R. REYNOLDS, M. D.,
Director of Streams Examination, for
the Sanitary District of Chicago.

As will be seen by the reports of Dr. Gehrmann and Professors Palmer, Jordan and Burrell, the investigation has been conducted on a plan of greater magnitude than that of any similar inquiry of which I have any knowledge. The scientific gentlemen connected therewith have been untiring in their labors and inspired wholly by a desire to learn the truth. The task imposed upon them was one that appealed to their

interest as scientists and to their pride and loyalty as public-spirited citizens. They have performed an important work faithfully and well, and it is believed that the reward and recognition which are theirs by right will be freely and gladly given, now that the result of their labors and research are to be published in permanent form.

In this connection, too, a word should be said as to the scientific value of the facts and data collated in the present volume.

While sanitary authorities now agree, in the main, on the self-purification of water under certain conditions, there have been those who have taken the negative side of the proposition. The results herein presented and the deductions to be made from the laboratory records of the University of Illinois, the University of Chicago, and the Municipal Laboratory, may be said to be conclusive as establishing the affirmative side of the question. In other words, it is now clearly proven that running water, if not too heavily charged with organic pollution, will purify itself through the natural bio-chemic processes, of which bacterial action and insolation are the most important. Indeed, it is now conceded that unless this were true there would be no such thing as pure water in streams affected by human habitancy.

It only remains for me to acknowledge the valuable services of Dr. John R. Neely, until recently connected with the Department of Health, and to Mr. E. R. Pritchard, secretary of the Department, for work done in the final preparation of the present volume for publication.

The sewage disposal problem of Chicago has been satisfactorily and scientifically solved. Very respectfully,

ARTHUR R. REYNOLDS, M. D.

REPORT OF THE LABORATORY OF THE CITY OF CHICAGO.

BY

PROF. ADOLPH GEHRMANN, M. D.

Arthur R. Reynolds, M. D., Director Streams Investigation, Sanitary District of Chicago.

Dear Sir:—Following your instructions, several conferences between Professor Arthur W. Palmer and Professor T. J. Burrill, of the University of Illinois, and Professor Edwin O. Jordan, of the University of Chicago, and myself, were held during January and February, 1899. These conferences, supplemented by considerable correspondence, resulted in decisions upon the following points connected with the investigation:

First—Localities from which samples would be collected.

Second—That all matters relating to the exact points in the rivers from which the samples would be taken and the selection of collectors would be left to the mutual agreement of Professors Palmer and Jordan.

Third—That arrangements for transportation of samples and equipment of collectors with outfits would be conducted by the Health Department Laboratory.

Fourth—A decision as to the outfits to be used by the collectors.

Fifth—The formulation of a mutually agreed summary of procedure for the methods of analyses to be used by the different laboratories. These methods appear in detail elsewhere.

As a result of these conferences the following communication was sent to you:

"March 28, 1899.

"Arthur R. Reynolds, M. D., Commissioner of Health.

"Dear Sir:—As the present outline of the plan for taking samples in the Illinois and Mississippi rivers contemplates the collection of about forty samples weekly, there will, therefore, be collected during the year 2,080 samples, to be delivered to each of the three laboratories. These samples are to be in duplicate for bacteriological and chemical purposes. They are to be collected in glass-stoppered glass bottles, and properly protected by covers over the stoppers, and sealed. The transportation is to be accomplished by sending wooden cases of such size as to hold the bottles. The samples for bacteriological analysis will be packed in a larger receptacle filled with ice during the warmer periods of the summer.

"The points at which samples are to be taken, as outlined in the present program, are as follows:

Sam		
I.	Bridgeport	Illinois and Michigan Canal.
2.	Lockport	Illinois and Michigan Canal.
3.	Toliet	Illinois and Michigan Canal, upper basin.
4.	Toliet	Illinois and Michigan Canal, middle basin.
5.	Channahon	
6.	Wilmington	Kankakee River.
7.	Morris	Illinois River.
8.	Ottawa	Fox River.
9.	La Salle River	Big Vermilion.
10.	La Salle	Little Vermilion.
II.	La Salle	Illinois River.
12.	Henry	Illinois River.
13.	Peoria	Illinois River.
14.	Pekin	Illinois River.
15.	Copperas Creek	Illinois River.
16.	Havana	Illinois River.
17.	Chandlerville	
18.	Beardstown	Illinois River.
19.	Pearl, C. & A. R. R	
20.	Grafton	Illinois River.
2I.	Grafton	Mississippi, above Illinois.
22.	Grafton	Mississippi, below Illinois.
23.	Alton	Mississippi, above city.
24.	Alton	
25.	Alton	
26.	Alton	
27.	Alton	
28.	Alton	Mississippi, below city.
29.	Bellefontaine	
30.	St. Louis	
31.	St. Louis	
32.	St. Louis	
33.	St. Louis	
34.	St. Louis	
35.	St. Louis	
36.	St. Louis	
37.	St. Louis	
38.	St. Louis	
39.	St. Louis	
40.	St. Louis	City water supply.

"I have obtained a rough estimate of the outfits that will be required for the collection of the samples, basing the general requirement upon the 40 samples a week to each laboratory, with sufficient cases and extra bottles to prevent any possible delay in sending the samples, or in case accidents should cause breakage. Two hundred complete outfits have been decided upon as the proper number to consider in the first estimate.

"Respectfully,

"ADOLPH GEHRMANN, M. D.,
"Director of Laboratory."

Supplies were purchased and the required number of outfits for the collectors was prepared. After some difficulty satisfactory arrangements for transportation of the shipping cases were made with the express companies. Upon advice from Professor Palmer and Professor Jordan, the collectors were supplied with outfits and instructions. The following general letter of instructions was sent to each man thus selected:

"Chicago, May 19, 1899.

"Mr. ———, Collector at ———.

"Dear Sir:—According to arrangements that have been made by Professors Jordan and Palmer, you are to make collection of samples of water for the Chicago Sanitary District.

"These samples are to be collected — of each week from the locality designated by them. From each spot from which water is obtained three separate samples must be collected; as each sample is represented by a large bottle and a small one, six bottles must be filled with water.

"The shipping cases and outfits have been sent to you. The tags, seals and numbers you will find in one of the three cans belonging to each sample. There is one press for the seals; this bears a number. This is your number as collector.

"The three samples are to be packed in the shipping cases and sent to the three addresses indicated upon the shipping envelopes. The shipping envelope is to be fastened to the neck of the large bottle by the wire seal; the small bottles do not require shipping envelopes.

"The samples are to be numbered with the tin tags bearing numbers. Use the numbers in regular order. As there are six bottles from each locality, there are six tags numbered alike, and one of these is to be fastened by the seal to each bottle. In putting on the seals pass the wire in and out through the holes in the cloth cover of the bottle.

"Read carefully the instructions on the cards that go in the shipping envelopes.

"The bills for this service are to be rendered monthly and must be made out in duplicate as follows:

"Chicago Sanitary District, "Arthur R. Reynolds, M. D.

"Director of Streams Examination, "411 City Hall, Chicago.

"The bill must state the dates and places of taking samples. Should you desire additional information upon any point or wish to give up the collection of samples at any time, please write to me.

"Very respectfully,

"(Signed.) ARTHUR R. REYNOLDS, M. D., "Director of Streams Examination."

Subsequent to this another circular letter was directed to each station calling special attention to certain points that apparently were not clearly understood. By this means and also through considerable personal

correspondence, the collectors came to attend to their duties with accuracy and expedition. Much credit is due these collectors for the success of the investigation.

Supplemental instructions sent to collectors:

"Chicago, May 31, 1899.

"Mr. ———. Collector at ———.

"Dear Sir:—It is desired that you proceed in the following manner regarding the packing of the small sample bottles in the cans: When the bottle is placed in the small can the lid must be pressed firmly in place so as to prevent water getting into it. The smaller can is then to be packed in ice in the large can. This can must be completely filled with ice. In the shipping case the packing paper goes around the can, and excelsior is to be stuffed into the corners and packed over the top of the can, filling up to the top of the shipping case. The ice must be kept from melting as well as possible.

"We are about to send you one complete outfit for sample to use in case of accident to any of the regular bottles or cans that are in use. This you will keep until used.

"We are also sending you the thermometers for taking temperature." These you will find in one of the shipping cases returned to you.

"Yours very truly,

In June of 1899 this part of the service was moving smoothly, and it was now considered that the men were sufficiently experienced to begin the official series of samples. At this time the complete list of collectors was as follows:

MONDAY.

I.	Ill. and Mich. CanalBridgeport, 500 feet west of pumping station.
2.	Ill. and Mich. CanalLockport.
3.	Desplaines RiverLockport.
1	Ill and Mich Canal (north) Ioliet F bank of N of Jackson street

- 5. Ill. and Mich. Canal (south). Joliet, E. bank of R. L. Ry bridge.
- 6. Kankakee River Wilmington.
- 8. Fox RiverOttawa, above canal viaduct.
- 9. Illinois RiverOttawa. Ill. and Mich. Canal..... Ottawa.

- 10. Big Vermilion River La Salle, at bridge.
- 12. Illinois RiverLa Salle, at wagon bridge.
- 14. Illinois River Peoria, Averyville bridge, at narrows No. 11.
- 16. Illinois River Pekin.
- Lake.
- 18. Sangamon River Chandlerville.

WEDNESDAY.

		WEDNESDA1,
19. 20. 21. 22. 23. 24. 25. 26. 27.	Mississippi River	Kampsville, above dam. Grafton, 2 miles above, in mouth of Illinois River. Grafton, above mouth of Illinois. Alton, Ill., 100 ft from E. bank Alton, Ill., E. of middle Alton, Ill., mid-stream Opposite new pumping
		THURSDAY.
28. 29. 30. 31. 32.	Mississippi River	Mitchell, inlet tower Mitchell, 400 ft. from Mo. shore W. Alton, Ill., at C., B. & Q. Ry bridge
33. 34. 35. 36. 37. 38.	Mississippi River	Jefferson Barracks, E. middle of stream. Jefferson Barracks, mid-stream. Jefferson Barracks, W. of middle. Jefferson Barracks, W. bank.
	COLLEC	CTION OF SAMPLES.
I. 2.	Dridgeport, I sample	Canal and Desplaines River, collector P. W.
3.		O'Brien, Lockport. Both from Canal, Ray Hurd; collector R. P. Elliott, Agt. U. S. Exp.
5.	Morris, 1 sample	Illinois River, J. W. Miller, Box 712; collector James Mack, Morris.
6.	Ottawa, 2 samples	Fox River and Canal, collector Dennis Foley, Ottawa, Ill.
7.	La Salle, 3 samples	Big Vermilion. Little Vermilion and Illinois Rivers, collector Dr. Wm. Fraser.
8.	Henry, I sample	Illinois River, collector Jas. McCune, Henry, Ill.
9.	Peoria, Averyville, 1 sample	Illinois River, collector D. J. Forbes, Peoria waterworks, Peoria.
10.		(Illinois River, collector D. H. Jansen, Co. Surveyor.
II.	Pekin, Ill., 1 sample	(
12. 13.	Havana, I sample	Illinois River, collector H. J. Heberling. Sangamon River, collector. E. O. Spink.
14.	Beardstown, 1 sample	Illinois River, collector J. A. Carney, C., B. &
15.	Kampsville, 1 sample	Illinois River, collector C. V. Brainard, Asst. U. S. Eng.
16.	Grafton, 3 samples	Illinois, Mississippi and Missouri Rivers, eollector B. F. Robinson Grafton.
17.	Alton, Ill., 5 samples	All from Mississippi River, collector Geo. Brooks, Alton.
18.	Mitchell, 4 samples	Mississippi River, collector Henry Atkins, Mitchell.

- 19. West Alton, Mo., 1 sample... Missouri River, collector Jas. Mathews, West Alton.
- 20. St. Louis, Mo., I sample.... Mississippi River and St. Louis tap, collector Aug. Johnson, 407 Espensheid street.

These general arrangements remained in force during the period of the investigation.

ANALYSES OF LAKE MICHIGAN WATER IN THE HEALTH DEPARTMENT LABORATORY.

A systematic analysis of the city water supply from Lake Michigan has been made since January, 1894. During 1894 samples were collected weekly, but from January, 1895, daily samples, exclusive of Sundays and holidays, have been examined. Full details of these analyses have been published from time to time in the reports of the Department of Health. Previous to the opening of the main drainage channel in January, 1900, a very close relation between rainfall and the sanitary quality as shown by analysis was demonstrated. Since that date this close relationship has failed to appear on numerous occasions, and although the evidence of contamination appears at irregular intervals, there is in general much improvement, especially as shown in the analyses of samples from the intakes farthest from shore. As a large area south of Thirty-fifth street and east of Clark street, and another north of Fullerton avenue, still drain into Lake Michigan, the source of pollution is still apparent, and not until the sewers in these areas are connected with the intercepting sewer system can much improvement over present conditions be expected.

The results of the analyses made in the laboratory of the department, which appear in the tabulations, give evidence of the purification taking place in the waters of the main drainage channel and in the Illinois River. On all of the main points these are in entire agreement with the results of our colleagues in the investigation. The slight variations in individual analyses are due to differences in time between collection and analysis and variations in temperature of the samples. The series of analyses made by us does not cover as long a period as those of our associates; this was occasioned by the increase in the regular work of the laboratory and also by changes in our working force. For the same reasons special summaries of the results were not prepared. Indeed, as these would be simply repetitions of those that are presented, they would not, in the least, change the outcome of the investigation.

CONDITIONS IN LAKE MICHIGAN.

The water front of the city is 21 miles in extent. Farther north are numerous towns, the sewage of which drains directly into the lake, while to the south the Calumet River, highly polluted with house drainage and manufacturing refuse, is an important source of pollution. However, all polluted water reaching the intakes is highly diluted and subject

to more or less digestion as it lays along near shore and purification by mixture with fresh lake water in passing into the lake two or four miles.

WHAT MAY BE CONSIDERED AS NORMAL LAKE WATER.

To determine this, samples have been collected at intervals at a distance of ten to twelve miles from shore and a varying depth from ten feet below the surface to a depth of fifty feet.

Analysis of Lake Michigan water taken twelve miles east of the mouth of Chicago River:

Total solids130.	00
Loss on ignition	.00
Fixed mineral solids106.	.00
Chlorine 5	.50
Free ammonia no	ne
Albuminoid ammonia	
Nitrogen as nitrates no	
Nitrogen as nitrites no	ne
	64
Oxygen consumed in 10 min 1.6	00
Bacteria per cc520	
Pathologic bacteria absent.	

(In parts per million.)

October 23, 1896—ten feet below the surface.

Day clear; wind west; wind movement for previous twenty-four hours. 315 miles; precipitation, none; mean temperature, 42° F.

This may be taken as an average analysis of the lake water under most favorable conditions. However, it may be said in regard to the number of bacteria per cubic centimeter that this will be lower in most instances. The count as usually found during the best periods is less than 100 per cubic centimeter.

The extent to which the area of contamination will pass into the lake has been shown by similar series of samples, and from these it has been found to extend ten miles from shore.

Samples collected ten miles from shore upon the days as noted showed presence of B. coli communis or an allied variety.

October 25, 1900, B. coli communis. November 23, 1900, colon group bacilli. December 6, 1900, B. coli communis. April 12, 1901, colon group bacilli.

Samples taken July 12, August 2, August 9, September 6, September 20, 1900, and on April 3 and June 26, 1901, at about the same place, 10 miles east from the mouth of the Chicago River, showed absence of the colon-typhoid group of bacteria.

The daily examination has included regular samples from the intake tunnels as follows:

Fourteenth street, four-mile tunnel off Peck court. Chicago avenue, two-mile tunnel off Chicago avenue. Hyde Park, two-mile tunnel off Sixty-seventh street. Lake View, two-mile tunnel off Montrose boulevard.

Since January 1st regular samples from Carter H. Harrison Crib, three miles off Oak street, and from Rogers Park water supply have been examined. These, however, are not important in this connection, as there are no earlier analyses with which to make comparisons.

In order to show the variation taking place in the quality of the water, samples have been collected every hour during twenty-four-hour periods. The following is a report that was rendered upon such a special series of samples:

"Chicago, Aug. 23, 1900.

"Dr. F. W. Reilly, Assistant Commissioner of Health.

"Dear Sir:—In accordance with your request a series of water samples was collected from the tap in the Fourteenth street pumping station every hour for a period of ten hours, beginning August 20 at 7 p. m. The result of the chemical examinations of these samples is as follows:

August 20—	Free	/ lbuminoid	Chlorine.	Oxygen consumed in
Sample No. Time-	ammonia.	ammonia.		min. at 100° C
8733— 7 p. m	.000	.005	.60	.260
8734— 8 p. m	.001	.005	.60	.260
8735— 9 p. m	.000	.008	.60	.230
8736—10 p. m	.000	.008	.60	.230
8737—11 p. m	.002	.008	.60	.300
8738—12 p. m	001	.008	.60	.320
August 21—				
8739— 1 a. m	. 000.	.009	.60	.250
8740— 2 a. m	001	.006	.60	.250
8741— 3 a. m	000.	.008	.60	.210
8742— 4 a. m		.007	.60	.240
	In parts p	er 100,000.		

August 20—	No. Minus Ourselles	33-43	0- 14
No. Time.	No. Micro-Organisms per cubic centimeter.	Pathogenic organisms found.	Sanitary quality.
8733— 7 p. m	300	None.	Good.
8734— 8 p. m	60	None.	Good.
8735— 9 p. m	125	None.	Good.
8736—10 p. m	115	None.	Good.
8737—11 p. m		None.	Good.
8738—12 p. m	45	None.	Good.
August 21—			
8739— 1 a. m	200	None.	Good.
8740— 2 a. m		None.	Good.
8741— 3 a. m	60	None.	Good.

"The result of this examination shows that during this period there was nothing more than slight changes in the character of the water. The sanitary quality of all the samples would be rated as 'good.'

"The presence of free ammonia in four of the samples cannot be taken as an indication of serious pollution; although the normal lake water seldom contains more than traces of free ammonia.

"The average amount of albuminoid ammonia as determined by our analyses of the samples taken as far as ten miles into the lake shows that it ranges between .007 and .009 per 100,000. As far as the albumin-

oid ammonia figures are concerned in this series, the indication would be plain that unpolluted lake water was being taken from the four-mile crib.

Very respectfully,

"ADOLPH GEHRMANN, M. D.,
"Director of Laboratory."

TABLES SHOWING VARIATIONS IN SANITARY QUALITY OF LAKE WATER.

The following tabulations are summaries of all the analyses of samples collected at the pumping stations during 1896 to 1901, inclusive, and present the average for each station for the years named:

			In Parts per 100,000			,		
Years	s. Stations	Free ammonia.	Albuminoid ammonla.		Oxygen consumed.		Per cent, times bacteria found.	
1896	14th Street Chicago Avenue. Hyde Park Lake View	00181	.00701 .00765 .00671 .00881	.317 .323 .325 .328	.157 .172 .158 .184	3,841 7,008 6,015 7,511	4.21 Patho 7.14 senic 7.14 for 6.92Anlmal	
1897	14th Street Chicago Avenue Hyde Park Lake View	0.0012 0.0002	.0060 .0065 .0062 .0065	.280 .327 .322 .309	.181 .229 .198 .193	6,840 10,516 8,405 10,095	16.5 24.5 20.3 20.4 B Coli	
1898	14th Street Chicago Avenue Hyde Park Lake View	$0010 \\ 0005$.0035 .0038 .0039 .0033	.325 .333 .335 .328	.177 .203 .207 .185	996 2,032 1,380 2,206	13.0 nis. 19.7 19.1 12.8	
1899	14th Street Chicago Avenue Hyde Park Lake View	0011	.0047 .0048 .0047 .0043	.337 .331 .335 .337	.147 .151 .167 .166	645 1,323 786 400	30.3 48.2 44.4 35.9	
1900	14th Street Chicago Avenue Hyde Park Lake View	0.0024 0.0019	.0124 .0123 .0117 .0119	.318 .316 .317 .317	.178 .207 .181 .183	227 518 409 339	23.2 53.4 Typho- 48.4 Colon 32.3 Group.	
1901	14th Street Chicago Avenue Hyde Park Lake View	0.0017 0.0014	.0094 .0099 .0091 .0091	.312 .319 .318 .318	.164 .191 .163 .165	1,161 1,558 1,673 1,570	23.0 38.1 35.0 26.5	

During the first three months of 1902 a marked improvement in quality is manifest, as shown by percentages of samples in which the pathogenic bacteria were found during this time.

1902.	January.	February.	March.	Average.
14th Street Station	. 11.5	0.0 54.5	0.0	3.8 33.6
Hyde Park Station Lake View Station	. 30.8	13.6	4.0	16.1 2.6

As each step in the general scheme for pure water has been made, a degree of improvement corresponding to it has been noted. Each time that serious contamination has been noted the evidence for it has been obvious, and as soon as the fault was corrected, the quality of the water returned rapidly to the status before the special period of pollution.

Respectfully submitted,

ADOLPH GEHRMANN, M. D., Director of Laboratory.

REPORT OF THE UNIVERSITY OF CHICAGO.

BY

PROF. E. O. JORDAN, Ph. D.

ARTHUR R. REYNOLDS, M. D., Director of Streams Examination.

Dear Sir:—I beg to submit herewith the final and complete report upon the chemical and bacterial analyses conducted under my direction in behalf of the Sanitary District of Chicago. These analyses cover a period extending from May 1, 1899, to July 1, 1900, and embrace about two thousand chemical and bacterial determinations. Regular weekly analyses have been made of water samples collected from the Illinois and Michigan Canal, from various parts of the Desplaines and Illinois rivers and from the principal tributaries of the Illinois River; regular examinations have also been made of samples from the Mississippi River at several points, from the Missouri River and from the St. Louis city water supply. The reasons governing our choice of the points for collection are stated in detail elsewhere (pp. 47-54). With two exceptions the collecting stations have been visited conjointly by Professor Palmer and myself, and all of the more important stations have also been visited separately by us both.

Throughout the investigation I have been assisted on the chemical side by Mr. F. L. Stevens, Ph. D. (Chicago), and on the bacterial side by Mr. E. E. Irons, S B. The continued faithfulness, zeal and high efficiency with which they have carried on their work deserve especial mention. I have also been assisted in special ways at various times by Messrs. W. G. Sackett, W. L. Sayer and C. B. Davis.

DESCRIPTION OF METHODS EMPLOYED AND RESULTS OBTAINED IN THE REGULAR ROUTINE EXAMINATION.

(a) Method of Collection. Explicit instructions were given to all collectors concerning the methods to be employed, and, in addition, printed directions were sent out weekly with each collecting bottle. The bacterial samples were collected in wide-mouth four-ounce glass bottles, which were sterilized in the laboratory, enclosed in a tight-fitting metal case, which was itself placed in a large packing canister, ten inches deep and six and one-half inches in diameter, and the whole, together with the

corresponding bottle for the chemical samples, fitted into a wooden box, one such outfit being shipped to the collector each week. Great pains were taken to insure that the sample was obtained with due bacterial precautions, and in order to emphasize the instructions given to the collector personal visits, involving careful supervision and recurrent demonstration, were made frequently to all the more important points of collection. The water sample was always taken at a point about eight inches below the surface and in midstream, except at Joliet and save where cross-sections were taken. The bottle of water was placed in its covered case as soon as collected, and this case was then packed in the canister and completely surrounded with ice. The hour of collection was always timed to permit of as speedy shipment as possible.

The chemical samples were collected in glass-stoppered bottles of one-gallon capacity. These were cleaned with chromic acid cleaning mixture each time before being sent out from the laboratory, thoroughly rinsed with ammonia-free water and drained. The stoppers were secured in place by tying tightly over them a piece of rubber cloth. After filling the bottle the cloth cap was fastened in place with wire which was sealed with stamped metal.

(b) Methods of Physical and Chemical Analysis. A serial number was assigned to each sample immediately upon its receipt in the laboratory, and this number was also placed upon the collector's certificate. The seal was then broken and the cover of the stopper removed. Any dust adhering was carefully wiped away, and a few cubic centimeters of the sample were poured out to wash the lip of the bottle. About 1,500 c.c. of the water were then filtered through a double filter (Munchfilter, No. 0), which had been carefully washed with a half liter or more of pure water. The determinations of the "nitrites" and "nitrates" have been made first, followed by the "ammonias" and oxygen consumed. The "date of collection" recorded in the table is taken from the collector's certificate that accompanies each sample; the "date of examination" refers to the date of making the nitrate and nitrite determinations and the ammonia analyses. The determinations of the residue and chlorine have been completed as soon as practicable. The results are all expressed in parts per million.

The loss on ignition of the residue on evaporation was omitted, since analysts are now generally agreed that the information furnished by this determination is not valuable enough to compensate for the labor involved.

Turbidity and Sediment.—The turbidity and sediment have been determined in the laboratory merely by ocular inspection and are expressed by the conventional adjectives "slight," "distinct," "decided" or "muddy," as regards turbidity: "very little," "little," "considerable" or "much," to indicate the sediment.

In the absence of any recognized standard at the time this investigation was undertaken this seemed to be the best method available, although it proved far from satisfactory.

Color.—For the sake of uniformity with previous examinations of the waters dealt with, the color of the filtered water has been ascertained by

comparison with nessler standards, in which .o1 m. g. of N. as ammonia is taken as a unit of color.

Odor.—The odor is recorded either as "gassy" or as "none," with no attempt to classify further.

Residue.—The total and dissolved residues on evaporation have been determined by evaporating 100 c.c. of the unfiltered and filtered waters, respectively, in a weighed platinum dish over the steam bath at about 100° C. The dish with the residue is then heated to a temperature of 170°-180° for one hour and the weight taken, after cooling in a desiccator over calcium chloride.

Chlorine.—The ordinary process of titration with silver nitrate has been followed (Report of Massachusetts State Board of Health, 1896, Purification of Sewage and Water, p. 723), 50 c.c. of water being used. Potassium chromate, 1 c.c. of a 5 per cent solution, is used to indicate the end of the reaction. When the amount of chlorine was presumed to be low (under fifteen parts per million) a measured quantity, usually 250 c.c. of the sample, was concentrated to less than 50 c.c. before titrating.

Oxygen Consumed.—One hundred cubic centimeters of the sample were measured into an Erlenmeyer flask of about 300 c.c. capacity and acidulated by the addition of 5 c.c. of concentrated sulphuric acid. Ten cubic centimeters of standard potassium permanganate solution were then added and the sample allowed to remain in boiling water for thirty minutes, more permanganate being added, if needed, to prevent complete loss of color. Upon removal from the bath, ten cubic centimeters of standard oxalic acid solution were added and the sample titrated back to a just perceptible pink color with permanganate. Great care was taken to insure that the process was carried out in the same way at each determination, and to this end quantity, time and temperature were made strictly uniform.

Nitrogen as Free Ammonia.—A half liter of the water, or less if the nitrogen content was very high, was rendered alkaline by the addition of 5 c.c. of 20 per cent sodium carbonate solution and distilled in glass flasks through a block tin tube condenser; the rate of distillation was such that 50 c.c. were collected in about ten minutes. The distillate was caught in three 50-c.c. nessler tubes and the tubes nesslerized separately unless the ammonia was high, in which case 200 c.c. of the distillate were caught in a graduated flask and, after thorough mixing, an appropriate aliquot portion was taken in a tube for nesslerization. Before each analysis the flasks and condenser were steamed until free from ammonia.

Nitrogen as Albuminoid Ammonia.—To the residue from the last operation 50 c.c. of the ordinary alkaline permanganate solution were added and the distillation continued, the distillate being collected either in tubes or flasks, according to the quantity of ammonia present. A small amount of rather coarse, thoroughly burned pumice has been found very useful in this process as a preventive of bumping, which is otherwise extremely troublesome.

GAAFION Edwin O. Jordan . MAMPSVILLE HAVANA BEARDSTOWN CHLORINE - PARTS DER MILLION , IN ILLINOIS RIVER. 135 150 165 180 WESLEY CITY PEXIN AVERYVILLE MENRY OTTAWA MORRIS BRIDGEPORT

Nesslerization.—The determination of the amount of ammonia in the distillate was made in the usual way by comparison with standard ammonia tubes. This process is greatly facilitated by the use of a nesslerizing cabinet, which is constructed, in a somewhat modified form, upon the same principle as one which has been in use for some years in Professor Palmer's laboratory at the University of Illinois. Eighteen standards are used, ranging in value from .004 to .13 parts per million. Fresh standards are made for each nesslerization, and the nessler solution is added to them and simultaneously to the distillates, which have been cooled to the room temperature. The readings are made about thirty minutes later.

Nitrogen as Nitrates.—The nitrates have been determined by the aluminum reduction method, which has not proven very satisfactory for many of the waters dealt with, but which has been adhered to for the sake of uniformity. Two and one half cubic centimeters of 33 per cent nitrogenfree sodium hydroxide solution were added to 100 c.c. of the sample and the mixture boiled rapidly to a volume less than 50 c.c., in order to drive off the free ammonia. After cooling, the residue was poured into a tall tube, its volume made up to 50 c.c., and a strip of aluminum added. After about twelve hours, when reduction is complete, the aluminum is removed and the tube allowed to stand till the supernatant fluid is clear. A measured portion, usually 10 c.c. or 20 c.c., is then pipetted into a nessler tube, diluted to the mark and nesslerized directly.

Nitrogen as Nitrites.—A modification of the Griess method was employed. Two stock solutions were prepared:

- I. A saturated solution of sulfanilic acid in 5 per cent HCl.
- 2. Eight gr. of napththylamine, 8 c.c. of HCl, made to 1,000 c.c.

These solutions were mixed in quantities of about 100 c.c. at a time, and the mixed solution was employed as the test solution, thus avoiding unnecessary pipetting. One cubic centimeter of the mixture was added to each tube.

About 50 c.c. of the sample were clarified by the use of one cubic centimeter each of decinormal alum and soda solutions and filtration through a washed filter paper. Then 50 c.c., or less if the nitrite was high, were placed in a nessler tube and compared with standards, the reading being taken about forty-five minutes after the addition of the test solution.

The standards were made with sodium nitrite prepared from silver nitrite.

(c) Methods of Bacterial Analysis. Dilution.—Preliminary experiments showed that when water containing a large number of bacteria was mixed directly with the nutrient medium a considerable proportion of the organisms present failed to develop visible colonies. Consequently, all waters were diluted with a known quantity of sterile water in small flasks, so that from I c.c. of the dilution used approximately one hundred colonies developed on a plate.*

^{*} See Notes on Bacterial Water Analysis. Jordan and Irons. Reports and Papers of American Public Health Association, 1899, XXV, p. 564.

Plating.—Nutrient agar, prepared as described below, was used as the standard medium throughout the work. Tubes containing 6-8 c.c. were heated in the water bath at 100° C. for about ten minutes and then placed in a water bath at 40° C. The tubes, after being allowed to cool to 40° C., were inoculated with 1 c.c. of a suitable dilution of the water under examination, and quickly poured into sterile Petri dishes, care being taken that as little as possible of the medium remained in the tubes. The plates were incubated in a dark culture room, the air of which was kept moist, at 20°-23° C. for eight days, and finally counted with the aid of a small hand lens. During the early part of the work an incubation period of ten days was allowed, but the shorter period of eight days was found to give as constant results, and was finally adopted.

Media.—Nutrient agar was made, with some slight modifications, after the standard method. Lean beef (1,200 gr.), chopped fine, was immersed in twice its weight of distilled water in the cold for twenty-four hours and the resulting infusion strained through cloth, yielding about 2,400 c.c. of filtrate. Of this filtrate, 2,000 c.c. were heated and the insoluble albumens precipitated. On filtration, about 1,600 to 1,800 c.c. of clear filtrate were obtained. The remaining 400 c.c. of the original infusion was added to 1,600 c.c. of this filtrate, together with 1 per cent Witte's peptone and 1½ per cent of agar-agar, previously dissolved by boiling for ten minutes in one liter of distilled water. The whole was boiled over a free flame for half an hour and then neutralized.

The total precipitation of albumens in the 2,000 c.c. of the original infusion was found advantageous, for the reason that, when all the albumen was allowed to remain, it interfered greatly with the recognition of the end point in titration, and also hindered filtration. Total precipitation of part of the infusion was found preferable to partial precipitation of the whole, because the flocculent precipitate which first comes down is much better adapted to final clearing of the agar than is the later, less flocculent matter. For these reasons the albumen in one-fifth of the infusion was regularly used as a final coagulant for the whole.

After the first neutralization, the agar was boiled for one and one-half hours over a free flame, the loss of water by evaporation, measured by weight, being made up from time to time with distilled water. After the second boiling the agar was neutralized, boiled for five minutes and filtered through paper in an ordinary glass funnel. The process usually yielded about 1,600 c.c. of clear agar, which filtered in from ten to twenty minutes, the last few cubic centimeters taking slightly longer.

The finished agar was made up to standard plating reaction (10 c.c. normal acid per liter), with normal HCl, and was then immediately tubed and sterilized in the autoclave at 120° C, for five minutes.

Sodium hydrate $\binom{n}{20}$ was used in titration with phenolphthalein (1 gr. in 1 l. of 50 per cent alc.) as an indicator. A permanent faint rose tint was regarded as the end point. In titration, 5 c.c. of the medium were boiled with 50 c.c. of distilled water in a white evaporating dish for three minutes over a free flame, and titrated while hot. All titrations

were made in duplicate. Normal sodium hydrate was used in neutralization.

- (d) See tables 81-158 (Appendix).
- (e) Methods and Results of the Examination for the Presence of B. coli communis. It is well known that fresh sewage always contains large numbers of the common colon bacillus. It is also true that when a water source is polluted with any considerable quantity of fresh sewage it is usually possible to demonstrate the presence of the colon bacillus in such water. Upon these familiar facts have been based various methods and conclusions of greater or less value to public hygiene.

The particular method of gauging the so-called self-purification of a stream by the relative abundance of B. coli communis at different points is not a new one, but, so far as the writer is aware, it has not been often applied on a large scale.

Theobald Smith's ingenious method of estimating the approximate number of fecal bacteria in water by the fermentation tube was first used in the study of the self-purification of streams by Smith and Brown* (1893).

The work of these authors was carried out upon the Mohawk and Hudson rivers, under the auspices of the New York State Board of Health, and aided materially in the solution of local problems. Owing, however, to the relatively low degree of pollution obtaining in these particular river waters, as is shown both by the chlorine determinations and by the small number of fecal bacteria, the contrast between different points along the course of the river is not very marked.

Some investigations also have been carried out by Hammerl* upon the river Mur above and below Gratz, but the number of this author's determinations, as recorded in his article, are so few, and his method for the detection and enumeration of colon bacilli** is so inadequate that not much weight can be attached to his conclusions.

From a general standpoint, it is clear that the question of self-purification can be most advantageously studied where the proportion of sewage added to river water is very high, so that slight fluctuations, due to temporary and local conditions, to incomplete mingling and to other minor factors, are wholly submerged by a gross and constant pollution. A particularly favorable opportunity has been afforded in the course of the investigation of the conditions created in the Illinois Valley by the discharge of the sewage of the city of Chicago into the Desplaines River. The enormous initial pollution and the fact that during certain seasons of the year the dilution from rainfall and run-off is slight* render it a comparatively easy task to trace the progressive purification in the flowing stream. Few other rivers, in which the process of self-purification has been studied, present so fortunate a union of three important conditions—extreme pollution, relatively little dilution and great length.


^{*} New York State Board of Health, Report for 1893, p. 680.

^{*} Hyg. Rundschau, 1897. XI, p. 529.

^{**}Suspicious colonies upon a gelatin plate were fished and tested "for ability to grow at blood temperature, to curdle milk and to produce gas in media containing sugar," op. cit., p. 537.

^{*} A chlorine content as high as 40 (parts per million) has been found at the mouth of the Illinois.

Comin O. JORDAN. ALBUMINOID AMMONIH - - - PARTS PER MILLION ILLINOIS RIVER.



Methods. In the beginning of the work use was made of the method of direct inoculation of water into the fermentation tube, as suggested by Theobald Smith in 1893, but this procedure was soon abandoned in favor of another method, which was continued through the major part of the investigation. This consisted in a preliminary incubation of a measured quantity of water in carbol-broth. The carbol-broth was prepared by adding I c.c. of a I per cent solution of carbolic acid in sterile water to tubes containing o c.c. of sterile broth of the standard composition. The use of measured quantities of fluid in this way, of course, necessitates allowance for evaporation during sterilization. By careful attention to the size of the tube and to the period of sterilization in the Arnold steam sterilizer, it has been found possible to calculate very closely the loss of fluid during heating and to make due allowance for it; subsequent evaporation before use has been guarded against. The carbol-broth, which we have used, has been first rendered neutral to phenolphthalein and then acidified by the addition of 5.5 c.c. of normal acid per liter.

In carrying out the method I c.c. of a suitable dilution of the water has been added to the tube of carbol-broth and incubated at 38° for 18-24 hours. Platings from this broth have then been made in litmus-lacrose-agar (5 c.c. normal alkali per liter). If red colonies developed on the medium at 38° they were transferred to tubes and tested at once for (1) Gas-production in dextrose broth in the fermentation tube; (2) Indol-production in sugar-free broth; (3) Coagulation of milk; (4) Liquefaction of gelatin.

During a part of the investigation another method was employed, consisting of the introduction of water directly into dextrose broth fermentation tubes without preliminary incubation. The dextrose broth was prepared with fresh meat from which the muscle-sugar had been removed by Smith's method, and to this sugar-free broth I per cent of dextrose was subsequently added. The broth was made neutral to phenolphthalein. After inoculation with the water the tubes were incubated at 38° for 48 hours, gas readings being taken at 24-hour intervals. At the end of 48 hours all tubes showing the formation of gas were removed from the incubator, cooled to the room temperature and the absorption of CO₂ determined by the addition of a 2 per cent solution of NaOH. It has been found necessary to take precautions against incomplete absorption of the CO₂, especially where large tubes are used and the amount of gas formed is considerable.

The use of litmus-lactose-agar for plating water direct proved entirely unadapted to the conditions of our work, and frequently failed to reveal the presence of the colon bacillus, when the two methods above mentioned showed conclusively that this bacillus was present.

Although many other methods were experimented with and carefully compared during the course of the work, only findings obtained by the carbol-broth and dextrose fermentation methods are included in the following tables. A comparison of the two methods (Irons,* 1900) showed that while in the main the results tally closely, the carbol-broth

^{*}American Public Health Association Report, 1900.

method is in general to be preferred for highly polluted waters, while for relatively pure waters the use of the fermentation tube direct appears to be a slightly more delicate test.

The interpretation of the results obtained by the respective methods demands some explanation, since the whole inquiry hinges upon the meaning of the records. It may be said at the outset that no attempt is made in this paper to record separately the occurrence of various members of the colon group of organisms or to pass judgment upon the sanitary significance attaching to the presence of various kinds of colon and paracolon bacilli. For reasons easily understood, such a subdivision of material would be entirely foreign to the problem under consideration. A general and arbitrary standard has of necessity been chosen. In the present state of uncertainty among bacteriologists regarding classification within the colon group, I have thought it best for the purposes of this investigation to adopt a somewhat comprehensive grouping and to include among "colon bacilli" (or "fecal bacteria") certain colon-like organisms showing a fundamental biological relationship. It should perhaps be expressly stated that the term "colon bacillus" is employed in this paper in this general sense, and is not used to designate a sharply defined single "species."

The carbol-broth method, as described above, has given results variously designed in the tables as +, - or? The sign +, as here used, indicates that colonies have been isolated which gave the typical characters of B. coli in (1) Fermentation tube (dextrose broth); (2) Sugarfree Broth for indol; (3) Milk; (4) Gelatin.

The sign — is used to denote those cases where, upon plating in litnus-lactose-agar, after incubation in the carbol-broth, careful search failed to reveal any red colonies; under this head also are placed those cases, not very numerous, where pure cultures obtained from a red colony have failed to yield an excess of H in the fermentation tube.

In the doubtful class are included those instances sometimes encountered where the organism isolated from a red colony produces the typical mixture of gases in the fermentation tube, but fails to respond positively to one or two of the other characteristic biological tests, (2) being the determination most frequently at variance.

The results obtained by the dextrose-broth method are tabulated on the following basis: The sign + is used for those inoculations yielding a total gas-production of more than 20 per cent of the tube length and showing on absorption an appreciable excess of H. If pure cultures are isolated from such tubes, organisms possessed of the biological characters above cited will almost invariably be found. There is a mixture of different kinds of organisms in these tubes, and it is sometimes necessary to examine a great many colonies. When this has been done we have rarely failed to find the colon bacillus. The error involved in the assumption that a member of the colon group is always present in these cases is probably less than 5 per cent. The close agreement of the results obtained by this method with those reached by the carbol-broth method, which appears on the face to be more rigorous, lends further countenance to this view.

Under the sign — are included those determinations in which no gas or only a small amount of gas—less than 10 per cent—was produced. There is perhaps a larger measure of uncertainty regarding the determinations classified under this head; it is possible that the colon bacillus was present in some instances where no gas or only a slight amount was formed, but these cases must have been rare, since we have never been able to isolate the colon bacillus on gelatin or litmus-lactose-agar plates made from such tubes. An exception must, of course, be made to this statement in cases where sewage or highly polluted water has been inoculated into the fermentation tube without proper dilution, since in such cases it sometimes occurs that only a small amount of gas collects in 48 hours, the colon bacillus being apparently overgrown by other sewage bacteria.

In the doubtful class are placed those determinations giving a total gas-production of 10-20 per cent, and those with a total gas-production of more than 20 per cent and absorption test showing an appreciable excess of CO. The majority of the determinations so classified might fairly be regarded as negative, and I believe that an error of not more than 10 per cent would be incurred if this were done. I have preferred, howver, to adopt the more unequivocal arrangement.

SPECIAL OBSERVATIONS AND EXPERIMENTS.

All the results that have been recorded in the foregoing tables have been obtained with transported samples of water, and, so far as the bacteriological side is concerned, they are unquestionably open to criticism on this score. It will be observed, also, upon reference to the detailed tables (tables 81-158) that at most stations the first examinations of water were made toward the end of the high-water period, when the numbers of bacteria were relatively higher than during the prolonged low-water period that followed. It was found impossible to arrange for the collections of water samples to begin at all the stations at the same time, although every effort was made to this end. At West Alton, owing to local difficulties, regular collections of Missouri River water were not instituted until July 27, so that the average given in the table is considerably lower than would have been the case if high-water figures for the Missouri had been obtained. The resulting averages for the Missouri River are hence considerably lower than those for the cross-section at the Chain of Rocks, and a comparison would be misleading.* The averages for all the stations on the Illinois River are, however, quite strictly comparable throughout.

As has been already stated, we have endeavored to supplement and control the results obtained from transported samples by numerous examinations made immediately at the point of collection. A series of samples collected at Bridgeport during the summer months and plated directly gave much larger counts than those resulting from the plating of the

^{*}The average at the 1nlet Tower at the Chain of Rocks for a period corresponding with that covered by the Missonri River analysis is 6,900 (Missonri River at West Alton, 8,200).

270 285 300 315 KAMPSVILLE FREE AMMONIA-PARTS RES MILLION IN ILLINOIS RIVER. 120 135 150 165 180 195 210 225 240 255 HAYAND BEARDSTOWN 150 "AVERYVILE WESLEKCITY WESLEKCITY BRIDGEPORT-JOLIET- MORRIS LA SALLE MENRY LOCKPORT OTTAWA 90 105 15 30 45 60 75 0/ 73 12 9 3

transported samples. This is undoubtedly due to the destruction of bacterial life in the ice-packed sample.*

The correct average from May to August would unquestionably be upward of 2,000,000. The Lockport samples also, especially those for August, show in a marked degree the diminution due to ice-packing. The Morris samples for July 17, August 21, August 29, September 16 and October 16 also show the effect of transportation; a series of twelve samples collected at different times during these same months and plated immediately never afforded numbers so low as those recorded on these dates. On one occasion, for example, three samples of water were plated at Morris immediately after collection (initial temperature of the water, 28° C.), and gave respectively 535,000, 412,000 and 329,000 colonies per c.c. The bottles were packed in ice by the ordinary method and shipped at once to Chicago, where the samples were plated in the usual routine. The count obtained after transportation was, respectively, 54,500, 50,500 and 73,500. If it were necessary, examples of this sort might be multiplied indefinitely.

The diminution of numbers that takes place in ice-packed samples does not, however, result in a stable condition; after a time renewed bacterial reproduction sets in, even when the water is kept constantly at a low temperature; in fact, the numbers may rise to a point higher than that originally obtaining. This secondary multiplication occurred not infrequently in the waters from the lower end of the Illinois River and in those from the Mississippi River, since these waters had to be transported nearly 400 miles before reaching the laboratory. At Grafton, direct platings from the Illinois and Mississippi Rivers gave almost invariably lower counts than were obtained from the transported samples.

For example:

GRAFTON. Plat	ings Dir	ect.	Platings from Same Bottles after Shipment.		
Illinois River October 255	345	270	1,200	800	440
Mississippi River "	2,020		4,500	1,500	• • •
Illinois River November 225	160	325	1,500	830	580
Mississippi River " 1.200	850	1.150	2,600	5.800	540

Multiplication of bacteria in transit was also shown in a marked degree in the samples collected at the Chain of Rocks, where, owing to the fact that the place of collection is difficult of access and a long boatrow is necessary, the packing in ice was unavoidably delayed, and the laboratory counts were uniformly higher than those made on samples plated immediately after collection. There is ample evidence, therefore, to support the view that during most of the period covered by these analyses the recorded averages range lower than the true figures as regards the collecting stations near Chicago, and are higher than is actually the case as regards the more distant points. The apparent difference between the number of colonies found, for instance, in the Illinois River at Averyville and Grafton may be explained in this way; the real difference is inconsiderable.

^{*}c.f. Jordan and Irons. Notes on Bacteriai Water Analysis. American Public Heaith Association Reports, 1899, XXV., p. 564.

It will be observed that the analyses of the water samples from Pekin and Wesley City show great irregularities from week to week. The peculiar local conditions are responsible for these marked fluctuations. The major part of the refuse poured into the river at Peoria enters on the right bank, but is deflected in various ways by sandbars and cross-currents, so that the sample from midstream at Wesley City sometimes shows great pollution, sometimes very little. The influence of these intricate and continually changing conditions is frequently evident in the Pekin samples as well. A cross-section taken at Wesley City on June 13, 1900, gave the following result:

	Free Ammonia.	Albuminoid Ammonia.	No. Colonies per c.c.
Right Bank	504	.864	4.340,000
Center		.248	40,000
Left Bank	168	.240	2.400

Slight changes in the position of sandbars and in the height of water deflect the main mass of polluted water now to one side, now to the other, and so lead to sudden and marked fluctuations in the analyses. Even at Pekin the mingling is often far from complete.

Early in the investigation the importance of following, so far as practicable, the changes taking place in one and the same body of water was recognized, but the pressure of routine work rendered such studies few in number. The most important of these special investigations were carried out between Morris and Ottawa, where laboratory experience had shown us that a change took place that might be properly denominated as purification. Several series of observations were made upon this stretch of river, but as they all led to the same result, only the two most important will be here described.*

The first of these was carried out on October 7 upon a stretch of the Illinois River just below Morris. The day was bright and sunny, the temperature of the air being 7° C. at 6 o'clock in the morning and reaching 20.5° by midday. A slight breeze ruffled the surface of the water in the middle of the day, but was at no time strong. The river was very low (5 feet) and the current exceedingly sluggish. The upper cross-sections were taken at a point just above the Mazon River, the lower about threefourths of a mile below the mouth of the Waupecan Creek. (Neither of these streams was contributing any water to the Illinois at this date). This stretch of river is almost exactly three miles in length. The rate of flow between the two points was determined by weighted floats and by the use of fluorescein solution, and was found to be very close to one-half mile per hour. Four series of cross-sections at hourly intervals were taken at the upper station (A), and these were followed by a similar series at the lower station (B), beginning six hours later. Platings were made within the hour. The samples designated as from the "right" and "left" banks, respectively, were taken midway between the shore and the

^{*}I am greatly indebted to my chief assistant, Mr. E. E. Irons, for aid in the planning and conducting of these somewhat ardness observations, and I am glad to acknowledge that their accuracy and completeness are largely due to the signal zeal and ability with which he devoted himself to this work.

center of the stream. At A the river was about 150 yards wide, at B about 125.

The figures given for the number of colonies are the averages of counts of two separate platings.

UPPER STATION (A).

Hour		No. Colonies per c.c.	Turbidity (Hazen's scale).	Temp. Water °C.	Chlorine (parts per million).
6:15 а. м.	Right Bank	. 500,000	.16	13.	
• 4	Center	. 378,000	.12	13.	
4.4	Left Bank	. 42,000	.075	13.	
7:15 A. M.	Right Bank	. 368,000	.17	13.	
6.6	Center	. 344,000	.125	13.	
s 6	Left Bank	. 35,000	.0775	13.	
8:15 л. м.	Right Bank	. 752,000	.16	13.5	91
k 6	Center	. 364.000	.11	13.	69
**	Left Bank	. 30,000	.0675	13.	45
9:15 а. м.	Right Bank	. 554,000	.16	14.	
4.4	Center	. 472,000	.11	14.	
6.6	Left Bank	. 79,000	.075	14.	

LOWER STATION (B)

(Three miles below A).

		No. Colonies	Turbidity	Temp. Water	Chlorine
Hour.		per c.c.	(Hazen's scale).	°C.	(parts per million).
12:15 р. м.	Right Bank	. 480,000	.13	16.	
4.6	Center	. 327,000	.15	16.	
4.6	Left Bank	. 87,000	.05	16.5	
1:15 р. м.	Right Bank	. 281,000	.1475	., 16.	• • • • • • •
4.4	Center	. 102,000	.09	16.	
**	Left Bank	. 19,000	.042	16.	
2:15 р. м.	Right Bank	. 400,000	.13	17.	87
4.4	Center	. 249,000	. 09	16.	72
4.6	Left Bank	. 22,000	.045	16.	52
3:15 р. м.	Right Bank	. 412,000	.136	17.	82
4.4	Center	. 416,000	.12	16.	78
1.4	Left Bank	. 11,000	.0433	17.	53


The averages may be tabulated as follows:

	No. Co	No. Colonies per c.c.			
Distance.	Right Bank.	Center.	Left Bank.	Analyses.	
Upper Station (A)	543,700	389,700	46,500	4	
Lower Station (B) 3 miles (6 hours). 393,250	273,500	34,750	4	
Percentage Decrease	27.6	29.8	25.3		

The chlorine determinations show that the mixing of the Kankakee water and the Desplaines is very incomplete both at the upper station (9.7 miles below the junction of the rivers) and at the lower, and this is entirely confirmed by the bacterial and turbidity cross-sections at the two points. Between A and B a great bacterial diminution occurs, and this in almost equal degree both along the seriously polluted right bank and along the comparatively uncontaminated left bank.

A second series of observations was carried out in a similar fashion upon a longer stretch of river. The distance from the regular collecting station at Morris to the regular collecting station at Ottawa (see map) is about 24 miles, and the rate of flow between the points averaged one-half mile per hour at the time our observations were made. A point midway between Morris and Ottawa was selected (Seneca) and a three-day series of observations was planned. The sun was wholly obscured by

NUMBER BACTERIA PER CUBIC CENTIMETER AT GRAFTON.



clouds during these three days, but no rain fell. The results are given as before in tabular form.*

UPPER STATION (MORRIS).

Hour.		No. Colonics per c.c.	Turbidity (Hazen's scaie).	Temp. Water °C.	Chlorine (parts per millon).
7:15 л. м.	Right Bank	433,000	. 153	7.	••••
6.6	Center	. 337,000	.13	7.	
b 6	Left Bank	30,000	.046	7.	
11:30 а. м.	Right Bank	177,000	.17	7.25	67.5
44	Center	. 145,000	.18	7.25	47.5
44	Left Bank	17,000	.02	7.25	8.
2:00 р. м.	Right Bank	174,000	.15	3.	
6.6	Center	131,000	.135	3.	
6.6	Left Bank	49,000	.05	3.	

MIDDLE STATION (SENECA).

Hour			Turbidity (Hazen's scale).		Chlorine. (parts per million).
9:00 A. M., Nov. 10.	Right Bank	134,000	**	9.5	51
66 66	Center	47,000 {	*Less than .09, more than .07.	9.5	44
44	Left Bank	. 23,000	more, thun .or.	9.5	35
1:30 р. м., Nov. 10.	Right Bank	67,000		11.	
66 66	Center	52,000		11.	
4.6	Left Bank	52,000	• • • • • • •	11.	

^{*} Turbidity readings could not be taken accurately at this point and at Ottawa owing to presence of water weeds.

LOWER STATION (OTTAWA).

Hour.			No. Colonies per c. c.	Turbidity (Hazen's scale	Temp. Water	Chlorine (parts per million).
10:00 A. M.,	Nov. 11.	Right Bank	. 11,000		9.5	49
6.6	6.6	Center	10,500	Less than .04.	9.5	46
4.6	4.6	Left Bank	3,900 (9.5	43
1:00 р. м.,	Nov. 11.	Right Bank	. 12,000		9.5	
4.6	6.6	Center	11,000	• • • • • • •	9.5	
44	4.6	Left Bank	. 18,000		9.5	

The averages are as follows:

Distance	No. Co	No. of Hourly.		
from Morris.	Right Bank.	Center.	Left Bank.	Analyses.
Upper Station (Morris)	261,000	204,000	29,000	3
Middle Station (Seneca) 12 miles. 1 24 hours.	103,000	49,000	35,000	2
Lower Station (Ottawa) \ 24 miles. \ 48 hours	11,500	10,700	13,500	2

During this flow of 24 miles, therefore, the Illinois River became nearly free from the great mass of sewage bacteria with which it was originally laden. In fact, the bacterial content of the Illinois at Ottawa was not greatly in excess of that of the local tributary streams. The number of colonies found in the water of the Fox River on November II was 6,850 (av.), a number not much lower than that found in the Illinois (II,900).

* In connection with this series may be given the averages of the regular chemical determinations for the period between October 23 and November 20. These are:

					Ox	ygen Consum	ed.
	No. Determi-	Res	idue on Eva	poration.	j	By Dissolved	By Suspended
Station.	nations.	Total.	Dissolved.	Suspended.	Total.	Matter.	Matter.
Morris	. 5	398	380	18.	10.7	8.4	2.3
Ottawa	. 5	356	353.5	2.5	7.5	7.3	.2
	Nitrog	en as					
Free Ammon	ia.	Alb	ouminoid An	ımonia.	nonia. Nitrites. Nitra		
		Total.	Dissolved.	Suspended.		Mirites.	Nitrates.
7.98		.860	.478	.382		.028	.340
6.6		.364	.315	.049		.382	.648

It was thought desirable to supplement these observations with another series made after the Sanitary Canal was opened, and accordingly, on May 24-26, 1900, another series of observations was made upon the stretch of river between Morris and Ottawa. The sun was partially obscured on May 25, but shone brightly on May 24 and 26; no precipitation occurred during the period.

UPPER STATION (MORRIS).

		No. Colonies	Temp. Water	Chlorine
Hours.		per e. e.	°C.	(Parts per million).
7:30 A. M., May	5. Right Bank	45.000	15.5	20
11 14	Center	74,000	17.5	13
4.4	Left Bank	42,000	18.	8
9:30 A. M., May	25. Right Bank	60,000	15.5	
"	Center	34,000	17.	
44 44	Left Bank	39,000	18.	
11:30 A. M., May	25. Right Bank	98,000	15.75	****
"	Center	62,000	17.	
11 (1	Left Bank	30,000	18.	
*2:00 г. м., Мау	25. Rlght Bank	70,000	16.	21
	Center	63,000	17.	14
64 46	Left Bank	48,000	18.	6

* Samples were taken at this time and submitted to chemleal examination with the following result:

	Right Bank.	Center.	Left Bank.
Total residue on evaporation	232	246	308
Dissolved residue on evaporation	230	244	285
Suspended residue on evaporation	. 2	2	23
Chlorine	. 21	14	6
Oxygen consumed—Total	6.6	7.2	8.8
Oxygen consumed by dissolved matter		6.0	4.7
Oxygen consumed by suspended matter	5	1.2	4.1
Free ammonla	2.28	1.08	.24
Albumlnoid ammonia—Total	. 256	.296	.240
Albuminoid ammonia—Dissolved		.224	.168
Albumlnoid ammonia suspended		.072	.072
Nltrites	. 072	.050	.040
Nltrates	4	1.1	. 95

MIDDLE STATION (SENECA).

Hours.		No. Colonies per e.e.	Temp. Water	Chlorine (parts per million).
7:30 A. M., May 25.	Right Bank	13,800	16.5	19.5
6.6	Center	11,700	* * * * * *	16
6.6	Left Bank	4,100		13
9:30 л. м., Мау 25.	Right Bank	12,700	17.	
66 66	Center	14,000		
66 66	Left Bank	16,000		
H:30 л. м., May 25.	Right Bank	4,500	17.5	••••
64 64	Center	4,100		
	Left Bank	3,000		****
1:30 г. м., Мау 25.	Right Bank	5,500	18.5	16
6.6 6.4	Center	6,700	18.8	15
4.6 4.6	Left Bank	13,100	19.7	12

LOWER STATION (OTTAWA).

Hours.			No. Colonles per c.c.	Temp. Water	Chlorine (parts per million)
8:30 A. M.,	May 26.	Right Bank	7,400	18.5	15
6.6	6.6	Center	10,000	18.5	14
6.6	6.4	Left Bank	8,300	18.5	13
#1:00 A. M.,	May 26.	Right Bank	8,100	19.	17
4.4	11	Center	6.800	19.2	15
4 t	1.1	Left Bank,	6,400	19.2	14
2:00 г. м.,	May 26.	Right Bank	9,000	19.5	17
6.6	6.4	Center	6,300	19.5	16
11	£ 6	Left Bank	1,200	20.	14


	Distanee	No. Colonies per c.e.			No. of Sep-
	from Morris.	Right Bank.	Center.	Left Bank.	arate Analyses.
Upper Station (Morris)		68,200	58,200	39,800	4
Middle Station (Seneca)	12 miles.	9,100	9,100	10.300	4
Lower Station (Ottawa)	24 miles.	8,200	7,700	6,300	3

GENERAL SUMMARY.

In proceeding to summarize our results from the special point of view of this investigation, namely, the effect of the opening of the Sanitary Canal upon the condition of the Illinois and Mississippi Rivers, it is important to fix attention upon the main features of the problem. particular comparison that it is desired to institute here is based, perforce, on a comparison of the period before the Sanitary Canal was opened with the period after the canal had been put in operation. This would at first sight seem to necessitate a comparison of the results obtained from May I, 1899, to January I, 1900, with those obtained from January I, 1900, to July 1, 1900. It will be recognized, however, that such a comparison would be misleading, since it is well known that different seasons of year are not properly comparable with one another, owing to differences in the temperature of the water, the rate of flow and other controlling factors. (See, for instance, the diagram of the colony count in the Illinois and Mississippi Rivers at Grafton). Thus a comparison of the winter conditions of 1899-1900 with the summer conditions of 1899 could have no practical bearing upon the problem under consideration. Fortunately, there is a partial seasonal overlapping which permits direct and proper comparison of the conditions obtaining in the river waters before the Sanitary Canal was put into operation with those after that event. analytical results secured during the months of May and June, 1899, may fitly be compared with those obtained in the same months of the following year. For this reason I shall present in the following diagrams a comparison of certain averages obtained during the first period (1899, canal not open) with the corresponding averages for the second period (1900, canal in operation).

It is believed that these diagrams will readily explain themselves; nevertheless, attention may be drawn to one or two features. The marked drop in chlorine in the upper part of the river during the second period witnesses to the greatly increased dilution of the sewage with the purer water of Lake Michigan. The increased volume of chlorine-laden water in the latter period, on the other hand, betrays itself in the increased amount of chlorine in the lower stretches of the river. The diagrams representing the relative amounts of free ammonia and albuminoid ammonia and the number of bacteria per cubic centimeter show similarly the effects of the increased dilution, and they also show that the change occurring in the significant substances in the water, both soluble and suspended, runs an essentially parallel course in the two years. For the period under consideration, therefore, there is substantial evidence that the amount of nitrogen in the form of free and albuminoid ammonia, and the number of bacteria per cubic centimeter, as shown by the ordinary plate count.

NUMBER BACTERIA PER CUBIC CENTIMETER IN ILLINOIS RIVER.



were at least no greater in the Illinois River at its mouth after the canal was opened than before that time. It is possible to go even further than this and to assert that the amount of the specially significant organic constituents has at most periods been no higher than is to be reasonably expected of river waters in this part of the country, as, in fact, is shown by the analyses of water from the tributary streams of the Illinois and from the Mississippi and Missouri rivers.

Is it, then, to be concluded that the discharge of the Chicago sewage into the valley of the Illinois has no effect whatever upon the Illinois River at its mouth beyond increasing the chlorine and other mineral constituents? The question cannot be answered summarily. While it is true that much of the nitrogen content of the original sewer is converted into gaseous compounds which escape from the water, it is also true that a large part of the ammoniacal compounds is oxidized to nitrates. After nitrification has become complete, or nearly so, the river water is consequently rich in nitrates and affords an especially favorable medium for the growth of microscopic algæ and water plants. These organisms abound in the lower stretches of the river during the summer months, and a certain proportion of the nitrogen originally present in the Chicago sewage unquestionably finds its way into their bodies. Waters containing large numbers of such organisms will of course show "high albuminoid ammonia or organic nitrogen" on analysis, although the nitrogen in this condition obviously has not the same sanitary significance as has the "organic nitrogen" of fresh sewage. Many of these organisms, however, either have definite cycles of development through which they pass, or else they perish at the approach of cold weather or in the presence of other injurious influences. The nitrogen in their dead bodies now begins a new series of changes and passes through the stages of "free ammonia" and "nitrites" to "nitrates." There it may again be taken up by a fresh set of plant organisms and enter into a new cycle of changes, only to terminate eventually in the mineralized nitrates.

In view of these well-known facts, it might reasonably be expected that the nitrogen-content of the Illinois at its mouth would be different from what would have been the case if no sewage from Chicago had ever passed into the river. The considerable amount of nitrogen appearing as "free ammonia" at Grafton at certain seasons of the year can best be interpreted in this way. It is reasonable to believe that it has not come direct and unchanged from the Chicago River, but has passed, perhaps several times, through the bodies of microscopic, chlorophyll-bearing plants. The completeness of nitrification as far up the river as La Salle is a strong reason for maintaining this view. The reappearance of high ammonias in the lower stretches, especially at the onset of cold weather, can only be understood by taking into consideration the cycle of nitrogen above outlined.

Considering the problem as a whole, it must be remembered that it is not so much the history of the nitrogen compounds that is significant, especially where conditions are so intricate as in the present instance, as the story of bacterial life. If the question be plainly put as to whether

typhoid bacteria or similar pathogenic microbes are likely to pass from Chicago to Grafton in the water of the Illinois River under any of the conditions prevailing during our investigation, it must be plainly answered that all the evidence that we have been able to secure is against such an occurrence. It has been already pointed out (pp. -----) that a study of the death rate among the colon bacteria added to the river water in sewage lends no countenance to the view that typhoid bacteria will survive passage down river. The facts indicate that the colon bacteria, which are present in such large numbers in Chicago sewage—undoubtedly in much larger numbers than typhoid bacilli—disappear almost completely in less than 150 miles' flow. Since all investigators are agreed that the colon bacillus is more hardy than its relative, the typhoid bacillus, and can live in water for a longer time, there is every reason for supposing that the latter microbe dies out with at least the same rapidity. Even were typhoid bacteria found in the water at the mouth of the Illinois, there are scores of communities to which they might certainly be more plausibly traced than to Chicago.

Respectfully submitted,

EDWIN O. JORDAN.

Professor of Bacteriology, University of Chicago.

April 1, 1901.

REPORT OF THE UNIVERSITY OF ILLINOIS

BY

PROF. A. W. PALMER, S. C. D.

Arthur R. Reynolds, M. D., Director of Streams Examination, Chicago, Ill.

Dear Sir:—Herewith I submit a report upon the chemical and bacterial analyses conducted at the University of Illinois in behalf of the Sanitary District of Chicago. These analyses cover the period extending from the latter part of April, 1899, to the beginning of October, 1900, and include the chemical and bacterial examination of about three thousand samples.

Regular weekly analyses have been made of water samples collected from the Illinois and Michigan Canal, from various parts of the Desplaines and Illinois rivers, from certain of the principal tributaries of the Illinois River, from the Mississippi River at a number of different points near Alton and St. Louis, from the Missouri River and from the St. Louis water supply. During a portion of the time three or more samples a week were collected at several important points.

The points at which the samples were taken were determined upon in a general way at the conference held in your office upon March 15, 1899; they were more precisely located by Professor Jordan and myself just before the inauguration of the work.

The reasons for the selection of these particular places are indicated below.

The report which I submit is in three parts: Part A, relating briefly to the inauguration of the investigation and the selection of the points of collection. etc.; Part B, relating to the chemical analyses, including the determination of dissolved oxygen, and Part C, by Professor T. J. Burrill, relating to the bacterial examinations.

The report of the chemical examinations includes:

First—Complete tables of the data of the analyses; the few blanks which appear in the tables being due either to the breakage of a bottle and the loss of an entire sample, or to the fact that some mishap during the course of the examination has resulted in the loss of a portion of the sample which could not be replaced.

Second—Tables of averages of the results for various periods.

Third—Plates which exhibit graphically the general results as expressed in the averages of the tables.

In the consideration of the data of the chemical examinations, I

7

have briefly indicated the origin, the characteristics and the significance of some of the more important substances determined, in the hope that such intelligent laymen as are really interested in the problems involved in these investigations may thereby be enabled in some measure to understand the significance of the numerical data and the various lines of the plates.

Throughout the investigations I have been very ably assisted by Mr. R. W. Stark, B. S., and during the greater part of the time by Mr. C. V. Millar, M. S. The continued interest, the skill and the indefatigable zeal of these two gentlemen in conducting most of the routine work of these examinations deserve special commendation. During much of the time covered I have been further assisted by Mr. F. C. Koch, M. S.; Mr. E. P. Walters, B. S., and Mr. A. L. Marsh.


INAUGURATION OF THE EXAMINATIONS.

At the conference held in Chicago, March 15, 1899, at which there were present Mr. Isham Randolph, Chief Engineer of the Sanitary District; Dr. Arthur R. Reynolds, Commissioner of Health of Chicago and Director of Streams Examinations for the Sanitary District of Chicago; Dr. R. W. Riley, Assistant Commissioner of Health of Chicago; Dr. Adolph Gehrmann, Director of the Municipal Laboratory of Chicago; Professor E. O. Jordan, representing the University of Chicago, and Professor A. W. Palmer, representing the University of Illinois, the general plan and scope of the investigations were discussed, and it was decided that samples of water for analysis should be taken from the Illinois River and certain of its tributaries, the Missouri River and the Mississippi River, at a number of different points between Chicago and St. Louis or thereabouts.

It was decided that the work should be done in triplicate, i. e., that three sets of samples should be taken simultaneously at each place each week, and that independent analyses of these three sets of samples should be made by the three institutions represented: that is, that one set of samples should be analyzed at the University of Chicago, one at the Municipal Laboratory of Chicago and one at the University of Illinois.

At this conference the task of visiting and precisely locating the various points at which the samples should be taken, and of selecting, appointing and instructing persons who could be relied upon to collect and ship the samples satisfactorily, was assigned to Professor Jordan and Professor Palmer, who shortly thereafter inspected the localities in question and personally gave instructions and demonstrations to the collectors whom they selected for the work. All of the points except two, namely Kampsville, upon the Illinois River, and Chandlerville, upon the Sangamon River, were visited by Professor Jordan and Professor Palmer conjointly, and each of the more important points has been repeatedly visited by them, either conjointly or separately, at various times during the period covered by the series of examinations.

Each collector was personally instructed in the manner of making



the collections and the care to be exercised in handling the bottles and samples. He was taken to the precise spot at which he should make the collection, and the necessity of making every collection at that particular spot was impressed upon him. Moreover, he was shown precisely how to make the collection, stopper and seal the bottles and pack them for shipment. He was impressed with the importance of putting plenty of ice around the bacterial sample and arranging everything for the utmost promptness of shipment by the quickest route. He was provided with metallic seals and the proper heavy tool for pressing them. In addition to these instructions, a description was printed upon the back of the certificate which he was required to fill out and attach to each sample. A copy of the certificate is given below:

STREAMS EXAMINATION-CHICAGO SANITARY DISTRICT.

SAMPLE OF WATER.

Instructions for Collecting and Sending Samples of Water for Analysis.

The outfit for obtaining water samples consists of two parts. The large gallon bottle is for the chemical sample, and the small four-ounce bottle is for the bacterial sample. The large and the small bottle together constitute one sample or set. The two bottles must be filled at the same time, and at the same place, and in the same general manner. They must then be placed in their respective compartments of shipping cases for transportation.

In filling the bottles, the stoppers should be taken out only when the spot from which the water is to be taken is reached. The bottle is immersed with the stopper in place about one foot below the surface and is filled, leaving about one or two inches air space. The large and small bottles are then immediately sealed by placing the rubber cloth over the stopper and drawing the wire seal with the number for the sample tight about the neck. The small bottle is to be placed in the smaller can and the lid pressed down. It is then to be packed in ice within the large can. The card in the shipping envelope is then to be filled out and attached to the neck of the larger bottle. The bottles should not be filled until just before the time for delivery at the express office.

As series of samples are to be sent to different laboratories, all of the bottles should be filled at the same time, and at the same place, and the different sets placed together as described.

The certificate of collection on the reverse side is to be filled out completely in every case.

(OVER.)

The points determined upon and the collectors selected were as follows:

POINTS AT WHICH SAMPLES WERE TAKEN.

(See map of localities on page 8.)

No. I—At Bridgeport samples were taken from the Illinois and Michigan Canal a short distance below the pumping station. Before the opening of the Drainage Channel the fluid pumped from the Chicago River into the old canal at this point constituted by far the major part of Chicago's sewage, and contained, in addition to house sewage, an enormous amount of manufacturing wastes, including those from stockyards, soap factories, rendering establishments, metallurgical works, gas works, etc., etc.

No. 1a—After the opening of the Drainage Channel in January, 1900, samples were collected from it also at Bridgeport, at the crossing of Kedzie Avenue.

The collections at this point were made by Mr. Joseph Weis.

No. 2—Lockport. The fluid pumped into the Illinois and Michigan Canal flows sluggishly from Bridgeport to Lockport, a distance of about 29 miles. Generally there is practically no dilution and no notable additional pollution of the water. At certain seasons of the year, however, storm water enters the canal in somewhat large quantity. The chemical impurities in the water of the old canal change but very little between Bridgeport and Lockport, owing, probably, to the very great concentration of the sewage, and possibly, too, to the fact that some of the manufacturing wastes retard the incipient decompositions.

No. 3—At Lockport there is some discharge from the Illinois and Michigan Canal into the Desplaines River, a small stream of extremely variable flow. A sample of water was collected from the Desplaines River at a point beside the old stone bridge, a short distance above the Norton Mills, where the first discharge from the old canal into the river takes place. The Desplaines River itself receives some sewage from various suburban towns along its banks above this point. After the opening of the Drainage Channel (January, 1900) the sample collected at this point consisted of a mixture of Desplaines River water with the water of the Drainage Channel, of which mixture the water of the latter amounted to from 25 to 99 per cent, the former only occasionally, in time of freshet, the latter during the last quarter of the year, the average for the entire year being 91¾ per cent for 1901 and 89.9 per cent for 1900.

No. 3a—After the opening of the Drainage Channel samples were collected at Lockport from it also, at a point just above the controlling works, where the discharge from the Drainage Channel into the Desplaines River occurs.

All collections at Lockport were made by Mr. William O'Brien, express agent.

No. 4—The Desplaines River, at Joliet, just above the town. At this point, four miles below Lockport, the Illinois and Michigan Canal and the Desplaines River unite to form a large basin, the canal from this point on passing down beside the west bank of the Desplaines River. Before the opening of the Drainage Channel it was at this point that the sewage in the old canal received its first notable dilution. During the greater part of the year, particularly the summer and autumn seasons, this dilution was very slight, but in the earlier or flood seasons of the year the flow in the Desplaines River at Joliet would at times amount to as much as three or four hundred thousand cubic feet per minute, and the dilution would be correspondingly great.

The average flow from the Illinois and Michigan Canal was formerly about 45,000 cubic feet per minute, but since the opening of the Drainage Channel the flow from the old canal has been reduced to an average of 26,700 cubic feet per minute in 1900 and about 13,800 cubic feet per minute in 1901. The sample of water was taken in the basin just above the town.


No. 5—The Desplaines River at Joliet, just below the town. A large part of the sewage of Joliet, i. e., that from all of the town lying on the east side of the river, enters the Desplaines directly at Joliet. In addition to house sewage, this includes also the wastes from various manufactories. Late in the season of 1899, i. e., in August, the bed of the Desplaines River below Joliet was almost entirely dry, as the flow was practically all carried by the canal. The collections from the river below the town were, consequently, discontinued at this time, and for some time afterward the samples were taken from the canal itself, but at a corresponding point below the town.

The collections at Joliet were made by Mr. Raymond Hurd.

No. 6—The Kankakee River at Wilmington. Formerly the first really considerable dilution of the Chicago sewage resulted from the union of the sewage-laden Desplaines River with the Kankakee River. The mean discharge from the Kankakee River for the year is estimated at three hundred thousand cubic feet per minute, but the flow is subject to exceeding great variations, and at certain seasons it becomes almost negligible. The sample taken at Wilmington gives a fair idea of the composition of this water. The organic matters contained in the Kankakee River water are mainly of vegetable origin, and are derived from the extensive marshes which are drained by this stream. Some house sewage also enters the stream at Kankakee (population, 13,595), about 20 miles above Wilmington and at several other smaller towns along its course.

The collections at Wilmington were made by Mr. R. P. Elliott, express agent.

No. 7—The Illinois River at Morris. This is the first collecting station upon the Illinois River proper, and is $9\frac{1}{2}$ miles below the junction of the Kankakee and the Desplaines rivers. The river at this point is practically a more or less complete mixture of Chicago sewage and Kankakee River water. In a dry season, like that of 1809, the propor-



tion of Kankakee River water may be very small, and it is always comparatively small in the latter half of the year. The average volume of the Kankakee River was formerly four or five times that of the Desplaines River; since the opening of the Drainage Channel it is about equal, i. e., the average flow of each for the year is approximately 300,000 cubic feet per minute. Owing to the fact that the mixing at Morris is quite incomplete, the chlorines given in the various tables do not show very clearly the relative proportions of sewage and of Kankakee River water.

The collections at Morris were made by Mr. J. W. Miller.

No. 8—The Fox River at Ottawa. The Fox River constitutes another great diluting factor. The area drained by the Fox River, 2,697 square miles, is about one-half that drained by the Kankakee River, 5,146 square miles, but the volume of discharge is relatively large. The sample collected from this stream was taken at a point above the town and above the aqueduct which carries the water of the Illinois and Michigan Canal across the stream. The Fox River receives the sewage and manufacturing wastes of a number of towns, of which Elgin (population, 1900, 22,433) and Aurora (population, 24,147) are the largest.

No. 9—The Illinois River at Ottawa. Collection at this point was made at about a mile above the entrance of the Fox River and above the point where town sewage enters, and shows the change resulting from 24 miles' flow without notable dilution and without material addition of impurities, but the mixing of the discharges from the Desplaines and the Kankakee is not yet complete at this point, 33 miles below the confluence of these two streams.

The collections at Ottawa were made by Mr. Lester Horan.

No. 10—The Big Vermilion River at La Salle. This is another important tributary draining an area of 1,413 square miles and receiving sewage from the towns of Streator and Pontiac, as well as the wastes from a number of cement factories situated upon its banks a short distance above its mouth. The sample was collected about one-half mile within the mouth of the stream, which is three miles above the bridge at La Salle.

No. II—The Illinois River at La Salle. The sample at La Salle was taken beside the wagon bridge which crosses the stream at this point. This is about three miles below the mouth of the Big Vermilion, at a point where comparatively little of the sewage of La Salle has entered the stream, and above the point at which the Illinois and Michigan Canal, which receives most of the house sewage of La Salle, finally discharges into the river.

No. 12—The Illinois and Michigan Canal at La Salle. The sample was taken from the lower basin of the canal, just above the point at which it discharges into the river and where it has received considerable of the sewage of the town of La Salle.

The collections at these three points, i. e., Nos. 10, 11 and 12, were made by Dr. William A. Fraser of La Salle.

No. 13—The Illinois River at Henry. Between La Salle and Henry there is comparatively little dilution, but the river receives the final discharge from the Illinois and Michigan Canal and most of the sewage of La Salle in addition to the sewage of Peru. The sewage from both of these towns includes considerable manufacturing wastes from zinc works, fertilizer works, etc., etc., in addition to house sewage. The sample at Henry was taken above the dam.

The collection at Henry was made by Mr. H. McCune, keeper of the state lock.

No. 14—The Illinois River at Averyville. The collecting station was at the bridge over the Narrows, about three miles above the city of Peoria. The results obtained here show the degree of purity, both bacterial and chemical, attained during the flow of 130 miles from Lockport, the point at which the sewage is first discharged into the Desplaines and Illinois River Valley.

The collections were made by Mr. Robert Martin, engineer, under the supervision of the superintendent of the Peoria Waterworks, the wells and pumping station of which are located quite near the bridge.

No. 15—The Illinois River at Wesley City. The city of Peoria (population, 1900, 56,100) contributes a large amount of organic refuse to the river between points No. 14 and No. 15. Not only does the main part of the house sewage enter, but there is also a great addition of manufacturing wastes and distillery slops and discharges from glucose factories and the drainage from extensive cattle sheds and stockyards, The Wesley City collection was made about four miles below this outpour of pollution. The amount of the Peoria pollution varies greatly at different seasons in the year and at different hours of the day, a fact that aids in explaining irregularities and fluctuations, particularly in the number of bacteria. Kickapoo Creek discharges into the Illinois River just above this point; it carries considerable organic matters derived from distilleries which drain to it. At certain seasons of the year its flow is considerable, and as the distance from its mouth to the collecting point, Wesley City, is not great, considerable variations are often due to the incomplete mixing of its waters with those of the Illinois.

No. 16—The Illinois River at Pekin. The samples were collected at the bend in the river several hundred yards below the last point at which sewage is discharged into the stream. The house sewage of Pekin, the discharges of distillery slop and those from large cattle-sheds and stockyards, glucose factories, fertilizer works, and so on, enter the river here.

The collections at Pekin and at Wesley City were made by Mr. D. H. Jansen of Pekin, county surveyor of Tazewell County.

No. 17—The Illinois River at Havana. The collection here was made above the town and above the mouth of the Spoon River. The results of the examinations show the changes which take place in the impurities contained in the water during the flow of 30 miles from Pekin to this point. Between Pekin and this point there is no notable addition

of impurities, and the dilution from small tributaries is ordinarily comparatively slight.

The collections at Havana were made by Mr. H. G. Heberling.

No. 18—The Sangamon River at Chandlerville (drainage area, 5,592 square miles). The sewage of various towns, some of them of considerable size, enters this stream at various points along its course, notably that of Springfield, with a population of 34,159 (1900), which enters about 45 miles above Chandlerville.

The collections at Chandlerville were made by Mr. E. O. Spink.

No. 19—The Illinois River at Beardstown. Little organic impurity is added to the river below the mouth of the Sangamon. The examinations of the water at Beardstown serve to show the changes wrought by the natural purifying processes and by the mixing of the Illinois River water with the waters of the Sangamon River.

The collections at Beardstown were made by Mr. G. W. Barney-castle, under the supervision of Mr. J. A. Carney of the C., B. & Q. Railroad.

No. 20—The Illinois River at Kampsville. The sample at this point was taken just above the United States dam, which is about 30 miles above the mouth of the river. There is no notable discharge of impurities into the river between Beardstown and Kampsville, and, indeed, between Beardstown and Grafton, but a number of small streams discharge into the Illinois between these points.


The collection at Kampsville was made under the direction of Mr. C. V. Brainard, assistant United States engineer, who is in charge of the dam and of the work of the improvement of the Lower Illinois River.

No. 21—The Illinois River at Grafton. A sample of water was taken at a point about two miles within the mouth of the Illinois River and above the point at which any mixing of the water of the Mississippi with the water of the Illinois occurs.

No. 22—The Mississippi River at Graffon. A sample was taken from mid-channel of the Mississippi River at a point above the mouth of the Illinois and where there can be no mixing of the Illinois River water with that of the Mississippi.

The collections at Grafton were made by Mr. B. F. Robinson.

Nos. 23-27—Mississippi River at Alton. At Alton five samples of water were taken from the Mississippi River above the town and at points opposite the new pumping station of the Alton waterworks, as follows: The first sample at 100 feet from the Illinois shore, the second at one-fourth distance from the Illinois shore across stream, the third sample from mid-channel, the fourth sample at one-fourth distance from the Missouri shore across stream, and the fifth sample at 100 feet from the Missouri shore. No sewage from Alton enters the river within half a mile or more of this point, and the drainage from the two or three hamlets between Alton and the mouth of the Illinois is so inconsiderable that the data from these cross-section samples would seem to fairly represent the amount of mixing of the Mississippi and the Illinois River



waters during the flow from Grafton to Alton, a distance of about 17 miles.

The collections at Alton were made by Mr. George Brooks.

No. 28—The Missouri River at West Alton. The condition of the Missouri River just before it mixes with the Mississippi is shown by the collections at this point. The samples were taken just beside the C., B. & Q. Railway bridge at Bellefontaine.

The collections at West Alton were made by Mr. James Mathews.

Nos. 29-32—The Mississippi River at "The Chain of Rocks." At Mitchell, or "The Chain of Rocks" that is, at the point where the pumping station of the St. Louis waterworks is situated, four samples from the Mississippi River were taken, as follows: The first 400 feet from the Illinois shore, the second from mid-channel of the Mississippi River, i. e., about midway between the intake tower of the St. Louis waterworks and the Illinois shore; the third sample immediately beside the intake tower of the St. Louis waterworks pumping plant, and the fourth sample at about 400 feet from the Missouri shore, between the intake tower and the Missouri shore.

The collections at this place were made by Mr. Henry Atkins, a prominent farmer residing upon the banks of the Mississippi at this point, the nearest railway station being at Mitchell, Ill., about four miles away.

No. 33—St. Louis Tap Water. The water drawn from the river at the intake at the "Chain of Rocks" is passed through several settling basins before it is delivered to the consumer in the city. The grosser effect of subsidence is shown by the great reduction of suspended matters from 876 parts at the intake tower to 98 parts in the tap water as delivered in the city in 1899 and 1,330 and 97 respectively in 1900.

Nos. 34-38—Mississippi River at Jefferson Barracks. A cross-section of the Mississippi River was made six miles below the city of St. Louis, i. e., nine miles below the Eads bridge, for the purpose of determining the effect produced upon the Mississippi River water by the addition of sewage and manufacturing wastes of the city.

The collections at Jefferson Barracks and from the St. Louis tap were made by Mr. Emil A. Appel and Mr. August Johnson.

The collectors at La Salle, Averyville, Kampsville, Grafton and Alton had already had considerable experience in collecting water samples for the State Water Survey.

The map on page 8 shows the relative situations of the various points of collection.

METHODS OF PHYSICAL AND CHEMICAL EXAMINATION.

When the samples were received at the laboratory a serial number was immediately placed upon each bottle and upon the tag or certificate which accompanied it; then the rubber cloth which covered the stopper was removed, the stopper and neck of the bottle cleaned, and, after

withdrawing the stopper, some of the water was so poured out as to rinse off the lip of the bottle.

After noting the turbidity, but before beginning the analysis, the sample was thoroughly shaken and every effort was made to keep all solid matters in suspension while the portions were being taken for the various determinations. Nearly half of the sample was immediately filtered through heavy Swedish filters, which had been previously washed with nitrogen free water. Often it was necessary to filter more than once.

The nitrites were always determined immediately upon reception of the sample in the laboratory; the determinations of nitrates, the ammonias and oxygen consumed were begun at once also, and the others were started as soon as possible. Some of the determinations, as the total solids and chlorine, were ordinarily not finished until several days after that upon which the sample was received in the laboratory.

In the tables of results, the date of collection indicates the date placed upon the collector's tag at the time the sample was taken. The date of analysis refers to the time of the receipt of the sample in the laboratory, which also invariably represents the day when the analysis of the sample was begun.

Turbidity and Sediment.—The amount of sediment and the degree of turbidity were noted from mere visual inspection at the time the sample was received and again in a portion of the sample after standing over night, and are indicated in the tables of results by the very approximate terms, "slight," "distinct," "decided" and "much," to indicate the degree of turbidity, and the terms, "very little," "little," "considerable" and "much," to indicate the relative quantity of sediment. A more definite idea of the amount and nature of the suspended matters is, of course, to be had from the figures recorded in the respective columns under the headings: "Total Solids," "Loss on Ignition," "Oxygen Consumed," "Albuminoid Ammonia," etc.

Odor.—Note was made of the odor after thoroughly shaking the water in the bottle just before the portions of the sample were poured out for the determination of the various constituents, and the result of the observation was roughly expressed as "oily," "gassy," "musty," "none," etc.

The Color.—The color of the water was determined by comparison with the color developed in the ammonia standard solution used in ness-lerization; in most cases it was necessary to filter the water for this purpose.

The figure recorded in each case represents the volume of standard ammonium chloride solution required to develop the same tint when diluted to fifty cubic centimeters with ammonia free water and treated with the usual amount of nessler reagent; that is, the color recorded as I. represents the color developed by nesslerization of a solution containing one cubic centimeter of the standard ammonic chloride solution diluted to fifty cubic centimeters with ammonia free water; or, in other words, fifty cubic centimeters of a solution which contains ammonic chloride equivalent to one one-hundredth of a milligram of nitrogen.

The tubes employed were those used in the regular nesslerization. They are of colorless glass, ten inches in extreme length and eight inches high to the mark; bottoms are ground smooth and polished.

Total Solids.—For the determination of the total solids, to two hundred and fifty cubic centimeters of the water, five cubic centimeters of a four-tenths per cent sodium carbonate solution were added, and the liquid boiled to dryness in a platinum dish upon the water bath. When dry, the dish and its contents were placed in the air bath, kept at 180 degrees Centigrade, and were heated until the weight was essentially constant, the time of heating ordinarily being one hour.

Loss on Ignition.—The loss on ignition was determined by heating the residue from evaporation in a radiator to low redness. No attempt was made to entirely burn away all carbonaceous matter contained in the residue, and the residues frequently were quite dark in appearancee from presence of minute particles of carbon.


Chlorine.—The chlorine determinations were made by the ordinary standard process by titration with silver nitrate solution. The indicator employed was potassium chromate of five per cent strength, one cubic centimeter of the solution being used with each lot of water titrated.

The end point was in all cases determined by close comparison with a blank. Fifty cubic centimeters of water were ordinarily taken for the determination, but in case there was reason to suppose that the water contained less than ten parts of chlorine per million, a larger quantity was used. In all such cases two hundred and fifty cubic centimeters or more were employed. To the measured water, five cubic centimeters of sodium carbonate solution (four grams Na₂CO₃ to the liter) were added and the liquid boiled down, the final volume being brought to fifty cubic centimeters before the determination was made.

Oxygen Consumed.—For the determination of the oxygen consumed, one hundred cubic centimeters of the water were used. To this, two cubic centimeters of pure concentrated sulphuric acid were added, and then ten cubic centimeters of the standard potassium permanganate solution, the strength used being such that one cubic centimeter was equivalent to one-tenth milligram of oxygen.

Every effort was made to have all determinations of oxygen consumed conducted with the greatest uniformity throughout. The heating was effected by immersing the flasks containing the mixture in the boiling water of a large bath. The time of heating was made exactly thirty minutes in each case, and the determination was carried out immediately upon removing the flasks with their contents from the bath.

Nitrogen as Free or Saline Ammonia.—Two hundred and fifty cubic centimeters of the water sample were diluted to 500 cubic centimeters with nitrogen free distilled water. Five cubic centimeters of a twenty per cent sodium carbonate solution were added and the liquid distilled from round-bottomed Jena glass flasks of capacity 900 cubic centimeters. The flasks were supported upon asbestos rings and heated by direct application of the flame. Connection with the condenser was



made by means of the modified form of Reitmair and Stütsen safety bulb designed by Hopkins.

We at first employed condensing tubes of block tin, three-eighths of an inch internal diameter, with a cooling surface twenty inches in length, but for most of the work we employed tubes of aluminium of the same dimensions. The tubes pass through a galvanized iron tank, through which a constant current of cold water is kept flowing. Before each determination the entire apparatus was thoroughly steamed until free of ammonia.

As all of the river waters contained considerable nitrogen as free ammonia, the distillate was collected in flasks of 200 cubic centimeters capacity, the distillation being continued until the flasks were full to the mark, and at such rate that from thirty to forty minutes lapsed between the appearance of the first drops of distillate and the completion of the distillation of the 200 cubic centimeters. The distillates thus caught were thoroughly mixed, the flasks stoppered and set aside for the nesslerization.

Nitrogen as Albuminoid Ammonia.—The determination of the albuminoid ammonia was made in the usual manner upon the residue remaining from the determination of free ammonia. The apparatus and contents having been somewhat cooled, fifty cubic centimeters of the usual alkaline permanganate solution were added through a funnel, the flask immediately connected again with the still, and distillation proceeded with at the same rate as in the determination of the free or saline ammonia. The distillate was caught in flasks of 200 cubic centimeters capacity, and the distillation was considered complete when two hundred cubic centimeters had come over.


Nesslerization.—In conducting the nesslerization, care was always taken that the distillates and the standards were of the same temperature. Commonly those distillates obtained in the afternoon were allowed to stand in a cool place until the next morning before proceeding with the determination.

The ammonium chloride solution used for the comparisons was of such strength that one cubic centimeter contained ammonium chloride corresponding to one one-hundredth of a milligram of nitrogen.

The eighteen standards used in the comparison were made of the lowing strengths. i. e., the quantities of standard ammonium chloride solution used were .05, .1, .2, .4, .6, .8, 1. cubic centimeter, 1.2, 1.4, 1.6, 1.8, 2., 2.5, 3., 3.5, 4., 4.5, 5. cubic centimeters.

In nesslerizing, one cubic centimeter of the nessler solution was added to the contents of each nessler tube and the mixture allowed to stand twenty minutes for the development of the full color. The reagent was always added to the samples and the standards simultaneously, and the readings were all taken within one hour of the time when the reagent was added.

The comparisons have been greatly facilitated by the use of a black wooden box or camera, which cuts out all side lights, the tubes being illuminated from the bottom by means of a mirror reflecting light from



the northern sky, the cross-section of the tubes being brought to the eye by another mirror placed just above the tubes.

Total Organic Nitrogen.—The total organic nitrogen in the original water and in the filtered water was determined by the Kjeldahl process, as follows: Two hundred and fifty cubic centimeters of the water were diluted with 250 cubic centimeters of nitrogen free distilled water; then five cubic centimeters of twenty per cent sodium carbonate solution were added and the mixture distilled as usual for the removal of all free ammonia, the distillation being pushed to precisely the same point as that reached in the distilling over of free ammonia for the determination of free or saline ammonia and albuminoid ammonia. To the residue in the flask ten or twenty cubic centimeters of pure nitrogen free sulphuric acid were added and the solutions heated under the proper precautions until the water was all expelled and the organic matters completely destroyed.

After cooling, 250 cubic centimeters of ammonia free water were added and then an excess (usually about fifty cubic centimeters) of strong nitrogen free sodium hydroxide solution. The flask was immediately connected with the condenser, the contents mixed by thorough shaking and the distillation, which was conducted at first very slowly, was continued until 200 cubic centimeters were distilled over. After thorough mixing, an aliquot portion of this distillate was employed for nesslerization in the ordinary manner.

Nitrogen as Nitrites.—Fifty cubic centimeters of the water were placed in a nessler tube, one cubic centimeter of an acid solution of naphthylamine hydrochloride (eight grains of naphthylamine, eight cubic centimeters of strong hydrochloric acid and sufficient water to make one liter of solution) and one cubic centimeter of a saturated solution of sulphanilic acid in water containing five per cent of strong hydrochloric acid were added, and the mixture allowed to stand for one hour.

Simultaneously with the addition of the reagents to the water which was being examined, the same quantities of reagents were added to a series of solutions which contained accurately known quantities of pure sodium nitrite. If a color appeared in the water sample in course of twenty minutes to one hour after addition of the reagents, it was matched with some one of the tints produced in the series of standards, and the quantity of nitrite contained in the original water was regarded as being the same as that contained in the standard which produced the same tint. If no color developed in the course of an hour, the water was considered free of nitrites.

Many of the river waters examined contained so much nitrites that the color developed in the undiluted sample was too deep for accurate comparison. In such cases quantities of from one to ten cubic centimeters of the sample were diluted to fifty cubic centimeters with nitrogen free water before adding the reagents.

Standard solution of sodium nitrite was prepared from pure silver nitrite by reaction with sodium chloride, and for convenience in making up the standards was made in two strengths, one solution containing in one cubic centimeter the equivalent of .005 milligram of nitrogen, the other .0005 milligram of nitrogen.

Waters which were turbid or deeply colored were clarified and decolorized by treatment with aluminium hydroxide and filtration before testing for nitrites. The comparison of tints was made in the camera described under Nesslerization.

Nitrogen as Nitrates.—One hundred cubic centimeters of the water were treated with two cubic centimeters of nitrogen free sodium hydroxide solution of thirty-three per cent strength, and the solution boiled down rapidly to about one-third its volume to remove free ammonia; the liquid was then diluted to one hundred cubic centimeters with nitrogen free distilled water, cooled and put into a large test tube.

One gram of aluminium in the form of a strip of thin foil was introduced and the tube with its contents placed in a water bath, the temperature of which was kept at about 25 degrees Centigrade, where it was allowed to remain over night. The reductions to ammonia were ordinarily complete when the examinations were continued next morning.

The contents of the reduction tube, including such portion of the aluminium foil as remained, were transferred to a distillation flask, 250 cubic centimeters of nitrogen free water being used to wash out the tube and dilute the liquid. The distillation and subsequent nesslerization were conducted precisely as for the determination of free or saline ammonia.

Dissolved Oxygen.—For the determination of dissolved oxygen, we have found the method of Albert Levy most satisfactory.* The process involves the use of a special pipette, with glass cock at each end. The capacity of the pipettes which we have used is exactly 107 c.c. The reagents employed consist of a solution of 150 grams of caustic potash in a liter of water, a solution of 22 grams of ammonio ferrous sulphate in a liter of water, a fifty per cent solution of sulphuric acid and a standard solution of potassium permanganate of such strength that one cubic centimeter is exactly equivalent to one-tenth of a milligram of oxygen.

The method of procedure is as follows: The pipettes are filled with the water either by immersing in the stream itself or by use of a rubber syringe. Then two cubic centimeters of the caustic alkali solution are put into the funnel at the top, and, by careful manipulation of the two cocks, are allowed to enter and mix with the water without admitting air. The funnel is then rinsed out and five cubic centimeters of the ammonio ferrous sulphate solution introduced into the funnel and then into the pipette by the same manipulation as before. The water run out of the pipette at the bottom as the reagents are admitted at the top is caught in the beaker in which the subsequent titration with permanganate is to be effected and which already contains two cubic centimeters of fifty per cent sulphuric acid.

It is assumed that the alkali and the iron solution in entering the pipette displace their own volume of the water, and with careful manipulation this undoubtedly is essentially effected, so that we may assume that

^{*}This method is as described in the Aunuaire de L'Observatoire de Mont-Souris, for 1883 and subsequent years.

within the pipette there remain 100 cubic centimeters of the original water with the seven cubic centimeters of the reagents.

The mixing of the liquids within the pipette is effected by shaking the pipette with an eccentric rotatory motion. In the course of a few minutes the action is completed, and from the color of the precipitate one may gather an idea as to the relative amount of oxygen contained in the solution; that is, if the water is about saturated, the precipitate is apt to show a somewhat brownish color, due to the ferric hydroxide, while if the quantity of oxygen is very small the precipitate is likely to be black, showing the preponderance of the ferrous hydroxide in the precipitate.

After a few minutes, when the action is thought to be complete, five cubic centimeters of sulphuric acid are introduced into the funnel, and, the cock between the funnel and the pipette being opened, the sulphuric acid, by reason of its greater gravity, passes from the funnel down into the interior, and mixing with the liquid dissolves the hydroxides of iron and renders the entire liquid acid.


When this reaction is complete, as shown by the clearing up of the solution, the contents of the pipette are run into the beaker, and the excess of ferrous salt determined by titration with the standard permanganate solution. A blank is run upon 107 cubic centimeters of the original water for every determination that is made, this being easily done while the reactions are taking place within the pipette.

In running the blank, 107 cubic centimeters of the water are measured into a beaker, then seven cubic centimeters of the sulphuric acid are added and the liquid mixed; after this the caustic potash, two cubic centimeters, is added, and finally precisely five cubic centimeters of the ferrous sulphate solution. Then the titration is effected as in the actual determination. The difference between the two readings—i. e., that of the blank and that of the direct determination—represents the quantity of dissolved oxygen in one hundred cubic centimeters of the water.

We have found the method of Levy more convenient than the Winkler method. Its advantages appear to us to be mainly due to the fact that the blank to accompany each determination is so easily made; whereas, with the Winkler method, the determination of the blank, which with the river waters concerned in these investigations is necessary for every sample examined, entails so much labor as to limit the applicability of the method.

As it has not been practicable for us to make all of the oxygen determinations upon the spot, we have had a great many samples of water shipped from the river to the laboratory. The determinations of the oxygen in these are, in most instances, made within twenty-four hours of the time of collection, but in that length of time the dissolved oxygen is found in most cases to diminish considerably in amount.

The waters of the Illinois River and its tributaries and those of the Mississippi contain a great deal of organic matter which is easily susceptible to the influence of dissolved oxygen. We have found, however, that it is perfectly practicable to treat the water samples with a



little mercuric chloride and thus prevent such reactions as result in the disappearance and consequent diminution of the dissolved oxygen; so that it has been practicable for us in the laboratory to make the comparison of the original quantity and also of the staying qualities of the dissolved oxygen in the water.

With a set of samples treated with a few drops of saturated solution of mercuric chloride we get a result which is essentially the same as that shown by the determination on the spot, while with the other set of samples which have been shipped in the original condition—i. e., merely in bottles which are entirely filled, but which have not been treated with mercuric chloride—it is found that the dissolved oxygen is considerably less in amount. The difference between the two is a rough indication of the condition of the water with respect to content of dissolved oxygen and content of such impurities as easily cause the disappearance of dissolved oxygen.

Our comparisons of the Winkler method with the Levy method show that the latter method gives somewhat higher result, but the differences are ordinarily very slight, and for comparative results the Levy method is so superior in economy of time and labor that of late we have used it almost exclusively.

CHEMICAL EXAMINATIONS.

Our sanitary chemical analyses have included fourteen different quantitative determinations upon each sample of water, the results of which appear in Tables Nos. I to 80, but for the present we need consider only those substances which are generally regarded as of the greatest significance in indicating the relative condition of the waters of streams—namely, chlorine, oxygen consumed, free ammonia, albuminoid ammonia, organic nitrogen and nitrites.

As is well known, and as also appears from the data of our examinations, the water of the canal at Bridgeport is exceedingly foul, being charged with all sorts of refuse matters and sewage. During the period May 1, 1899, to December 31, 1899 (for details see Tables 1-38 and the averages of Table A, page ...), there was comparatively little change in the character of the canal water as it flowed from Bridgeport to Lockport, i. e., the organic matters were present in practically the same quantity at Lockport as they were at Bridgeport.

The effect of the dilution of the canal water by mixing with the waters of the Desplaines River at Joliet was to diminish the proportions of the various constituents and render it possible for the putrefactive decomposition to proceed more actively.

At Morris, the first point upon the Illinois River proper, we found considerably smaller proportions of five of the significant substances above referred to, due largely to the dilution of the water of the Desplaines River by mixing with the waters of the Kankakee, nine miles above Morris; but the slight increase in nitrites shows that the second stage in oxidation is beginning. Continuing down the course of the

stream, the five appear in diminished quantity at Ottawa, while the nitrites have enormously increased, which fact shows that at this point the purifying oxidation caused by the action of bacteria is in full course and at the maximum.

Below Ottawa, the oxygen consumed and the albuminoid ammonia and tota, organic nitrogen increase as we approach La Salle, while the free ammonia and the nitrites are diminishing, partly through dilution with the waters of the Fox River, which enters at Ottawa, partly by continuance of the natural oxidation processes. The increase in the albuminoid ammonia and organic nitrogen continues as we proceed down the river to Henry, partly from fresh accessions of sewage, but also by reason of the growth of grosser vegetable organisms than bacteria, the plankton, which now begins to increase enormously, but all the constituents now under consideration diminish on the way to Averyville.

At Averyville each of the various constituents, except the nitrites, which, however, are not present in considerable proportions until the second stage of oxidation begins vigorously below Morris, has now reached the smallest proportion thus far found. The station at Averyville is about three miles above the Peoria city hall.

The effect of Peoria's contribution of sewage and refuse is seen in the increase of each one of the significant substances between Averyville, just above, and Wesley City, just below. From this vicinity there is a continuous diminution in the quantity of the various constituents until we reach the mouth of the river at Grafton, although large quantities of refuse matters enter the river at Pekin and cause an increase in quantity of oxygen consumed and free ammonia, etc., at that point.

The study of the numerous data of our analyses shows that there is a very considerable purification of the waters of the Illinois River as they flow slowly to the vicinity of Peoria, and that again below Peoria the purification proceeds to such an extent that the water discharged from the Illinois River into the Mississippi at Grafton contains less organic matters than does the water of the Mississippi River itself, and less than is contained in the water of the Missouri near its mouth.

The facts referred to are exhibited by means of the curves of Plate No. 1, page 45, which presents most of the more significant of the averages of Table A, pages i-x. In considering the evidence thus presented, the peculiar significance of the several constituents must be kept in mind.

Chlorine.—This substance is contained in the waters in combination with various basic elements, but chiefly in the form of sodium chloride or common salt. Chlorides occur in very small quantity in ordinary soils, and hence are present in small proportions in the waters of streams in their natural condition. Most animal matters contain more or less chlorides, and chlorides are constant and considerable constituents of sewage.

The only natural causes of change in the proportion of chlorine, aside from such slight concentration as may result from long continued hot weather during the season of minimum flow, would be its decrease by dilution with waters which contain less, or increase by mixing with

waters and other liquids, as sewage, etc., which contain more. The diminution between Lockport and Joliet was due to dilution with the waters of the Desplaines River, which water during this period contained but about one-fifteenth as much chlorine as did the canal water.


The water of the Kankakee, uniting with the Desplaines above Morris, contained 3.9 parts of chlorine per million, and it was the addition of this water, as well as the water of smaller rivers and creeks, which brought about a very notable decrease in the proportion of chlorine between Joliet and Morris. The diminution in chlorine continued until Peoria (Averyville) was reached, resulting from the dilution of the river water with the water of smaller streams, and also undoubtedly by the underground flow to the river from the country lying on either side. This underground flow is known to be considerable along the middle stretch of the Illinois River between La Salle and Beardstown, and particularly between Peoria and Havana, and in times of low water may have quite an appreciable effect in diluting the waters of the stream, an effect which, of course, was more notable before the flow and the initial dilution were increased by the inflow of lake water secured by the opening of the Drainage Channel in January, 1900.

The increase in chlorine between Averyville and Wesley City resulted from the contribution of Peoria's sewage to the river, and in small part was probably due to the inflow of the water of certain saline artesian wells and springs. From this point to the mouth of the river there was a gradual but considerable diminution of the chlorine.

Oxygen Consumed is the quantity of oxygen required for the burning of the organic matters contained in the water, or rather the quantity of oxygen required for oxidation of the readily affected organic matters contained in the water; for, as is well known, not all of the organic matters are affected, i. e., completely oxidized by the chemical process employed. It is evident from the curves that the organic matters which at Lockport required a great deal of oxygen were so affected by dilution and natural oxidation as to appear in much smaller proportions at Morris and Ottawa. A notable increase occurs between Ottawa and La Salle, but the organic matters then diminish until we reach Averyville. There is again an increase until we pass Pekin, and from there to the mouth of the river there is a gradual but definite diminution.

Free Ammonia or Saline Ammonia, as it is sometimes designated, is contained in comparatively great quantity in fresh sewage or sewage which is in the first stages of putrefaction. The rapid decrease in quantity of free ammonia between Joliet and Morris results mainly from dilution with the waters of the Kankakee, but between Morris and Ottawa and from there on down to the mouth, it is chiefly due to the natural exidation, but doubtless is partly due to assimilation by the plant life which is abundant from Ottawa down.

The Nitrites contained in surface waters are products of the action of bacteria upon nitrogenous organic matters and constitute the second step in the series of decompositions which convert these organic matters finally into the comparatively unobjectionable mineral matters, which are




chiefly carbonic acid and nitrates. The comparatively small proportion of nitrites in the upper river to the neighborhood of Morris shows that this second stage of bacterial purification is to this point but beginning, but the great increase below Morris witnesses to the fact that from here on this highly important feature of nature's purification processes is efficiently active. The gradual diminution of nitrites to Averyville indicates a gradual progressive exhaustion of the food supply of the bacteria which are responsible for this oxidation, and thus serves as evidence that the purification is proceeding. The continuous decrease in nitrites from Pekin to Grafton clearly indicates the great improvement in the sanitary condition of the water during its passage from Pekin to the point of discharge into the Mississippi River.

The Total Organic Nitrogen is one of the most important data, inasmuch as the nitrogenous organic matters are the most significant and constant constituents of sewage, and include all sorts of animal refuse, tissues, etc., as well as constituting essential parts of all living things, great or microscopic, both vegetable and animal. It is substances of this class in particular which serve as the nutrients upon which bacteria thrive and multiply.

The Albuminoid Ammonia varies in conformity with the total organic nitrogen, though the quantity is but half that of the latter. Its significance is similar to that of the total organic nitrogen, but the total organic nitrogen represents all of the nitrogen that is contained in the organic matters; that is, as constituting part of the tissues living or dead, as well as various products of the vital activities of living organisms, while the albuminoid ammonia represents only the nitrogen of such part (about one-half) of these substances as is comparatively easily affected by the chemicals and process used in the albuminoid ammonia determination.

COMPARISON OF THE ILLINOIS WITH ITS TRIBUTARIES AND WITH THE MISSISSIPPI AND MISSOURI.

In Groups 2 and 3 of Plate No. 1, page 45, the various columns show the comparative proportions of total organic nitrogen, albuminoid ammonia and free ammonia contained in, and oxygen consumed by, the waters of the more important tributaries of the Illinois River, those of the Mississippi River above the mouth of the Illinois and those of the Missouri just above its mouth. It appears that during this period the waters of the Illinois at its mouth at Grafton contained more free ammonia than those of the Mississippi and the Missouri, and considerably more than those of its tributaries except the Big Vermilion (the Illinois and Michigan Canal is not here considered). On the other hand, the columns show that the waters of the Mississippi and the Missouri contained considerably more total organic nitrogen and albuminoid matters than did the waters of the Illinois, while the oxygen consumed was nearly twice as great. In these respects, too, the columns show that the Illinois compares favorably with its tributaries.



The Mississippi at Alton.—The curves of Group 4, Plate No. 1, referring to the cross-section of the Mississippi at Alton, represent the averages of analyses of samples of water taken at five points across the Mississippi at Alton, about 17 miles below the mouth of the Illinois.

It is evident from the consideration of the data (Tables 23-27 and A), and is graphically shown by the curves of Plate I, that the waters of the Illinois do not mix thoroughly with the waters of the Mississippi, but chiefly pass down close to the Illinois shore, and that the quantities of organic matters contained in the Mississippi River waters are greater than the quantities of the same substances contained in the Illinois River waters, the chlorine and ammonia, on the other hand, being higher in the Illinois River than in the Mississippi at Alton or at Grafton.

The Mississippi at "Chain of Rocks."—The waters of the Missouri do not average so high in chlorine as do the waters of the Illinois, but they contain an enormous amount of suspended matters, and notably greater quantities of various objectionable organic matters. From the data of Tables 28-31 and Table A, exhibited in Group 5, Plate No. 1, it would seem that at the "Chain of Rocks" the waters passing down the west bank and beside the intake tower consist mainly of Missouri River water, and contain much larger quantities of organic matters than do the waters of the Illinois.

Further Comparisons.—The consideration of the analyses of the individual samples shows that the proportions of the nitrogenous organic matters never, during the given period, were so great in the Illinois River as they were in the Missouri, and that the minimum quantity of these matters in the Mississippi and in the Missouri never reached quite the low point that was observed in the water of the Illinois River. The general conclusion from the consideration of these data which, however, it must be remembered, cover simply the period from the beginning of May, 1899, to December 31, 1899, is that the water discharged from the Illinois River is in these respects in purer condition than the water of the Mississippi River or the water of the Missouri River, and that it is in most respects quite similar to the waters of the larger tributaries of the Illinois. The table below gives the specific data:

EFFECT OF OPENING THE DRAINAGE CHANNEL.

The effect of the dilution of the sewage of Chicago with lake water before discharging it into the Desplaines and the Illinois is shown by the analyses made during the year 1900. The data of these analyses appear in tables 39 to 80 inclusively, and the averages are brought together in Table B, pages i-x. The two plates upon pages 49 and 53, Nos. II and III respectively, present the most important averages graphically, and in the light of the explanatory discussion above, page ..., with reference to the earlier year, may readily be understood. They are not especially discussed here, for the reason that the analyses made during 1900 covered the period January 1 to October 1, while those of 1899 covered a different period, namely, that from May 1 to December 31.

It is a well-known fact that the waters of streams vary greatly at different seasons, both as to the volume of water and the content and nature of substances carried by the water. These seasonal variations are particularly important in the cases of some of the streams of the Mississippi Valley, and especially so for the Illinois, the Missouri and the Mississippi. However, inasmuch as the two periods overlap in the summer, it is possible for us to institute a comparison for the most important season of the year—the four months, June, July, August and September—during which period the most important physical features to be borne in mind are the comparatively small volume of water or flow and the comparatively high temperature, not only of the atmosphere, but of the water itself, features which are especially important with reference to the Illinois.

For the purpose of making this comparison I have prepared averages of the data of the analyses made during June, July, August and September, 1899, or before the opening of the Sanitary Channel, and similar averages of the data obtained from the analyses made during the same four months in 1900, or after the opening of the Sanitary Channel. The data appear in detail in the Tables 1-80, and the particular averages in question appear in Tables C and D respectively, upon pages i-x. In order to facilitate the comparison, the plates, Nos. 4 to 15, in which the various curves and columns represent the figures of the tables in a form which facilitates a realization of their significance, have been prepared.

In order to clearly understand the situation and the full significance of the data, reference should be frequently made to the statement of the reasons for selecting the various points of collection (pages 47 to 54) and to the map upon page 8, which shows the relative situations of these points.

In each plate there are two sets of curves or columns, one set representing the data for the summer of 1899, and the other representing the data for the summer of 1900. By following the various proportions of the curves it is made evident that the conditions in the river have changed considerably since the opening of the Drainage Channel.

The portions of the curves and the sets of columns at the left hand of each plate, representing the conditions prevailing between Chicago and Lockport, are, of course, modified more greatly than the other parts of the curves, for the reason that until January, 1900, it was the old, or Illinois and Michigan Canal only which carried the discharge of Chicago sewage into the Illinois River Valley, while since the opening of the Drainage Channel the Sanitary Canal itself carries, not merely undiluted sewage and stockyards drainage, but a very large proportion of diluting lake water.

Inasmuch as the Illinois and Michigan Canal is still mainly fed from a fork of the South Branch, which is not yet well flushed with lake water, we find that the proportions of the various constituents are still excessive in the waters of the old, or Illinois and Michigan, Canal, both at Bridgeport and at Lockport. It is to be remembered that the discharge through the Illinois and Michigan Canal is now much less than it was formerly, for since the opening of the Drainage Channel the old canal is used chiefly for transportation, not particularly as a sewer, and the pumps at Bridgeport are supposed to raise only so much fluid as is required to maintain a serviceable depth for commerce. The average flow in the Illinois and Michigan Canal prior to 1900 was from 45,000 to 50,000 cubic feet per minute. In 1900 it was 26,700 and in 1901, 13,800 cubic feet per minute.


The figures of Tables Nos. 40 and 41 reveal occasional variations of considerable magnitude. These variations are chiefly due to the fact that the pumps at Bridgeport, which feed the Illinois and Michigan Canal, are situated but a few rods from the mouth of the South Fork, and that the diluted sewage, which, on its way through the South Branch to the Drainage Channel, passes the mouth of the South Fork, sometimes enters a short distance within the mouth and is thence pumped into the old canal. The short duration of such occasional conditions is accountable for the fact that the variations are less frequent and less considerable at Lockport, 29 miles below.

At Joliet, where the discharges through the two canals and that from the Upper Desplaines River have come together, there is frequent and considerable variation in the proportions of organic matters and other impurities contained in the waters of the Desplaines River, partly, perhaps, in consequence of variations in the flow of water from the Illinois and Michigan Canal through use of the locks above, partly from occasional closing of the gates at the beartrap dam, thus shutting off or reducing the flow of the much more dilute sewage from the Drainage Channel.

The accumulation of sewage in the Chicago River which follows a stoppage or reduction of the flow through the Drainage Channel is another cause of variation, for when the gates are again opened the sewage discharged is for a time necessarily less diluted than it would otherwise have been. The variations at Joliet result in large part also from incomplete commingling of the waters, for the discharge from the old canal, entering by the east bank, generally passes down beside that bank for some considerable distance, but at times, when the flow from the Drainage Channel is reduced, extends further out into the stream, or possibly entirely across the river bed.

Comparison for Chlorine.—Upon Plate 4, page 57, there are shown, first, general curves representing the proportions of chlorine and of nitrogen as albuminoid ammonia in the waters of the stream at different points between Chicago and the mouth of the river at Grafton.

Comparison of the two curves representing chlorine shows a much more considerable diminution in the proportions of chlorine contained in the water flowing through the bed of the Desplaines River at Joliet in 1900 than in 1899. This is obviously due to the dilution of Chicago sewage with lake water, which passes through the Drainage Channel. The continuous diminution in content of chlorine between Bridgeport (in both years, following the Illinois and Michigan Canal as far as



Joliet) and the mouth of the river at Grafton is, of course, due merely to dilution. (See discussion of Chlorine, page 72.)

The average quantity of chlorine contained in the natural surface waters of the regions in question has ranged from two to three parts, as in the water of Lake Michigan, to as much as five or six parts, which is frequently found in the waters of such streams as the Fox, the Kankakee and the Sangamon.


The waters which pass through the Drainage Channel contain a considerably smaller proportion of chlorine than is contained in the waters passing through the Desplaines and the Illinois River valleys, for the reason already noted, that the sewage carried by the Illinois and Michigan Canal and discharged by it into the Desplaines River just above Joliet is still very concentrated. Moreover, in and about Lockport and Joliet there are various saline artesian wells, sewers and manufacturing establishments which discharge into the old canal or the river between the collecting point at Lockport and that at Joliet.

From Joliet down the river, the slight fall in the chlorine for 1900, as we reach Morris, and the subsequent apparent increase in the content of chlorine at Ottawa, notwithstanding the inflow of the waters of the Kankakee above Morris, are, in some measure, misleading and untrustworthy, for the mixing of the waters of the two streams is, at these points, still incomplete, as subsequent work has shown.

To the incomplete mixing of the main body of water from the Drainage Channel with the waters of the various tributaries is due many of the variations in both directions which appear in the results for 1900 for the several points between and including Joliet and Ottawa. Various cross-sections made at La Salle, Averyville and Grafton, however, show that at these points the mixing is complete, for, particularly at Averyville and Grafton, which may be looked upon as critical points, determinations of chlorine in samples taken at a number of points across the stream have invariably shown that in these localities there is no difference in the chlorine content of the water anywhere between the two banks.

At the mouth, where the Illinois discharges into the Mississippi River, its water contains less chlorine than is found in it anywhere between that point and Chicago, except in the Sanitary Canal. The proportion contained in the water at the mouth of the river in 1900 is very slightly greater than that contained during the corresponding time in 1899.

Albuminoid Ammonia.—The curves representing the nitrogen as albuminoid ammonia show for 1899 a very rapid diminution in quantity as we pass from Bridgeport down the stream to Ottawa. Beyond Ottawa there is a notable increase, reaching the maximum at Henry, but beyond that point there occurs a considerable diminution to Averyville (Peoria). Below Averyville—i. e., at Wesley City and Pekin—the quantities of nitrogen as albuminoid ammonia were greatly increased as a result of the discharge of sewage and of wastes from distilleries, cattle-sheds, glucose works and such plants at Peoria and Pekin. Below Pekin there is again



a slight but constant diminution until we reach the mouth of the river at Grafton.

The curve for 1900 shows that there are much smaller quantities of albuminoid matters contained in the waters of the Sanitary Canal and in the waters of the Desplaines River at Joliet, and as we proceed down the stream from Joliet there is a decrease which reaches the limit at Ottawa. Beyond Ottawa the albuminoid nitrogen increases until Henry is reached, beyond which point it diminishes to Averyville. A slight increase occurs as we pass from Averyville to Wesley and Pekin and on to Havana.

Beyond Havana there is a gradual but continuous diminution until we reach the mouth of the river at Grafton, and the proporton of albuminoid nitrogen contained in the water discharged from the Illinois River at Grafton is less in 1900 than it was in the year 1899.

COMPARISON OF THE ILLINOIS WITH THE MISSISSIPPI AND THE MISSOURI.

Group 2, on Plate 4, represents, first, the comparison of the chlorine contained in the waters of the Illinois as it discharges into the Mississippi at Grafton for the summer months of 1899 and for the summer months of 1900. As is evident from inspection of the figures and the plate, the difference between the two years is so slight as to be hardly noticeable.

Second—A comparison of the chlorine contained in the Mississippi above the mouth of the Illinois River, for like periods in 1899 and 1900. This shows that the water of the Mississippi River contained a considerably greater proportion of chlorine during the summer of 1900 than it did during the summer of 1899, due, doubtless, to the fact that the water in the Upper Mississippi was at a lower stage during the summer of 1900. See Plate XVII for comparison of the stages of water in the Mississippi River in 1899 and 1900.

Third—A comparison of the chlorine contained in the waters of the Missouri River near the mouth for the same periods. This shows that the waters of the Missouri River contained a much higher proportion of chlorine during the summer of 1900 than during the corresponding season of 1899, but it must be noted that the period covered in 1899 is shorter than that for 1900, as in the earlier year the collections from the Missouri began July 10.

Fourth—Comparison of the content of albuminoid nitrogen in the Illinois River water for 1899 and 1900, referred to in the general statement above, is shown graphically in this group.

Fifth—Comparison of the content of nitrogen as albuminoid ammonia in the water of the Mississippi above the month of the Illinois shows a slightly smaller proportion contained in the water of the Mississippi in 1900 than was contained during the same period of 1899, and shows, further, that there was considerably more albuminoid nitrogen in the waters of the Mississippi River during each of these seasons, viz., 1899 and 1900,

than was contained in the waters of the Illinois River in either of the seasons in question.

Sixth—Comparison of the content of nitrogen as albuminoid ammonia in the waters of the Missouri shows that during the summer of 1900 the proportion was greater than during the summer of 1899, and shows, further, that more than twice as much was contained in the water of the Missouri River in each of these years than was contained in the waters of the Illinois River at Grafton.

CROSS-SECTION' OF THE MISSISSIPPI AT ALTON AND AT MITCHELL OR "CHAIN OF ROCKS."

Group 3, upon Plate 4, represents the content of chlorine and of nitrogen as albuminoid ammonia in the waters of the Mississippi River at four points across the stream at Mitchell, Ill. ("Chain of Rocks"), opposite the intake tower of the St. Louis waterworks. From consideration of the curves it is made evident that the content of chlorine and of albuminoid nitrogen was considerably greater upon the Missouri side of the river than it was upon the Illinois side, and shows, further, that there was a greater difference between the two sides of the stream at the St. Louis pumping station during the season of 1900 than there was during the same period of the preceding year, or the year 1899.

Plate 14 exhibits the differences between the proportions of chlorine at the different cross-section points at the "Chain of Rocks," and also shows the seasonal variations of chlorine at these points. The small circles place the dates and show also the proportions of chlorine.

Upon Plate 5 the proportions of chlorine at the several stations are represented by means of columns, and in addition to the various features discussed above (page 72), the cross-section of the Mississippi at Alton is shown in Group 3. This shows the fact repeatedly referred to, that the Illinois River water hugs the Illinois shore and does not mix very much with the Mississippi for a considerable distance below the mouth at Grafton.

Albuminoid Ammonia, Total and Dissolved.—Plate 6 gives similar representation to the facts concerning albuminoid ammonia, including also that part which is in solution.

Organic Nitrogen, Total and Dissolved.—Plates 7 and 8 present in general similar features to those already referred to in the description of Plates 4 and 6, but the particular substances represented upon Plates 7 and 8 are the total organic nitrogen and the dissolved organic nitrogen. By total organic nitrogen, we mean all the nitrogen contained as a constituent part of organic matters, including living organisms themselves. It includes that which is referred to above as albuminoid nitrogen, or nitrogen as albuminoid ammonia, and in general is from two to three times as much as the latter. (See also page 68.) An inspection of the curves and columns shows variations in the quantity of nitrogen in these forms, which are similar in most respects to those described above with reference to Plates 4 and 7, and the general conclusion there, as to the diminition

of these substances as the waters pass down the Illinois River Valley, applies here.

The dissolved organic nitrogen is, of course, derived from those nitrogenous organic matters which are actually in solution in the water, while the total organic nitrogen includes not only this but also the nitrogen contained in such matters as are suspended in the water. Group 2, Plate 7, and Group 3, Plate 8, include comparisons of the proportions of total organic nitrogen contained in the waters of the Illinois, the Missouri and the Mississippi rivers for the periods in question.

The Illinois River water we find contained a considerably smaller proportion of total organic nitrogen during the summer of 1900 than it did in 1899, and in each of these years the proportion was less than that contained in the waters of the Mississippi, and about one-fourth to one-third that contained in the waters of the Missouri River.


The Mississippi River water was, in this respect, somewhat better during the summer of 1900 than during the summer of 1899, but the Missouri River contained considerably greater proportions of total organic nitrogen during the summer of 1900 than it did during the summer of 1899.

These comparisons give expression to the fact that the nitrogenous organic matters in solution, which are here represented by the term organic nitrogen in solution, are contained in the Illinois River water in greater proportion than in the waters of the Mississippi or in the waters of the Missouri River, but it is noticeable that the difference between these streams has been considerably less during the summer of 1900 than it was during the summer of 1899. In the waters of the Missouri River there was somewhat more in 1900 than in 1899, but the difference is slight. For the Mississippi and the Illinois, however, there are, as represented, considerable differences, indicating a better condition of their waters in this respect during the summer of 1900 than during the summer of 1899. The striking differences between these three streams with respect to the proportions of suspended nitrogenous organic matters becomes clearly manifest upon inspection of the unshaded portions of the columns of Plate 8.

Group 3, of Plate 7, and Group 2, Plate 8, show comparisons of the quantities of total organic nitrogen and of dissolved organic nitrogen in the waters of the Mississippi at four different points across the stream at Mitchell ("Chain of Rocks"). The points are the same as those referred to on Plate 4. It is evident, from inspection of the lines, that the quantities of organic nitrogen in the water have been considerably less all across the stream during the summer of 1900 than they were during the summer of 1899, and that there is even greater difference between the waters passing down the Illinois side and those passing down beside the intake tower during the year 1900 than there was during the year 1899.

The curves and columns representing the dissolved organic nitrogen show less marked differences between the waters at the two sides of the stream, although in both years the proportion is greater on the Illinois side.

At the left hand of Group 3, Plate 7, and in Group 4, Plate 8, we have similar data for a cross-section of the stream at Alton, Ill. It should




be understood that at Alton and at the "Chain of Rocks," throughout the time that the examinations were under way, samples were collected each week or oftener, and that the curves are drawn to represent the averages covering the periods of the four summer months in question. The curves and columns which represent the conditions at Alton show a notable difference between the waters at the two sides of the main stream of the Mississippi River, the total nitrogenous organic matters being present in greater proportion toward the Missouri side, that part which is in solution being present in greater proportion toward the Illinois side, while for nitrogenous matters in suspension the conditions are reversed, the proportion being far greater toward the Missouri side than it is toward the Illinois side of the stream.

Free Ammonia and Nitrites.—Plates 9 and 10 represent the proportion of nitrogen as free or saline ammonia and of nitrogen as nitrites contained in the waters of the Sanitary Canal, the Desplaines River and the Illinois River at various points between Chicago and the mouth of the river at Grafton. After what has been said concerning similar plates presenting other important constituents, but little explanation is needed in order to understand the meaning of the curves of Plate 9 or the columns of Plate 10, with reference to the purification of the stream.

The total organic nitrogen and the albuminoid nitrogen referred to in the description of Plates 4, 6, 7 and 8 represent waste organic matters, which may be derived from either vegetable or animal sources. All vegetable and animal organisms contain as essential constituents substances which in their original condition, and also in certain various stages of decomposition, are classed as nitrogenous organic matters. Such matters are included in feces and urine, in wastes from the household, in wastes from all industries which utilize plant and animal substances or products, as well as in the vegetable substances contained in soils, and which accumulate in swamps and in forests. Those substances which make sewage and animal wastes in general offensive to the senses and dangerous to the health are, as a rule, nitrogenous organic substances, either living organisms or the products or wastes of living things. The natural processes of purification of sewage and similar organic wastes involve the oxidation of these nitrogenous matters. The first stage in the purification process includes such a decomposition of these substances as results in the liberation of a portion of their nitrogen, together with hydrogen in the form of ammonia. This may remain merely dissolved in the water, or it may unite with acids, particularly with the carbonic acid. which in these decompositions of the organic matters may be formed simultaneously with ammonia and appear as salt of ammonia. Ammonia is not ordinarily contained in surface waters in notable quantities, except it be derived from the sources above mentioned. The proportion of free ammonia contained in the water of a stream or in sewage indicates, on the one hand, the relative quantity of refuse matters contained and, on the other hand, shows the stage of decomposition.

Referring to Plates 9 and 10, we may observe that in 1899 a considerable proportion of nitrogen as free ammonia was contained in the water of the Illinois and Michigan Canal, and that, although the propor-



tion contained in the Desplaines River at Joliet was considerably diminished through mixing with the waters of the Upper Desplaines River, yet considerable proportions of free ammonia were found in the river as far down as La Salle, but that the diminution still proceeded as we pass down the river till we find the minimum at Averyville. Below Averyville the free ammonia increased in proportion again, reaching the maximum at Pekin, diminishing then more rapidly and continuously until the mouth of the river at Grafton is reached.

The curve for 1900 shows, of course, the effect of dilution with lake water in the Sanitary Canal, i. e., the proportions of nitrogen as free ammonia never were so great in the water flowing between the upper end of the canal and the points between that and Grafton as they were in the preceding year in the Illinois and Michigan Canal and the Desplaines and the Illinois rivers. Otherwise the curves show similar features, i. e., the very rapid diminution of free ammonia until we reach Averyville shows that the natural decomposition of the organic matters is taking place very vigorously in the upper reaches of the stream; that this action diminishes as we proceed down the stream, the indication being that the diminution of the bacterial action is caused mainly by the exhaustion of the nutrients, i. e., the supply of substances which are being converted by bacteria through ammonia into other innocuous forms of matter, and which may be regarded as the food supply of the bacteria, is more rapidly diminishing.

It is notable that the proportion of free ammonia increases again below Averyville, but afterward diminishes, until we find at the mouth of the river considerably less than elsewhere along the course of the stream, and less during the summer of 1900 than during the summer of 1899.

Upon Plate 9 there are two curves, and upon Plate 11 various columns representing the proportions of nitrogen as nitrites. Nitrites are oxidation products derived from the nitrogenous organic matters above referred to. In the conversion of objectionable and harmful nitrogenous organic matters into innocuous substances, the nitrites mark a stage following that which is indicated by the presence of free ammonia.

The curve for 1899 shows the maximum content of nitrites at Ottawa. From there on considerable diminution occurs until we reach Averyville and Wesley, but beyond that point there is again an increase; we find the maximum at Pekin, beyond which point there is a gradual decrease until we reach the minimum at the mouth of the river at Grafton.

The two maxima indicate the two zones in the river at which the second stage of the natural purifying processes is most vigorously under way.

Referring now to the curve for 1900, it becomes evident that the increase in the proportion of nitrites becomes marked very much nearer the source of the sewage, i. e., since most of the sewage has been diluted before it is discharged into the Desplaines River Valley, those natural decompositions and oxidations of the organic matters which result in the formation of nitrites, and subsequently nitrates, have begun vigorously at a point much nearer Chicago.

The point at which the maximum quantity of nitrites is found during the summer of 1900 is again at Ottawa, as during 1899. From there the curve varies in general in the same direction, but it is notable that the minimum is less than during the earlier season, and that the maximum never reaches the high proportion found during 1899.

Consideration of the curves shows that the zones at which the purifying action takes place with the greatest vigor are in the vicinity of Ottawa and the stretch of river between Pekin and Havana, under both sets of conditions, i. e., those conditions prevailing during the summer of 1899 and those prevailing during the summer of 1900, or both before and after the opening of the Drainage Channel.

Study of the various figures of the tables and the curves of the plates reveals the fact that in each of the summers in question there was less nitrogenous organic matter contained in the water of the Illinois River at Ottawa, just above the mouth of the Fox River, than was contained in the river water at points farther down its course, except at Averyville and at Grafton. This may in part result from incomplete mixing in the upper river, but is undoubtedly due in part to the fact that organic matters of animal origin, such as constitute the chief organic matters of sewage. are more easily susceptible to the putrefactive and oxidizing influences which bring about the purification of waters; while vegetable matters, which constitute the greater part of the organic matters found in the natural waters of swamps and streams, such as are contributed in larger proportion to the Illinois by its tributaries, and by the run-off from its own slopes. are less easily decomposed and persist in the water of the lower river in greater proportion. Moreover, vegetation, particularly of the minuter and floating forms—i. e., the plankton—is very abundant in the waters of the lower stretches of the river from La Salle down. The nitrites and nitrates produced abundantly in the waters of the upper river, through the decompositions of the nitrogenous organic matters of sewage, are in turn utilized by the vegtation, and in part their nitrogen is built up into more stable and less objectionable nitrogenous organic matters, which remain floating or dissolved in the waters, and appear in the analytical results, but must be regarded with much less disfavor and objection than would be accorded a like quantity of nitrogenous matters of animal origin.

Oxygen Consumed.—Upon Plates 10 and 13 are shown curves and columns representing the proportions of oxygen consumed by the water in the Illinois River at the various points as indicated on the other charts. It is obvious that the water in the upper stretches of the river contains a great deal more of such organic matters as are easily susceptible to oxidizing agencies than does the water in the lower stretches of the river.

The diminution in 1899 was very rapid until we reach Ottawa and Averyville. Below Averyville it increases in quantity, owing to the discharge of sewage and wastes into the river at Peoria and at Pekin. Below Pekin there is a continuous diminution in these matters until the mouth of the river is reached. Similar points appear when we consider the total oxygen consumed by the matters in solution in the water, i. e., the filtered water.

The curves for 1900 show variations at different points along the river and make manifest the fact that the proportion of easily affected organic matters in the water is considerably less than during the corresponding season of 1899, though there is not such a great relative diminution along the course of the river.

Group 2 of Plate 12 represents a comparison of the relative quantities of oxygen consumed by the waters of the Illinois River, those of the Mississippi river and those of the Missouri River. It is evident that the waters of the Illinois River contained considerably less readily oxidizable matter in the summer of 1900 than in the summer of 1899. In the case of the Mississippi River water, it is evident that the proportion of oxidizable matters far exceeds that contained in the Illinois River water, and that the proportion is practically the same for the two seasons, viz., 1899 and 1900.

In the case of the Missouri River water, the proportion of oxidizable matters is greater in 1900 than in 1899; in both years it is much greater than the proportion contained in the Mississippi River, and at least twice as great as the proportion contained in the water of the Illinois River. On the other hand, comparison of the proportionate quantities of oxygen required by dissolved organic matters shows that there is less required by the waters of the Missouri River than by the waters of the Mississippi or those of the Illinois River.


In 1899 the quantity required by the water of the Illinois River was considerably greater than in 1900; also by the Mississippi River water more was required during the summer of 1899 than during the summer of 1900; but in each year the quantity required by the Mississippi River water was greater than the quantity required by the Illinois River water.


Group 3 of Plate 12 shows the relative quantities of oxygen consumed by water samples collected at the four points across the stream of the Mississippi River at the "Chain of Rocks," or Mitchell. It is evident that the water along the Missouri shore, and in the vicinity of the intake tower, contains much greater quantities of organic matters than does the water which passes down near the Illinois shore. On the other hand, the dissolved, readily-oxidizable matters contained in the water near the intake tower are less in proportion than those contained in the water near the Illinois shore.

Plate 13 presents similar facts by means of columns.

Considering the evidence represented by the columns in Group 1, and that of the curves of Group 3, of Plate 12, it appears that the water passing down near the Missouri shore and past the intake tower of the St. Louis waterworks, is much more nearly like the water of the Missouri River than it is like that of either the Mississippi River in its upper stretches or the Illinois River in its lower stretches.

Dissolved Oxygen.—The results of our determinations of dissolved oxygen in some four hundred samples of water from the Illinois River have shown that the waters of the Illinois River at all times contain a considerable percentage of the quantity of oxygen which is required for saturation, and that at times it contains much more than the saturation





figure; indeed, occasionally, at certain points, it contains even more than double the saturation quantity. The supersaturation of the water with oxygen, if we may so term it, is due to the liberation of oxygen by the chlorophyl-containing micro-organisms and other minute plants, which are very abundant in the waters of the Illinois at certain seasons of the year; in fact, oxygen-evolving organisms seem to be always present and almost always active in these waters.

Many of the determinations have been made upon the spot, but inasmuch as circumstances rendered it impracticable for us to make very extensive series of determinations in this way, the samples of water were, throughout a considerable portion of the time, shipped to the laboratory and the determinations of dissolved oxygen made there ordinarily in about twenty-four hours after the time of collection, but sometimes not until forty or forty-eight hours had lapsed from the time of collection. These shipped samples gave results which are always less than the saturation figure, except upon a few occasions when the original water was highly supersaturated with oxygen.

Comparison of the results of determinations made upon the spot with those made upon duplicate samples, which were either shipped to the laboratory or kept under similar conditions to those involved in shipping, show that there was in all cases a notable diminution of dissolved oxygen during the time which elapsed between the time of collection and the time of making the examinations. Consequently, it is certain that the actual content of dissolved oxygen in all these waters was in reality greater than the results here recorded. There was but one case in which a greater percentage of dissolved oxygen was found in a sample which had been kept. (See page 65, Averyville.)

Dissolved Oxygen in the Water of the Illinois River at Its Mouth at Grafton.—The series of determinations of dissolved oxygen which appear in the tables upon pages 95a and 95g show that during the period January 4 to June 27, 1900, the percentage of saturation in the Illinois River water ranged from 43.1 to 95.7, the average being 76.47 per cent for the entire series of 141 shipped samples. As is seen in the date columns the determinations were made upon samples shipped to the laboratory and usually about twenty-four hours after collection. The minimum, 43.1 per cent, was found in the case of four samples collected May 16 and examined May 18. In no other case was less than 52.7 per cent found, and, indeed, in but five other samples was less than 61 per cent dissolved oxygen found.

Dissolved Oxygen in the Waters of the Mississippi at Grafton.—A parallel series of determinations of dissolved oxygen in the waters of the Mississippi, just above the mouth of the Illinois, was made upon samples collected at the same time—i. e., within a few minutes—and shipped and examined under similar conditions. The details appear in the tables upon pages 95c and 95e. The degree of saturation in these ranged from 55.78 per cent to 108.84 per cent, the average for the entire series of 127 samples being 82 per cent.

Comparison of the data obtained from these series of shipped samples

shows that the waters of the Illinois contained 93.2 per cent of the quantity of dissolved oxygen found in the waters of the Mississippi.


Analyses made upon the spot (see tables, pages 95c and 95e) show that both the Illinois River water at Grafton and the Mississippi River water at Grafton frequently contain a greater proportion of dissolved oxygen than is required for saturation, but each of them also frequently contains less dissolved oxygen than the saturation figure requires. In general, it is true that the waters of the Illinois do not retain their dissolved oxygen so long as do the waters of the Mississippi—that is, the dissolved oxygen diminishes more quickly and to a greater extent in the water of the Illinois than it does in the water of the Mississippi River, although the difference between the two river waters in this respect is not very great, as may be seen by study of the data of the tables.

The Upper Illinois.—In the Illinois River at La Salle, just above the town, upon August 16, 1899, we found the water 18 inches below the surface just saturated with dissolved oxygen, the actual figures being 100.12 per cent, 100.12 per cent and 100.36 per cent, in samples taken at different points across the stream. A sample of the water taken at the same time, but examined twenty-four hours later, contained but 75.7 per cent of dissolved oxygen, while still another sample collected at the same time as the three above mentioned, but kept forty-one hours, contained but 58 per cent of the amount required for saturation. The evidence would seem to show that although during sunshine oxygen was being evolved by the vegetable life in the water with sufficient rapidity to keep the water saturated, yet when the conditions were changed by placing the sample in the dark, such action ceased, and the oxygen began rapidly to disappear.

Similar conditions were found with respect to the water of the Illinois and Michigan Canal at La Salle. The water of the upper basin at La Salle was, before the opening of the Sanitary Canal, in better condition ordinarily than was the water of the Illinois River at this point, for the reason that the canal above La Salle is fed from the Fox River and other streams. The records with respect to the water in this place, as given in Table 95a, show that the water in its original condition contained 102.42 per cent, while after standing twenty-four hours it was reduced to 62.26 per cent.

Upon August 17 somewhat striking conditions were found at Henry. The samples were taken above the dam at 1:15 p. m. upon a bright day, the temperature of the water, 32 degrees C., being very nearly as high as was that of the air, which in the shade was 34 degrees (?). A sample of the water taken one inch below the surface was found to contain considerably more than twice the amount of oxygen required for saturation, presuming, of course, the saturation figure should be that for an equilibrium between the oxygen of the atmosphere and that of the water.

The water was full of minute vegetable organisms and it had a very distinct yellowish green tint. The temperature of the top inch of water was 32 degrees C. The quantities of dissolved oxygen found by the Levy method were 220.9 per cent and 211.4 per cent respectively. Samples taken at the same time and place, but at 18 inches below the surface, had a temperature of 28 degrees C., and were found, by the Levy method, to



120000	
115000	
1100001	
105000	
000001	
95000	
00006	
85000	
80000	
75000	
70000	
65000	
00009	PLATE XVI
55000	DISCHARGE CURVES, 1900.
20000	IIIINOIS RIVER AT KAMPSVIIIF
45000	SANITARY CHANNEL AT BEAR. TRAP DAM LOCKPORT
40000	DESPLAINES RIVER AT JOILET
35000	One division represents 5000 cubic feet per second.
30000	COTOMINATION OF THE MENT OF THE PROPERTY OF TH
25000	CELANICE OF CHEINSTRY CHIVERSITY OF ILLINOIS
200000	20000
00051	. 15000
V 00001	Minossy River at W
5000 Desplaines River	Sol Round at Boar Transland and
7 14 21 28 4 11 18 25 4 11 18 25 1 19 15 22 6 13 20 27	3 10 17 24 1 6 15 22 29 5 12 19 36 2 9 16 23 7 14 21 28 4 11 18 25 2 9 16 23 1 1 14 21 28 4 11 18 25 2 9 16 23 1 1 14 21 28 4 11 18 25 2 9 16 23 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

contain dissolved oxygen corresponding to 200 per cent of saturation and 208.3 per cent of saturation respectively; while the determination made by the Winkler method showed 208.35 per cent. A sample collected at the same time and kept until the next day, the analysis being made twentyfour hours after the time of collection, during which time the sample was kept in the dark, was found to contain IIO.I per cent of the saturation figure. At the same time and place a collection was also made at two feet from the bottom, or ten feet below the surface. At this depth also the water was found to be supersaturated, but far less so than the water at the surface; it contained 129.1 per cent. The dissolved oxygen in the water from near the bottom disappeared much more rapidly and was reduced far more proportionately than in the water from near the surface, the percentage remaining at the end of twenty-four hours being 47.42. This difference as to retention of dissolved oxygen was undoubtedly due to a difference in either the plankton or the other organic matters contained in the water in these two different strata.

Averyville.—Several visits were made to Averyville, just above Peoria, for the purpose of making determinations upon the spot, and some of these have developed quite interesting results. Thus, upon July 21, 1899, the water, 18 inches below the surface, was found, late in the afternoon, to contain about 171 per cent of the saturation figure of dissolved oxygen, and the water was still supersaturated with oxygen when samples, which were kept in the dark, were again examined eighteen hours and twenty-four hours later, after the lapse of this time the figure being 115 to 117 per cent.

The quantity of dissolved oxygen in samples collected at the same spot the following morning was found to be only 87.8 per cent, but during the morning it increased slightly to 95.6 per cent at 10:10, reaching 111.7 per cent at 12:20 p. m. and 114.3 per cent at 1:10 p. m., at which time the examinations were stopped by the necessity for catching a train. Samples collected at the same time were kept over for two days, and it was found that the quantity of dissolved oxygen had diminished in forty-eight hours from 114 per cent to from 38 to 40 per cent. (See Table on page 95a.)

Upon August 17, late in the afternoon, the quantity of dissolved oxygen found in the water at the depth of 18 inches, at Averyville, was 80.5 per cent, while the water coming from about two feet above the bottom in fourteen feet of water contained 74 per cent of the saturation figure. A sample of the water taken 18 inches below the surface at 6:30 p. m., which was kept until the next day, was found to contain a considerably higher quantity of dissolved oxygen than the sample which was examined at the time of collection. The determination on the spot gave 80.49 per cent, and that made twenty-four hours later gave 97.53 per cent. It may be that this is an error, but it is possible that in this case some more hardy oxygen-evolving organisms than usual were contained in the water, and that during the following day, which was excessively hot, light was not entirely excluded from the sample, or the original content of oxygen may possibly have been quite different, for, as is evident from the records for

the 19th, when this point was again visited, the water, earlier in the afternoon, was supersaturated.


Upon August 19, at 12:30 p. m., a sample taken 18 inches below the surface contained 184 per cent; another taken at the same point, but three hours later, contained slightly more than 200 per cent, and a sample collected at 4 o'clock contained 215 per cent; another collected at 4:25 contained 215.7 per cent, while samples of water collected one and one-half hours later at the same point and under the same conditions, were found to contain 95.6 per cent dissolved oxygen.

During the time that these examinations were made the surface of the water was marked here, there and everywhere with patches of blue-green scum, the size of a man's hand, apparently consisting of masses of filamentous algae, etc. The current was quite rapid, and these patches were about two or three feet apart in various directions. The samples were taken under such conditions as to exclude these floating matters from the sample of water upon which the test was made. The sun was shining very brightly, and the temperature was very high. These examinations were made at the Averyville bridge, where the river is very narrow and the current rapid. Between this point and Henry, about thirty miles above, the river spreads out to a considerable width, forming a series of so-called pools, lakes or lagoons, in which the current is very sluggish. In these broad stretches the conditions of depth, spread of surface, temperature and rate of movement are in the summer season especially favorable for the development of plant life, and the water consequently so teems with green and blue-green organiss that oxygen is evolved in great abundance; in fact, may frequently be seen coming from the floating plants in streams of minute bubbles.

During the middle of the day, when the chemical activity of the sun's rays is at the maximum, the evolution of oxygen naturally will be at the maximum and the water then become supersaturated, but as the sun sinks toward the west the action slackens and the excess of oxygen begins to disappear; partly, perhaps, by diffusion into the atmosphere, but undoubtedly in the main through absorption by the organic matters either living or dead contained in the water.

The water which was found to be supersaturated at Averyville doubtless received its oxygen mainly at some distance above, and the rather sudden drop from 215 per cent to 95.6 per cent was, in all probability, due to the fact that a new body of water, which in the late hours of the afternoon had not been exposed to such vigorous action as prevailed earlier in the day, began to reach the point of collection. This change may also have been partly due to the fact that as evening approached the sun was in part obscured from the particular point at which these examinations were made by dropping below the high wooded bluffs which line the west bank of the river in this vicinity.

Later in the year—i. e., upon October 16—another series of examinations was made at Averyville, but at this time the conditions were entirely different. The quantity of dissolved oxygen reached but 74.5 per cent to 77.5 per cent.



Examinations which were made during January, 1900, gave similar results to those just mentioned for October, 1900. Examinations made in June, 1900, showed figures which were considerably below those obtained in July and August of the preceding year. It is unfortunate that we did not have an opportunity to make determinations in 1900 upon the same dates as in 1899, for in June the season is not sufficiently advanced to bring about the conditions which are commonly encountered in the latter part of July and the middle of August.

The results obtained in October, 1900, were practically identical with those obtained in October, 1899, which would seem to imply that the opening of the Drainage Channel has not materially changed the conditions with respect to dissolved oxygen at this point and for this season of the year.

Dissolved Oxygen Just Below Peoria.—The few examinations made at Wesley City have always shown that the water of the river at this point contains far less than the saturation quantity of dissolved oxygen, a result due, of course, to the fact that so much sewage is discharged into the river about two or three miles above. Somewhat similar conditions were found with respect to the water at Pekin. The effect of bright sunlight is strikingly shown by comparison of the figures obtained upon August 18, in the middle of the morning, with those obtained August 19th late in the afternoon, the figures for the morning being 24 per cent, those for the afternoon averaging about 82 per cent.

Examinations of water collected early in the morning of August 19, at Havana, thirty miles below Pekin, showed but about one-third as much dissolved oxygen as would correspond to the saturation figure, while later in the day determinations by both the Levy and the Winkler methods, made at Copperas Creek dam and at Lancaster Landing, points between Havana and Pekin, showed considerable excess of dissolved oxygen. The weather was hot, 100 degrees to 108 degrees F., in the shade, and there was considerable plankton in the water, much of it appearing to consist of filamentous and other algæ.

Dissolved Oxygen in the Water of the Drainage Channel.—Several determinations were made of the dissolved oxygen in the waters of the Sanitary Canal itself. Upon August 13, 1900, an examination was made of the water in the upper end of the Sanitary Canal at Kedzie Avenue. At 5:15 in the afternoon a sample taken 18 inches below the surface was found to contain 29 per cent, and a little further down the canal—i.e., at the Belt Line bridge—at 5:40 o'clock, it was found to contain 28.2 per cent.

Upon June 29, 1900, at 5 o'clock in the afternoon, this water, just above the beartrap dam at Lockport, was found to contain 10.9 per cent of dissolved oxygen, while at 200 yards below the dam, after the water had been agitated with air in its escape from the higher level of the Drainage Channel to the bed of the Desplaines River below, the aëration had been sufficient to bring the proportion of dissolved oxygen up to 55.5 per cent of the saturation figure. Upon August 14, 1900, similar determinations showed that the water above the dam contained 5.9 per cent, and that below the dam 70 per cent of dissolved oxygen.

At Joliet, examinations made upon June 30, 1900, showed 27.8 per cent of dissolved oxygen just above Dam No. 1, while just below the dam, where the water has fallen twelve or thirteen feet, the percentage of dissolved oxygen was found to be 81.28.

In making these determinations, due care was exercised to prevent any minute bubbles of air being retained by the water which was made the subject of the examination. The entire pipette was filled with the water and was allowed to stand for a few minutes, until such bubbles as were contained in the sample collected in the upper part of the pipette and were allowed to escape.

The results obtained at Lockport and at Joliet show quite strikingly the effect of agitation of the water with air in falling over the dams.

A single determination made upon the water of the Fox River at Ottawa, August 20, showed the water of this tributary to be quite super-saturated; it contained 138.25 per cent of dissolved oxygen. The water of the Illinois and Michigan Canal, just above Channahon, was examined upon August 21 and was found to contain 3 per cent of the saturation figure of dissolved oxygen. At this time there was practically no water flowing in the bed of the Upper Desplaines below Joliet, as at Joliet all of the flow both from the Desplaines River above and from the Illinois and Michigan Canal was carried down the west bank of the river in the Canal itself.

The following tables, designated I to X inclusive, include those prepared by Professors Palmer, Jordan and Burrill.

ARTHUR W. PALMER, Sc. D., Professor of Chemistry, University of Illinois.

SANITARY CANAL, LOCKPORT.

Date of Collection.	Date of Determination.	Depth of Collection.	Mg. per Ll	A ver.	Satura- tion.	marks.
1900 June 29, 5:00 p.m. '' 29, 5:45 '' Aug. 14 '' 11	1900 June 29, 5:00 p.m. 29, 5:45 " Aug. 14 4 14	18" below surface 18" " " 18" " " 18" " "	1. 1. 5.1 5.1 .55 .55 6.55 6.55		55.5 200 yds	, bel, dan above Frap dan
SANITARY CANAL,	Kedzie Avenue.		1			
1900 Aug. 13, 5:15 p.m. " 13, 5:40 "	1900 Aug. 13, 5:15 p.m. 13, 5:40	18" below surface	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	2.75 2.65	29, 28, 2	
DESPLAINES RIVER			<u> </u>	1		
1900 June 30, 5:50 a.m. '' 30, 6:00 ''	June 30, 5:30 a.m. " 30, 6:00 "	18" below surface	2.6 2.6 7.6 7.6	2.6 18°C	27.8 Just al No. 1. Below sluice	dam in
ILLINOIS AND MICI	HIGAN CANAL, JUST	ABOVE CHANNAHA	N			
1899 Ang. 21, 6:30 a.m.	1899 Ang. 21, 6:30 a.m.	18" below surface	.3 .2	.25 26° c	3.05	
1899 Aug. 20, 5:00 p.m.	1899 Aug. 20, 5:00 p.m.	18" below surface	10.2 10.4	10.3 31°c	138.25	
-	ATER, LA SALLE, ABO			<u> </u>	<u> </u>	
" 16, 3:05 " " 16, 4:00 " " 16, 3:30 " " 16, 3:30 "	" 16, 4:00 " " 17, 4:00 " " 18, 9:00 a m.	18" " " " " " " " " " " " " " " " " " "	8.4 8.2 8.2 8.2 8.2 8.25 6.2 6.2 4.6 4.9	8.2 "	100.12 100.12 100.36' 75.70 / Test 58.00 (Ing q)	
	HGAN CANAL, UPPEI	R BASIN, LA SALLI	E.	<u> </u>	1	
1899 Aug. 16, 4:00 p.m. 16, 1:00 '' 16, 4:00 ''	1899 Aug. 16, 4:00 p.m. 17, 4:60 " 18, 10:30 a.m.	18" " "	8.2 8.25 6.3 6.3 5. 5.	8.22 27°c 6.3 ::	102.42 78.45 Kept 24 62.26 40	
Illinois River W	ATER, HENRY, ABOV	е Дам.				
1899 Ang. 17, 1:15 p.m. 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 17, 1:15 1899	1899 Aug. 17, 1:15 p.m. 17, 1:15 " 17, 1:15 " 17, 1:15 " 17, 1:15 " 17, 1:15 " 18, 1:30 " 18, 1:30 "	1" below surface 1" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " "	17.3 17.6 16 7 16.7 15.8 15.8 16.6 16.3 16.46 8.9 8.5 10.1 10.2 3.7 3.8	16.7	129. 10 ing q	r Methoo of keep nalities, of keep
		-			ing q	nalities.
1899 July 21, 4:30 p.m. 21, 5:30 " 21, 5:30 " 21, 6:00 " 21, 6:00 " 21, 6:00 "	1899 July 21, 1:30 p.m.	18" below surface 18" " " " 18" " " 18" " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " 18" " 18" " 18" " 18" " 18" " " 18" " " "	13.2 13.1 13.3 13.2 13.1 13.2 9.1 19.0 8.9 Lost 9.0 9.0	13.25 " 13.3 " 9.05 " 8.9 "	Later Children	of keep
" 22, 850 a.m. " 22, 9:26 " " 22, 10:10 " " 22, 12:20 p.m. " 22, 1:10 " " 22, 1:10 " " 22, 1:10 "	22, 8:50 a.m. 22, 9:26 " 22, 10:10 "	14.	9.0 9.0 6.6 6.8 7.1 7.1 7.4 7.4 8.1 8.9 8.9 8.8 2.9 3. 3.02	9.0 6.7 7.1 7.4 8.65 8.85 2.95	87.85 91.73 95.60 111.75 111.31 38.11 39.02) Test	of keep ualities, of keep nalities,
A ng. 17, 6:30 p.m. 17, 6:30 p.m. 17, 6:30 ° 17, 6:30 °	· 21, 2:00 ·	18" 18" 2' from bottom 2'	2.98 7.0 6.3 8.2 8.4	6.85 6.3 8.3	37.85) Wink 80.49 21.03	der's.

ILLINOIS RIVER, AVERYVILLE.—Continued.

Date of Collection.		Date of Depth of Collection.		Mg. per Liter.			Temp. of Water.	Per cent Satura- tion.	Remarks.		
	Jonection.	Deter	mination.	Сопес	tion.	1	2	Aver.	Te Wa	Per Sat ti	
Aug	19, 3:20 " 19, 4:00 " 19, 4:25 " 19, 5:55 " 16, 8:35 a.m. 16, 9:25 " 16, 11:00 " 16, 2:30 p.m. 16, 3:30 " 16, 4:20 "	Aug. 19 19 19 19 19 19 19 10 10 11 10 11 11 11 11 11 11 11 11 11	6, 5:55 " 8, 8:35 a.m. 9, 9:25 " 6, 11:00 " 6, 2:30 p.m. 6, 3:30 " 6, 4:20 "	18" below 18" " 18" " 18" " 18" " 18" " 18" " 18" " 18" "	surface	14.5 15. 17.2 16.7 7.5 6.8 7. 7. 7.2 6.8	14.3 16. 16.1 16.7 7.3 6.9 7. 7.1 7.1 7.2	14.4 15.5 16.65 16.7 7.4 6.85 7. 7.05 7.05 7.2 6.9		76.17 76.71	
Jan	1900 4, 8:45 a.m. 4, 8:45 " 4, 8:45 " 4, 8:45 " 4, 8:45 " 4, 8:45 " 4, 8:45 " 4, 8:45 " 4, 11:45 " 4, 11:45 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 30, 9:35 " 31, 9:45 " 31, 1:10 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 9:45 " 31, 11:10 " 31, 4:25 " 31, 11:10 a.m. 31, 11:10 " 31, 4:25 " 31, 11:10 a.m. 31, 11:10 " 31, 4:25 "	Jan. 4 " 4 " 5 " 4 " 4 " 5 " 4 " 4 June 30 " 30 " 30 " 30 " 30 " 30 " 30 " 31 " 13 " 13 " 13 " 13 " 13 " 13 " 15 " 15	, 9:00 a.m., 9:35, 9:35, 9:35, 9:35, 9:35, 1:45 p.m., 9:2:40, 3:45, 4:35, 4:35, 9:45,	18" 18" 18" 8' 8' 8' 18" 18" 18" 18" 18" 2' from 18" below 2' from 18" below 2' from 18" below	surface "" "" "" "" bottom surface bottom surface bottom surface "" "" "" "" "" "" "" "" "" "" "" "" ""	11.1 10.9 10.8 11.1 11. 10.6 10.7 10.9 11.2 5.8 6. 6.2 6.5 6.8 6.4 6.7 6.7 6.7 6.8	11.1 10.9 10.7 11. 10.6 10.7 11.1 11.2 5.8 6.1 6.2 6.6 6.8 Lost 7.2 7.4 6.9 7.2 7.4 7.1 7.1 7.1 6.1 6.1 6.1 6.1 6.1 6.1 6.1 6.1 6.1 6	11.1 11.1 10.9 10.75 11.05 11.06 10.7 11.2 5.8 6.05 6.2 6.55 6.7 6.7 6.85 6.7 7.15 7.2 7.1 7.25 7.1 7.2 7.1 7.2 7.1 7.2 7.1 7.2 7.1 7.2 7.1 7.2 7.1 7.1 6.05	1°c 25°c 18.5°c 18.5°c 18.5°c	77.78 76.38 75.33 77.44 77.08 74.28 74.98 77.08 78.48 69.46 72.45 79.36 81.80 83.63 74.34 74.34 74.37 75.77 75.77 75.77 74.00 63.95	
ILLIN	ois River, W	ESLEY C	ITY.								
July Aug.	1899 22, 9:00 a.m. 22, 9:30 ··· 22, 10:00 ··· 18, 7:00 a.m. 18, 7:00 ··· 19, 6:30 p.m.	July 22 22 22 Aug. 18 18 19	7:00 a.m. 7:00 a.m. 7:00 6:30 p.m.	18" " 18" "	surface	2. 7.2 7.5 3.6 2.7 4.2	2.4 7.3 7.2 3.6 2.7 3.5	2.2 7.25 7.35 3.6 2.7 3.85	28°c 24°c 27.5°c	91.77 93.03	stream
Jan.	1900 4, 4:00 p.m. 4, 4:00 " 4, 4:00 " 4, 4:00 "	Jan. 4. 4. 5	900 , 4:00 p.m. , 4:00 '' , 12:40 '' , 12:40 ''	2' below 2' '' 2' '' 2' ''	surface	$ \begin{array}{c} 11.1 \\ 10.7 \\ 10.5 \\ 10.5 \end{array} $	10.4 10.7 10.4 10.4	10.75 10.7 10.45 10.45	1°c	75.33 74.98 73.22 73.22	East Bank. East Bank, kept 21 hours.
Illinois River, Pekin.											
Aug.	1899 18, 10:00 a.m. 18, 10:00 '' 19, 5:30 p.m. 19, 5:30 ''	Aug. 18, 18, 19,	10:00 '' 5:30 p.m.	18" "	surface	2. 2.6 6.4 6.04	1.9 2.6 6.4	1.95 2.6 6.4	26.5°c 30°c	24.04 32.06 84.21 79.74	Midstream.
Illinois River, Lancaster Landing,											
Aug.	1899 19, 4:20 p.m. 19, 4:20 ···	Aug. 19.	899 4:20 p.m. 4:20 °	18" below 18" ''	surface	9.7 9.13	9.8	9.75		128.29 120	Winkler Method
Illinois River, Copperas Creek Dam.											

HLLINOIS RIVER, HAVANA.

Date of Collection.	Date of Determination.	Depth of Collection.		per Li	A ver.	of of ater.	Per cent Satura tion.	Remarks.
		Concention.	1	2	Aver.	Te W.	Sal	
1899 Aug. 18, 3:30 p.m. 19, 6:00 a.m. 19, 6:00 19, 6:00 19, 6:00	19, 6:00	18" below surface 18" " " 18" " " 18" " " 18" " "	$ \begin{array}{c} 3.4 \\ 3.6 \\ 3.1 \end{array} $	3.7 3.5 3.7	3.7 3.45 3.65	31°c	42 89	Winkler Metho
LLINOIS RIVER, G	RAFTON.							
1899 Apr. 29, 10:30 a.m 29, 2:15 p.m 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 29, 3:15 22, 9:00 p.m 23, 4:30 p.m 24, 1:45 p.m 27, 3:30 p.m 27, 3:30 27, 3:30 27, 4:30 27, 4:30 27, 5:00 27, 5:00 27, 5:00 27, 5:00 27, 5:45 27, 8:30 27, 8:30 28, 6:30 28, 6:30 28, 6:30 28, 6:30 28, 6:30 28, 6:30 28, 11:30 28, 11:30 28, 11:30 28, 11:30 28, 4:30 28, 4:30 28, 4:30 29, 4:30 28, 4:30 28, 4:30 28, 4:30 28, 4:30 29, 4:30 29, 4:30 20, 21, 11:40 p.m. 21, 22, 3:40 22, 3:40 23, 4:30 24, 4:30 25, 11:40 p.m. 25, 11:40 p.m. 26, 3:40 27, 28, 4:30 28, 4:30 29, 4:30 29, 4:30 20, 21, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 23, 4:30 24, 1:30 25, 11:40 p.m. 25, 11:40 p.m. 26, 1:30 p.m. 27, 11:40 p.m. 28, 1:30 p.m. 29, 1:30 p.m. 20, 21, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 23, 4:30 24, 1:30 p.m. 25, 4:30 28, 1:30 p.m. 29, 1:30 p.m. 20, 20, 21, 21, 21, 21, 21, 21, 21, 21, 21, 21	1899 Apr. 29, 10:30 a.m. 29, 2:15 p.m. 29, 3:15 June 23, a. m. 23, q. m. 23, q. 30 p.m. 23, 4:30 p.m. 24, 1:45 p.m. 27, 3:30 p.m. 27, 3:30 p.m. 27, 3:30 p.m. 27, 4:00 p.m. 27, 5:45 p.m. 27, 5:45 p.m. 28, 6:30 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 29, 11:40 p.m. 20, 11:40 p.m. 20, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 25, 11:40 p.m. 26, 11:40 p.m. 27, 11:40 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 20, 11:40 p.m. 20, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 22, 11:40 p.m. 23, 11:40 p.m. 24, 11:40 p.m. 25, 11:40 p.m. 26, 11:40 p.m. 27, 11:40 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 20, 11:40 p.m. 20, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 23, 11:40 p.m. 24, 11:45 p.m. 25, 11:40 p.m. 26, 11:40 p.m. 27, 11:40 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 20, 11:40 p.m. 20, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 23, 11:40 p.m. 24, 11:43 p.m. 25, 11:40 p.m. 26, 11:40 p.m. 27, 11:40 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 20, 11:40 p.m. 20, 11:40 p.m. 21, 11:40 p.m. 22, 11:40 p.m. 23, 11:40 p.m. 24, 11:43 p.m. 25, 11:40 p.m. 26, 11:40 p.m. 26, 11:40 p.m. 27, 11:40 p.m. 28, 11:30 p.m. 28, 11:30 p.m. 29, 11:40 p.m. 20,	1' " " " 18" " " 18" " " " 18" " " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" " " 18" "	11.4 11.1 5.7 5.1 5.7 5.5 6.2 8.2 5.8 6.2 6.3 6.3 6.3 6.3 6.3 6.3 6.3 6.3	6.25 9.7	6.25 9.65 7.75 6.35	27° c 28.5° c 28.5° c 28.5° c 31° c	71.78 71.52 69.87 75.95 113.01 112.46 113.01 83.89 137.85 80.25 78.66 106.16 87.67 78.42 78.94 77.76 84.63 81.27 67.34 67.34 67.34 67.34 67.34 67.34 67.34 67.34 83.55 123.03 107.63 107.63 107.63 107.63 107.63 107.63 83.24 68.35 64.47 89.79 85.37 89.11	Winkler's Method. Winkler's Method.
4, 11,00	" 6, 2:20 p.m. " 6, 2:20 " " 6, 11:00 a.m. " 6, 11:00 m. " 18, 12:00 m. " 18, 2:00 p.m. " 18, 2:00 p.m. " 18, 2:00 o." " 25, 10:30 a.m. " 25, 10:30 a.m. " 25, 10:30 " " 31, 10:25 " " 31, 10:25 " " 31, 10:25 " " 14, 8:50 a.m. " 9, 8:50 a.m. " 9, 8:50 a.m. " 11, 8:45 " " 14, 8:45 " " 14, 9:45 " " 16, 8:30 " " 16, 8:30 " " 16, 8:30 " " 16, 9:20 " " 10, 9:20 "	18" 18" below surface 2" from bottom	11. 10.9 10.9 11.1 12. 12. 12. 12.1 10.8 10.9 10.8 10.9 11.3 11.3 11.3 11.3 11.5 11.7 11.5 11.6 11.1 11.8 11.7 11.6 11.8 11.9	11.1	11.05	6.6		

ILLINOIS RIVER, GRAFTON—Continued.

ILLINOIS RIVER. GRAFTON—Concluded.

Date of Collection.	Date of Determination.	Depth of Collection.	Mg. 1	per Liter. 2 Aver.	Temp. of Water.	Remarks.
1900 May 2, 11:30 a.m. " 2, 11:30 " " 9, 11:00 " " 9, 11:00 " " 9, 11:00 " " 16, 11:30 " " 16, 11:30 " " 16, 11:30 " " 30, 11:30 " " 30, 11:30 " " 13, 11:30 " " 13, 11:30 " " 27, 11:30 " " 27, 11:30 " " 27, 11:30 " " 27, 11:30 " " 7, 5:00 p.m. " 7, 5:00 " " 7, 5:45 "	" 4, 10:00 " " 10, 4!45 p.m. " 11, 8:10 a.m. " 10, 5:10 p.m. " 11, 8:35 a.m. " 18, p.m. " 18, p.m. " 18, p.m. " 31, 1:50 p.m. " 31, 1:50 p.m. " 31, 2:50 " " June 1, 10:15 a.m. " 7, 4:30 p.m. " 7, 4:30 " " 14, 4:30 " " 15, 9:00 a.m. " 28, 10:00 " " 28, 10:00 "	2" 18" below surface 18" 2" from' bottom 2" below surface 18" 2" from bottom 2" from bottom 2" from bottom 2" surface 18" 18" below surface 18" 18" below surface 18" 18" below surface 18"	6.3 6.9 6.5 3.7 3.8 3.7 3.8 5.1 5.1 5.2 5.2 5.2 5.3 5.4 7.3 7.2	8.8 8.2 8.2 6.8 6.75 6.3 6.3 6.5 6.5 3.8 3.75 3.6 5.1 5.1 5.1 5.1 5.2 5.2 5.2 5.2 5.2 5.2 5.2 5.2 5.2 5.2	20°c 99 19°c 77°c 44°c 44°c 44°c 46°c 99 19°c	55.75 89.23 72.03 77.23 73.64 99.37 14.91 14.91 15.51 51.08 51.08 51.08 51.08 52.69 52.27 54.75 71.61 77.97 57.24 57.24 57.24 57.87 55.4 94.7
Mississippi River.	Grafton.					
June 24, 3:00 p.m. July 27, 7:30 p.m. 28, 9:15 a.m. 28, 9:15 28, 2:30 p.m. Aug. 25, 10:08 a.m. 25, 2:05 p.m. 25, 2:30 p.m. 25, 2:30 p.m. 26, 2:30 p.m. 27, 2:30 p.m. 28, 1:30 p.m. 28, 1:30 p.m. 28, 1:30 p.m.		18" · · · · · · · · · · · · · · · · · · ·	8.9 7. 8.6 7.09 7.1 8.9 7.3 8.1 13.7 13.1	8.9 8.9 6.8 6.9 8.8 8.7 8.7 8.8 7.4 7.35 8.1 8.05 8.1 8.1 13.7 13.7 13. 13.05 14. 14.05	30°c 11 27°c 9 28°c 10 10 0°c 9	5.9 6.7 1.7 1.7 5.8 1.53
Jan. 4, 1:30 p.m. 4, 1:30 " 4, 1:30 " 4, 1:30 " 4, 1:30 " 4, 1:30 " 17, 1:30 " 17, 1:30 " 17, 1:30 " 17, 1:30 " 21, 2:30 " 24, 2:90 " 24, 2:90 " 24, 2:90 " 28, 12:15 " 28, 12:15 " 28, 12:15 " 28, 12:15 " 28, 12:15 " 12, 2:30 " 7, 2:30 " 7, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 12, 2:30 " 14, 1:00 " 14, 1:00 " 14, 1:00 " 14, 1:00 " 14, 1:30 " 21, 1:30 " 22, 11:30 " Mar. 2, 11:00 a.m. 22, 11:30 " 23, 11:00 " 45, 11:30 " 45, 11:30 " 45, 11:30 " 45, 11:30 " 45, 11:30 " 47, 2:00 p.m. 47, 2:00 p.m. 47, 2:00 p.m.	5, 12:00 m. 6, 8:30 a.m. 6, 10:30 18, 11:00 18, 11:15 18, 11:15 25, 11:00 25, 11:00 25, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 4, 9:15 9, 8:20 a.m. 9, 8:20 a.m. 9, 8:20 14, 9:15 14, 9:15 14, 9:15 15, 9:45 16, 8:50 17, 8:25 20, 5:00 p.m. 21, 8:15 a.m. 23, 11:10 24, 11:10 25, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 31, 11:30 32, 11:10 33, 5:15 p.m. 34, 9:10 35, 9:10 36, 11:30 37, 8:25 a.m. 38, 9:10 38, 9:10 39, 9:10 30, 9:10 31, 11:30 31, 11:30 31, 11:30 31, 11:30 32, 11:30 33, 5:15 p.m. 34, 9:10 35, 9:10 36, 11:30 37, 8:25 a.m.	2' from bottom 18" below surface 2' from bottom 18" below surface 2' from bottom 18" below surface	14.6 14.6 14.6 13.9 12.7 12.6 11.9 12.5 10.6 10.3 9.4 8.2 13. 12.8 13. 12.8 13. 12.8 13. 14.6 15. 16.6 16.3 17. 18. 19. 19. 19. 19. 19. 19. 19. 19	14.1 14.8 16. 16. 15.8 15.7 15.4 14.4 14.8 14.7 14.4 14.5 14.4 14.5 14.4 14.5 14.1 14.35 12.9 12.5 12.6 12.65 12.7 11.7 11.7 11.8 11.9 12.5 12.4 12.45 10.6 10.6 10.4 10.35 9.6 9.5 8.2 8.2 12.7 12.75 12.7 12.85 12.7 12.85 12.7 12.85 12.7 12.85 12.8 12.85 10.9 10.9 9.4 9.35 9.2 9.2 12.5 12.5 12.6 12.6 12.6 12.55 12.6 12.55 12.4	0°C 88 88 88 88 88 88 88 88 88 88 88 88 88	7.41 6.29 3.33

DISSOLVED OXYGEN IN THE WATER OF THE

MISSISSIPPI RIVER, GRAFTON—Contlinued.

	Date of ollection.		Date of ermination.	Depth of Collection.	1 Mg.	per Li	Λ ver.	Temp. of Water.	Per cent Satura- tion.	Remarks.
	1900		1900						-	
ar.	7, 2:00 p.m.		9, 8:45 a.m.	2' from bottom	13.	13.	13.	0°c		
6	7, 2:00 " 9, 1:00 "	6.6	9, 8:45 "	2' " " " 18" below surface	$\frac{13}{12.7}$	$\frac{12.9}{12.7}$	12.95 12.7	66	88.09	
	9, 1:00 "		12, 8:50 a.m.		12.7	12.1	12.05	6.6	81.97	
6.6	9, 1:00 "	6.6	10. 5:30 p.m.	2' from bottom	12.6	12.6	12.6	6.6	85.71	
66	9, 1:00 "	6.6	12, 8:30 a.m.	2' " "	12.	12.	12.	900	81.63	
44	12, 1:00 " 12, 1:00 "	44	13. 5:00 p.m. 14, 8:00 a.m.	18" below surface	$\frac{11.7}{11.2}$	11.8	11.75 11.25	2°c	84.71	
**	12, 1:00 "	4.6	13, 5:20 p.m.	2' from bottom	12.	12.	12.	. 6	86.52	
* *	12, 1:00 "	6.6	14, 8:35 a.m.	2' " " "	11.7	11.7	11.7	40 -	84.35	
66	14, 11:30 " 14, 11:30 "		15, 5:30 p.m. 16, 8:00 a.m.		$\frac{10.5}{10.1}$	10.5	10.5	4°c	79.84	
4.6	14, 11:30 "	6.6	16, 9:00 "	2' from bottom	10.2	10.3	10.25	6.6	77 94	
6.6	14, 11:30 ''		16, 9:00 "	2' " "	10.1	10.	10.05	46	76.42	
66	19, 2:00 " 19, 2:00 "		20, 4:15 p.m. 21, 8:15 a.m.		$\frac{10.5}{10.3}$	10.7	10.6 10.45	0°c	72.11	
	19, 2:00 "		20, 4:45 p.m.	2' from bottom	10.5	10.7	10.45	/	72.11	
6.6	19, 2:00 "	4.6	21, 8:40 a.m.	2' "	10.7	11.	10.85	**	73.81	
66	21, 11:30 a.m.			18" below surface	11.8	12.	11.9:	3°c		
4.6	21, 11:30 " 21, 11:30 "		23, 8:30 a.m. 22, 5:15 p.m.	10	$\frac{10.7}{11.8}$	11.1	$10.9 \\ 11.8$	4.4	80.74	
4.4	23, 1:00 p.m.	1.6	24, 4:50 "	18" below surface	10.4	10 6	10.5	66	77 77	
	23, 1:00 "		26, 8:45 a.m.	2' from bottom	9.4	9.5	9.45	40-	70.00	
66	26, 1:00 " 28, 1:00 "		28, 10:50 " = 28, 10:50 "	18" below surface 2' from bottom	10.6	10.6	$\begin{bmatrix} 10.6 \\ 10.85 \end{bmatrix}$	4°c	80.61	
66	26, 1:00 "		29, 9:00 "	2' " "	$\frac{10.9}{10.2}$	10.8 10.2	10.2	66	77 56	
6.6	28, 1:00 "	6.6	29, 5:00 p.m.		10.9	10.9	10.9	64	82.89	
66	28, 1:00 "		30, 8:20 a.m.	18" " "	10.7	10.6	10.65	46	80.98	
	28, 1:00 " 28, 1:00 "		30, 9:25 '' 30, 9:25 ''	2' from bottom	$\frac{11.1}{10.8}$	11.	11.05	66	84.03	
6.6	30, 1:00 "	April	001 11.00	18" below surface	10.6	10.6	10.6	4.6	80.61	
16	30, 1:00 "	166	2, 11:55 "	2' from bottom	10.9	10.9	10.9	**	82.89	
pril	2, 1:00 p.m. 2, 1:00 "	66		18" below surface 2' from bottom		12.2	12.2	5°c		
6.6	2, 1:00 ··· 2, 1:00 ···	6.6	4, 8:20 a.m. 4, 8:20 "	2' '' of oottom	$\frac{11.5}{11.1}$	11.5 $11.$	11.5		89.70	
6.6	4, 1:00 "	4.6		18" below surface	10.4	10.4	10.4	**	81.12	
	4, 1:00 "	6.6	6, 8:50 a.m.		10.4	10.3	10.35	100	80.73	
	6, 1:00 " 6, 1:00 "		7, 4:50 p.m. 9, 8:30 a.m.	18" below surface	$\begin{array}{c} 9.4 \\ 8.2 \end{array}$	$9.5 \\ 8.2$	$\begin{array}{ c c} 9.45 \\ 8.2 \end{array}$	4°c	71.86 62.35	
44	6, 1:00 "	6.6	7, 5:25 p.m.	2' from bottom	10.5	10.4	10.45	6.6	79.46	
6.6	9, 1:00 "		10, 5:00 "	18" bclow surface	9.5	9.5	9.5	9°c	82.04	
66	$\begin{array}{cccccccccccccccccccccccccccccccccccc$		11, 8:00 a.m.		9.4	9.5	9.45	66	81.61	
	9, 1:00 "		11, 9:10 " 11, 9:10 "	2' from bottom	$\begin{array}{c} 9.7 \\ 9.9 \end{array}$	$\frac{9.7}{9.9}$	$9.7 \\ 9.9$	66	83.76	
6.6	11, 1:00 "	4.4		18" below surface	9.9	9.8	9.85	8°c		
66	11, 1:00 "		12, 5:15	2' from bottom	9.3	Lost	10.4	66	78.35	
66	13, 1:00 " 13, 1:00 "		14, 4:40 " 14, 5:10 "	18" below surface	$10.4 \\ 10.5$	$\begin{array}{c} 10.4 \\ 10.5 \end{array}$	10.4	66	87.61	
44	13, 1:00 "		16, 8:20 a.m.		10.3	10.3	10.4		87.61	
6.6	13, 1:00 "	6.6	16, 8:50 "	2' '' ''	9.8	9.8	9.8	4.6	82.56	
66	16, 1:00 "			18" below surface		10.5	10.5		94.93	
6.6	16, 1:00 " 16, 1:00 "		18, 9:00 "	18" " " " " 2' from bottom	$\frac{9.8}{10.}$	$\frac{9.7}{10}$.	$\begin{vmatrix} 9.75 \\ 10. \end{vmatrix}$		88.15 90.41	
	16, 1:00 ,	6.6	18, 9:00 "	2'	10.3	10.3	10.3	4.4	93.13	
66	20, 1:00 "		21, 10:00 "	18" below surface	9.6	9.7	9.65	4.6	87.25	
	20, 1:00 "		21, 10:00 " 21, 11:15 "	18" " " " " " " " " " " " " " " " " " "	$\begin{array}{c} 9.7 \\ 8.9 \end{array}$	$\frac{9.7}{8.9}$	$\begin{array}{ c c } 9.7 \\ 8.9 \end{array}$	6.6	87.70	
	23, 1:00 "	6.6	24, 5:00 p.m.	18" below surface	8.6	8.6	8.6	14°c		
6 6	23, 1:00 "		25, 8:00 a.m.	18" " "	7.7	7.7	7.7	4.4	74.32	
6.6	23, 1:00 "		25, 9:10 "	2' from bottom	8.6	8.6	8.6	6.6	83.01	
ay	23, 1:00 ··· 2, 1:00 p.m.		25, 9:10 " 3, 10:30 a.m.	2' " " " " 18" below surface	$\begin{array}{c} 8.3 \\ 8.5 \end{array}$	$\frac{8.4}{8.5}$	8.35	19°c	82.53 90.71	
	2, 1:00 "	66	3, 10:30 "	18" " "	8.2	8.3	8.25	4.5	88.04	
	2, 1:00 "	66	3, 11:25 "	2' from bottom	8.9	8.9	8.9		94.98	
	2, 1:00 " 9. 1:00 "	ì	4. 10:00 "	2' '' 18" below surface	$\frac{8.1}{7.3}$	$\frac{8.2}{7.3}$	8.15	18°e	86 98 76.44	
	9, 1:00 "	6.6	11, 8:10 a.m.	18" " " "	7.2	7.2	7.2	10.0	75.39	
6.6	9, 1:00 "	1 66	10 5:10 n m	2' from bottom	7.9	7.9	7.9		82.72	
	9, 1:00 "	66	11, 8:35 a.m.	2' '' bolow surface	7.5	7.5	7.5	23°e.	78.53	
	16, 1:00 ··· 16, 1:00 ···		18, p.m. 18, p.m.	18" bclow surface 2' from bottom	$\frac{6.}{6.4}$	6.3	$\begin{bmatrix} 6.\\ 6.35 \end{bmatrix}$	23°C	69.20 73.24	
	30, 1:00 "	1.6	31. 2:30 p.m.	18" below surface	7.2	7.1	7.15	24°c	84.02	
	30, 1:00 "		31, 2:30 "	18" " "	7.1	7.1	7.1	66	83.43	
ine	30, 1:00 " = 6 1:00 p m	June	7 5:00 n m	2' from bottom 18" below surface	$\begin{bmatrix} 7. & \\ 5.8 & \end{bmatrix}$	7. 5.8	7. 5.8		82.25 6J.46	
ine	6, 1:00 p.m. 6, 1:00 "	6.6	7, 5:00	18" " "	6.	6.	6.	~i) C	71.86	
	6, 1:00	6.6	8, 11:00 a.m.	2' from bottom	6.4	6.4	6.1	41	76.65	
6.6	13, 1:00 "	6.	14, 4:40 p.m.	18" below surface	6.6	6.7	6.65	26° c		
	13. 1:00 "	• •	15, 9:00 a.m.	2' from bottom 18" below surface	$\begin{array}{c c} 6.4 \\ 6.5 \end{array}$	$\frac{6.1}{6.3}$	$\begin{array}{c} 6.25 \\ 6.4 \end{array}$	27°c	76.31 79.70	
	27, 1:00 ··· 27, 1:00 ···	6.6	28. 12:40	18" " " "	6.2	6.2	6.2	21 6	77.21	
	27, 1:00 "	6.6	98 1.00 "	9' from hottom	5.8	5.7	5.75	• •	71.61	
	27. 1:00 "	6.6	28. 1:00 "	2'	$\frac{5.9}{7.4}$	6.	5.95 7.35	200	74.09	
ıly	7, 5:45 p.m.	T 1								

DISSOLVED OXYGEN IN THE WATER OF THE

Mississippi River, Alton.

-	- Dat	e of			Dat	e of			Depth	ı of	Mg.	per Li	ter.	rer.	rent ira- n.	Remarks.
C	olle	etion.		1 k t	erm	inati	on.		Collec	tion.	1	0	Aver.	Ter Wa	Per Satu	Remarks.
	18	899			18	QL)								0		
June	25,	9:30 a	.m.	June	25.	9.30	a.m.	18"	below	surface		7.6	7.6	26° c		Missonri shore.
	25,	9:45	8	• •	25,	9:45		18"			7.3	7.2	7.25		88.5	Midstream.
6.6	25,	10:00		**	25.	10:00	6.6	18"	6.0	6.6	7.0	6.8	6.9	6.6		Missouri shore.
Aug.	26,	7:55 a	.m.	Aug.	26,	7:55	a.m.	18"	6.4	**	6.9	6.9	6.9	27°C	85.92	100' off Ill. shore.
"	26,	8:30	6.	6.1	26,	8:30	4.6	18"	4.4	4.6	7.3	7.4	7.35	28°c	93.03	y 4 distance from III, snore.
4.6	26,	9:10	6.0	6.4	26,	9:10	6.6	18"	6.6	4.6	7.8	7.9	7.85	6.6	99.36	Midstream.
44	26,	9:40	6.6		26,	9:40	6.	18"	6.6	4.6	7.7	7.8	7.75	٠.	98.10	14 distance 1 from Mo.shore.
4.4	26,	10:10	* *	* *		10:10		18"		6.6	7.8	7.7	5.75	6.6	98.10	100' fr. Mo. shore.
6.6	26,	1:151	n.m.	b b	26,	1:15	p.m.	18"	4.4	* *	7.5	7.5	7.5	23°c		100' fr. Ill. shore.
4.6	26,	1:45	••			1:45	4.1	18"	6.6	44	7.7	7.8	7.75		100.12	y 4 distance from Ill, shore.
4.6	26,	2:15	6.6	6.6	26.	2:15	4.6	18"	6.0	4.4	8.0	8.1	8.05		104.00	Midstream.
44		2:35				2:35	6.6	18"		s 6	8.3	8.2	8.25	29.5°c	107.56	1 ¹ ₃ distance 1 from Mo.shore.
6.6	26,	3:05	4.6	6.6	26,	3:05	4.4	18"	* * *	6.	8.3	8.3	8.3	6.6	108.21	100' fr. Mo. shore.
4.6		4:10	* *	**	26,	4:10	6.6	15"	• •	6.4	7.9	7.8	7 85	6.1	102.34	100' fr. Ill. snore.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

1899	
EAR	
-	
FOR	
RESULTS	
RES	
OF	
GES	
ERAGES	

University of Illinois.

Report of ARTHUR W. PALMER,

-01 -01	Nitrates	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
NITRO- GEN A	Nitrites.	0.000 0.000
	papuad sus-	25.11.25.25.25.25.25.25.25.25.25.25.25.25.25.
ORGANIC NITROGEN	Dis- solved.	\$\frac{1}{2}\frac{1}\frac{1}{2}\f
S S S S S S S S S S S S S S S S S S S	T'otal.	200.000
IA.	p,pud	100 100
NITROGEN AS AMMONIA	pio h'vios	1
N AS A	ToTT Pictor Picto	1.1 1.2
FROGE	Sinom Sinom	
	Freehm-	F-0
OXYGEN CONSUMED.	Bysuspd.	\(\frac{1}{2} \cdot \
ONYGEN	By Dis-	<u></u>
5	Total.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
.90	Съдотіт	400 00 00 00 00 00 00 00 00 00 00 00 00
	Sus- Sus-	1880 2188 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
EVAPORATION.	Dis- solv'd solv'd solv'd solv'd	84282488488844888448888888888888888888
VAPOI	Tot'l Loss	\$\$\$45\$?!! \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
ON	pəpuəd -sns	855744484646688846884658744888748888888888
SIDUE	Dis- solved.	28 28 28 28 28 28 28 28 28 28 28 28 28 2
RESU	Total.	28
10.0		
	AVERAG	85885188888888888888888888888888888888
l do	RING 99	
PERIOD	COVERING 1899	
1		
	SOURCE OF WATER.	Illinois and Michigan Canal—Bridgeport Desplaines River—Lockport Desplaines River—Joliet Desplaines River—Joliet Kankakee River—Joliet Kankakee River—Morris Illinois River—Ottawa Illinois River—La Salle Illinois River—La Salle Illinois River—La Salle Illinois River—La Salle Illinois River—Hary Illinois River—Herry Illinois River—Herry Illinois River—Herry Illinois River—Herry Illinois River—Herry Illinois River—Havana Sangamon River—Chandlerville Illinois River—Grafton Illinois River—Grafton Mississipi River—Grafton Miss. River—Alton, Midstream. Miss. River—Alton, Midstream. Miss. River—Alton, Mo. Shore Miss. River—Alton, Mo. Shore Miss. River—Alton, Mo. Shore Miss. River—Alton, Mo. Shore Miss. River—Mitchell, Midstream. Miss. River—Mitchell, Missouri Shore Miss. River—Mitchell, Missouri Shore Miss. River—Jeff. Bar, East Shore Miss. River—Jeff. Bar, E. of Center Miss. River—Jeff. Bar, West Shore Miss. River—Jeff. Bar, West Shore
) E	O ON STATIC	

TABLE B.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

AVERAGES OF RESULTS FOR YEAR 1900.

	-
	0
	Z
	-
	bo
- 1	_
	_
	-
	۰-
	~
	C
	P .
	'n
	•
	press
	4
	U
	ė.
	~
	М.
	1
۰	-
	_
	5
- 1	-
- i	
	-

Report of ARTHUR W. PALMER,

se	Sitrates	8 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
NITRO- GEN A	Nitrites.	000 000 000 000 000 000 000 000 000 00
	beuged	86.50 86
ORGANIC NITROGEN.	solved.	\$2.50 \$3.50 \$3.50 \$3.50 \$4.50 \$5.50
ORG. NITR	-sid	4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
	Total,	888.888.888.888.888.888.888.888.888.88
AMMONIA.	P.pud.	=
AS AM	T'ot'T pis- polv'd solv'd	25
NITROGEN	T'10'T	### ### ### ### ######################
NITR	Ргеели». топів.	10101 10
EN MED.	BySuspd Matter.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN	By Dis-	$\mu \cdot \omega + \alpha - \omega + \alpha \circ \omega \cdot \omega \cdot \omega \circ \omega \omega \cdot \sigma \circ \omega \circ \omega \circ \omega \circ \omega \cdot \omega \cdot \sigma \circ \circ$
	laio'f	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
16,	Chlorin	<u> </u>
	gas- gas-	
RATIO	Dis- Bolv'd Sas- Bea'd Dis- Deu'd	8-8-9-4-4-4-4-4-4-4-4-4-4-4-4-4-4-4-4-4-
EVAPORATION.	l'10T	888498882488999999999999999999999999999
0.N	pepued sus-	884 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
RESIDUE	Dis- beyied,	18 28 28 28 28 28 28 28 28 28 28 28 28 28
24	Total.	888 888 888 888 888 888 888 888 888 88
E OE	улевуе	8895151515895555558888855985858555555555
	5	88888844808 6 4 4 4 4 4 8 8 8 8 8 8 8 8 8 8 8 8 8
PERIOD	COVERING 1900	12-0-61-7-4-4-7-3-8-8-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-1-8-9-1-
	Co	e
'NG	SOURCE OF WATER.	Illinois and Michigan Canal—Bridgeport sanitary Canal Kedzie Avenue. Illinois and Michigan Canal—Lockport. Desplaines River—Lock port Desplaines River—Moliet Canal Fox Minhigton. Illinois River—Morris Canal Las Alle. Illinois River—Ottawa. Illinois River—Ottawa. Illinois River—La Salle. Illinois River—Henry. Illinois River—Henry. Illinois River—Henry. Illinois River—Henry. Illinois River—Pekin. Illinois River—Chandlerville. Illinois River—Chandlerville. Illinois River—Chandlerville. Illinois River—Chandlerville. Illinois River—Chandlerville. Illinois River—Readstown. Illinois River—Manpsville. Illinois River—Mon, Mids. From Mo. Shore Miss. River—Alton, Mids. From Mo. Shore. Miss. River—Alton, Midstream. Miss. River—Mitchell, Missourf Shore. Miss. River—Jeff. Bar. Center. Miss. River—Jeff. Bar. West Shore. Miss. River—Jeff. Bar., West Shore.
el ((百円) (日本)

TABLE C.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

AVERAGES OF RESULTS-For Period June, July, August and September, 1899.

ALMER,
PAL
¥.
UR
RTHU
AB
0 3
eport
2

University of Illinois.

.N(R	RESIDUE ON	Ev	APORATION.	TON.		,9£	၁၀	CONSUMED	.α	4	AMMONIA.	SN AS NIA.		SN	ORGANIC NITROGEN.	÷	NITRO- GEN AS
)] L	Source of Water		-p	eg	Loss o	NO IGNITION	TION.	iiro	J.		pd pd		Albu	Albuminoid	Am	l.			sa
ATS	SOUNCE OF WALEN.	rioT	Si([Solve	suS	['10T]	Dis-	p,pud -sns	Срјо	rioT	By Dis	BySus Matte	FreeA	[,10,],	p.vlos	p,pud ·sus	[E10T]	Dis-	-sng	Nitrite
-6	Illinois and Michigan Canal—Bridgeport	590.2	505.2	35.	60 8	51.6	15.1	119.	39.9	20.6	19.3	17.22	25.38	1.01	1.37	67.75		1 77 7	820
	Desplaines River—Lockport	335.6	316.6		5.4	84		6.4	14.1	12.9	ં	.07	.52	44.	38.	1.13	00.00		010234
	Desplaines River—Joliet (above town)	493.5	473.3	28 75 23 75	4.2.4 5.5	34.5		3.7	5. E.	17.6		ي اي اي	1.76	7.00	1.02	3.81		2.23	0124
	Illinois River—Morris	419.3	396.9	33.4		40.9		68.5	16.4	13.8	9:0	9.3		jro	.35	.8.		; ₹	0242
_	Fox River—Ottawa	380.9	298.1	85.5 85.5 85.5 85.5 85.5 85.5 85.5 85.5		49.3		6.4	13.1	10.1	eo +	.053	.53	.31	종:	1.28		.615	013
_	Illinois River—Orlawa Rig Vermillion River—La Salle	746.	708.3	38.1	0 x			23.62	7.00	3 7	1.6	718.9	0.6	17	1.61	1.13		98.	691
_	Illinois River-La Salle.	417.9	372.2	45.7		43.4		50.8	14.1	11.2	2.9	1.68	57.	98.	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1.31		j re	5338
2	Illinois and Michigan Canal-La Salle		307.1	39.3		43.9		14.9	11.9	9.5	2.7	.578	ۍ. د	88.	.17	1.231		.371	0443
	Illinois River—Henry.		332 S	4.47		10.1		433.20	14.6	11.1	က က်	1.02	.64	œ;	98:	1.57		.65	4515
	Illinois River—Averyville	253.5 253.0	240.2	19.0		42.1		30.8 - 80.8	3.0	10.4	200	36	٠. و	સ્કું ક જ	.15	 83:8		ج ج ا	906
	Illinois River—Pekin.		313.3	41.3	8.77	45.1		36.7	13.9	11.1	. €0 . ∞	707	3 3	000	2 %	5.5		٠ و تو	9266
	Illinois River-Havana		330.	46.7	48.9	£3.8		29.9	12.9	10.1	2.8	17:	3.	.34	.16	1.31		44.	164
000	Sangamon River - Chandlerville		273.2	106.3	50.4	36.3			₹- ¢	4.9	ლ ი დ. ი	.033	88:	11.	.14	.851		.463	0287
	Illinois River—Beardstown		291.1	108.7		34.3		: ;;	20.00		0. c	-	7.	98.	<u>∞</u> ;	1.086		.476	1257
_	Illinois River—Randpsvine	v C	0000	73.×	? o	31.6		20.7	201	0 00 0 00	3	1008	65.5	3.5	21.	1.004		65.	# 10.00 # 1.00 #
0.00	Mississiphi River - Graffon	-	156.2	170.9		30.00		2.6	16.57	1.0	- 6	0406	46	0.00	26.	908		600	1000
	Mississippi River—Alton, Illinois Shore	343.3	198.4	144.9				7.00	13.	8.6	ec.	.031	2.4.	6	3 00	1.066		00.00	0000
	Mississippi River-Alton, 14 dis. from Ill. Shore	319.5	174.6	134.9		28.5	×	4.34	15.1	10.7	4.4	.023	.43	.22.	.21	1.136		9	0103
٠ ان	Mississippi River-Alton, Midstream	353.5		191.4	39.7	31.	 00	2.6	15.7	10.8	4.9	.032	.45	ં	.25	1.211		.736	0058
	Mississippi River-Alton, 14 dis. from Mo. Shore	3.00 3.00 3.00 3.00 3.00 3.00 3.00 3.00	159.8	213.5		30.7	3	6.5 6.5	15.7	10.8	4.9	860.	.47	.21	98.	1.287		785	0045
_	Mississippi River—Alton, Missouri Shore	371.2	162.6	208.6		31.4	14 9	65 70.	15.6	10.8	8.7	.033	.45	રો સ		1.223		.726	0041
	Missouri River—West Allon	2207.6		2041.9		20.1	2.5	χ. 1-1	21.1	0.5	15.4	120	£ 1	= :	39.	2.177		1.884	.0071
	Mississippi Kiver Mitchell, Illinois Snore	756.0		1239.9		£ 6	54.5	5.1	25.25	x : 0x	12.7	.0. 10.	.77	61.		2.162		1.727	.0087
02.5	Mississippi Kiver—Mitchell, Midstream	1930.5	216.8 215.8	16.17.0	0 c	20.3	45.55 C. C. C.	. c	233	5. C	15.1	220.	*	91:	æ 8	2.38		1.982	.0071
-	Mississippi Miver—Mitchell, Missouri Shore	9338	200 ×	9118		98.	53.0	2.5	94.0	- 1-	17.4	620.	. o	<u> </u>	0.8	2.00 9.00 9.00		200	.000±
_	THE POST OF THE PARTY OF THE PA					-													

TABLE D.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

AVERAGES OF RESULTS-For Period June, July, August and September, 1900,

Report of ARTHUR W. PALMER,
University of Illinois.

Source of Water. Although Canal Principles and Mobilean Canal	A 36	solariiN	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
Constitute Con	NITRO	Solitiv	
Source of Water. Fig. 12		рэриэс	
Constitute Con	NIC GEN.	-sus	0
Constitute Con	ORGA		
Courte of Water Courte of		Total.	01-01
Courties Courties		Sus.	
Courties Courties		Dis- Dis-	5. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
Source of Wayer Constitution C	AMMO	T'10T	28 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 - 4 -
Source op Water. Source op W	7.		
Source of Waver. Source of W		Matter	$\frac{1}{1} \cos \alpha \widehat{\alpha} + \alpha \widehat{\alpha} $
A	CGEN UMED.	pealos	
1	Oxy		
18 19 19 19 19 19 19 19			
Source of Water. Source of W	.et	1.1	
Course of Water. Course of W		Softial Softia Softial Softial Softial Softial Softial Softial Softial Softial	
Source of Warer. Source of W	ATION.		នុ ១ ភ ភ ជន់ និង ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ ភ
Thinois and Michigan Canal—Bridgeport. 174 20 20 20 20 20 20 20 2		Lorr Log	
Source of Water. 10	s on Ea	paptiad -snS	1
Source of Water. 10	RESIDUI	.bevies	41-68226822622222222222222222222222222222
82888888888888888888888888888888888888	H	Total.	201.02 201.02 201.02 201.02 201.03 20
82888888888888888888888888888888888888		SOURCE OF WATER.	llinols and Michigan Canal—Bridgeport anitary Canal—Fedzie Avenue. Ilinols and Michigan Canal—Lockport. Seplaines River—Jollet. Ilinois River—Morris Sir Vermillion River—La Salle Sir Vermillion River—La Salle Sir Vermillion River—La Salle Sir Vermillion River—La Salle Ilinois River—Tenry Ilinois River—Tenry Ilinois River—Fekhn Ilinois River—Fekhn Ilinois River—Fekhn Ilinois River—Fekhn Ilinois River—Fekhn Ilinois River—Fekhn Ilinois River—Grafton Ilinois River—Grafton Ilinois River—Grafton Ilinois River—Alton, Illinois Shore Ississippi River—Alton, Adis, from Ill. Shore Ississippi River—Alton, Missouri Shore Ississippi River—Alton, Missouri Shore Ississippi River—Alton, Missouri Shore Ississippi River—Alton, Missouri Shore Ississippi River—Mitchell, Illinois Shore Ississippi River—Mitchell, Missouri Shore
11.			42000-00-00-00-00-00-00-00-00-00-00-00-00

TABLE E.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO, SANITARY WATER ANALYSIS-PARTS PER MILLION

COMPARATIVE AVERAGES, 1899 AND 1900.

University of Chieago.

Report of EDWIN O. JORDAN,

631.000 650.000 28,200 12,500 492,000 744,286 5,000 16.100 400,000 445.000 6,500 20.600 27.470 116.000 7.970 19.200 8,8.0 13,290 64.203 54,600 51,800 36,800 492,300 68,400 17,470 23,100 5.080 94,000 ,755,000 Colonies per Cubic Centi-No. of meter. NITROGEN AS trates ż trites. .0136 $\frac{\cdot}{N}$ 031. 1336 p,pud -sus 191 19 <u>25</u> Albuminoid Am. NITROGEN AS -si(I 149 301 301 AMMONIA
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000
 200000 Total 7.08 1.60 1.60 1.30 1.31 1.31 1.31 1.32 1.33 1.34 1.35 .262 999 999 999 996 996 27.72 1.046 2.008 пошв FreeAm-p.puad OXYGEN CONSUMED solved -siġ 80. 30.3 16.6 32.1 18.1 10.1 80.68 4.60 4.4.62 4.63 4.4.63 Total. 96.6 96.6 124.5 124.5 6.1 6.1 41.5 CHLORINE, pended Sus-RESIDUE ON EVAPORATION. 268. 394.2 242.8 297.7 264. solved. 481.8 389. 485.1 451.8 353.4 333. 480.1 274. 292. 957.3 267.2 360. 326.9 249. 353. 368.1 250.9 370.1 313. 336.1 Dis-355.4 306. 375.6 562.7 421.8 562. 491. 371. 307.4 343.8 413.1 326.7 395.8 320. 996.6 441. 417.6 307.7 399.1 351. 387.6 303. 394.9 348. 388.6 303. 335.1 4409. 385 6 369.3 500.5 500.5 Total. 390.4 351. 313. 1900 1800 1500 1900 1900 1000 Jan.-June, 1900 May-Dec., 1899 1900 1900 6681 899 6681 668 190) 1899 6681 0061 6681 1900 COVERING May-Dec., 1 Jan.-June, 1 May-Dec., 1 Jan.-June, 1 Jan.-June, May-Dec., Jan.-June, 1 May-Dec.; 1 Jan.-June, 1 June-Dec., Jan.-June, Period Jan.-June, Jan.-June, May-Dec., May-Dec., Jan.-June, Jan.-June, May-Dec., Jan.-June, May-Dec., Jan.-June, Jan.-June, May-Dec., Jan.-June, May-Dec., June-Dec., May-Dec., May-Dec. May-Dec., May-Dec., May-Dec., Illinois and Michigan Canal—Bridgeport..... Sangamon River—Chandlerville Desplaines River—Loekport..... Illinois and Miehigan Canal—La Salie. Desplaines River—Joliet (above town) Kankakee River-Wilmington.... Illinois River—Ottawa (above town)..... Big Vermilion River-La Salle..... Illinois River—Averyville..... Illinois and Miehigan Canal—Lockport...... Illinois River—Henry..... Hinois River-Pekin..... Illinois River-Wesley City Fox River-Ottawa.... Minois River-Mavana..... Illinois River-Morris Illinois River-Kampsville..... Illinois River-Grafton.... Illinois River-Beardstown Illinois River-La Salle..... SOURCE OF WATER.

TABLE E-Continued.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

COMPARATIVE AVERAGES, 1899 AND 1900.

Report of Edwin O. Jordan, University of Chicago.

SOURCE OF WATER.	;	N SI	EVAPORATION	ON.	'ME'	Co	CONSUMED		ANNONI	AMMONIA	A	EI V.	NITROGEN	ن د	olonies
	PERIOD COVERING	Total.	Dis- solved.	Sus-	Сигокі	Total	Dis.	Sus- Sus- Freekin-	вінош	Albuminoid sid sid sid sid sid sid sid sid sid s	Sus -sus	pnd'd NI.	NI- trates.	1	per Cubic Centi- meter.
Mississippi Biver Grafton.		407.3	165 8	241.5	© 1 1 00 0	12.7	8.6	1.1	0) (=	7.600
Mississippi River-Alton, East Bank	JanJune, 1800 May-Dec., 1890	35.77		247.	-1 2 6 6	10.4	- 1- - 00:	3.45	076					~ ±C	30,600
	_	587.	C.5	371.	٠٠. ن	6.01	6.1	8.8	01			-		- 66	30,300
Mississippi River-Alton, East of Center	May-Dec., 1899	326.2	183.9	143.3	rυ π ≟ α	1 1 1	بر من در	4. 4					_	SS 15	7,720
Mississ ppi River-Alton, Midstream		379.2		214.3) <u>}-</u>	11.5	. *	3	051	165	298 . 237		800	3 88	6.650
		80c		346.	0.4 0.5	23.2	. 1	6.6		_	_			120	51,800
Mississippi Kiver—Alton, West of Center	May-1.ec., 1899	2 2 3 3 4 4 4		348	7. 9	9. 8. 9. 8.	2.5	ص ص ص ح ح		_		_		 R Ž	6,350
Mississippi River-Alton, West Ban commen	-	391.2		227.4	2 74	0.11	1.7	· · ·						92	7,760
-	-	489.		. 1	4. 60.	13.	0.0	6.		_	_			379	42.500
	July-Dec., 1899	1121.5	204.7		<u>10 1</u>	∞ <u>c</u>	0) t	∞ - ∞ -	-					17.3	8,170
Wississippi Biver-Chain of Bocks, East Bank		1503.1				1 to	- oc	1.5						و و و	43,100
	_	917.			6.7	13.7	6.2	→						2 00	36.200
M. ssissippi River Chain of Rocks, Midstream	_	1619.2		1387.4		15.6	20 I	8.0		_	_	_		34	12,390
Westernian Picer Chain of Poeks Intake Tower	Anr - Dec 1800	1045 0		1600 7	ο α ν ι·	ν. τ. ο	5 15 5 15	 ;		_				٠ •	46.600
		1626.		1399	10.3	(-) · ·	3.4.0	. co						 2 99	000.14
	- ,	1993.9		1752.4	φ. φ.	15.3	4.6	7.01						12	1.966
	Tun June, 1900	1 2055 1 949 9		17.81	× 0		- ·							9,3	19.700
	-	881.6	2 Kg	101	13.0	0.0	0 7	3 9							255.25
Missi sippi River Jefferson Barracks, East Bank		1328.1		1025 9	5.6	15.	6.5	20.						33.	17.590
		686.		495.	1-1 63 0	9:2:	6.3	xc :		_				7.6	65.600
Mississippa River—Jenefson barracks, Last of Center	Jan Apr. 1900	1001		1.00.c	:0 O	 	ه وه چ نه			_				27	13,120
Mississippi River-Jefferson Barracks, Midstream	_	1502.2		1286.8	6.1	3. c	5.9							34	13 390
	_	× 25.7		641.		13.8	∞.℃			_	_			200	69.500
Mississippi River-Jefferson Barracks, West of Center		1633.		8.0681	 	6: 3	6.1,	.0		_	1			10	21,850
Missleinni Blyer-Jefferson Barracks, West Bank	May-Dec. 1899	1750.5		1050	= x	x x	 		•	513	0				74.800
	-	1087.		776.	12.6	2 22		. j.		80	08			200	10. 100 10. 000
-	AprJune, 1900	215.8	88.	85	0.87	∞ (B)	5.01	2.9	33	ã: 66	63	_		289	1.332,000
Sanitary (anal. Dum at Lockboth	AprJune, 1800	E024		25.00	1.2.0	5.27	26.7	 	25. 25. 26.		: ::	10.		118	1,167.00
Josephannes Elvel, John Lown, John John Holler below Rock Island Bridge or Jefferson Street		250.	22.1	ē - 86 2 86	22.00	G ∝	5 1 1 C	1.0	25 C. S.	0.1.0	55 - 1.4	10. 78.		990	766.700

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-MONTHLY AVERAGES OF COLONY COUNTS.

TABLE F.

Report of T. J. Burnill, University of Illinois.

December.	2,151,666	16,300	1.026,250	25,300	170,000	261,766	101,400	223,000	12,757	9,300	20,666	301,333	10,266	114,100
Хочетьет.	2,497,000	9,640	1,023,000	11,650	747,000	18,070	11,910	164,400	14,050	1,640	705,000	159,040	7,100	26,070
October.	2,150,000	25,783	1,145,000	11,075	505,000	33,050	27,112	131,250	3,987	3,768	2,373,750	12,433	3,683	5,000
September.	1,186,665 2,483,333 315,000 696,666	24,160	651,666 217,000 203,330	985,000 5,387 4,950	2,186,100	34,875 4,825	9,375	166,375	3,102 3,200 11,100	136,700 3,100 4,466	484,575 264,375	1,061,875	20,500 20,300 316	3,039 5,550
nsugu v	1,233,700 1,215,000 3,215,000	38,850	1,205,666 3,220,000 134,750	4,611,666	4,882,375	7,512 18,830 4,560	2,963 11,875	241,000	1,480 7,000 26,350	2,4850 8,060 8,060	2,355,000 99,250 1,98,33	286,500 5,725	19,783	10,287 15,090
July.	863,888 3,895,555 5,323,750	11,162	601,250 2,848,666 170,000	3,201,666 4,720 950	930,750	3,950 9,950 9,670	3,266 6,370	38,375	4,510 5,475 22,362	40,433 9,358 4,233	3,925,000 96,000	313,375	9,900 13,900	1,352
June.	377,000 1,035,500 3,885,750	14,912	1,445,000	1,369.375 22,837 3,650	999,875	20.550 14.237 11.027	2.412	170,050 31,062	7.062 8,550 64,350	3,487 8,700 1,637	1,476,480 58,625 1,500,000	150,625	7,125 29,562	4,400
May.	1,474,000	5,637	485,000	3.300	46,200	15,125	5,800	25,300	22,270	14,260	70,400	106,500	23,550	12,880
Li _{T(} f A	6,037,500	21,375	1.655,000	52,000	392,500	68,500	49,250	157,750	33,000	39,875	22,750	27,250	20.900	28,375
March,	4,240,000	167,500	2,000,000	75 250	1,593,750	116,750	84,166	212,000	103,125	183,625	107,750	125,125	105.875	145,750
February.	6,063,333		1.448.333	000 601	1,731,250	228,000	0.00	169,625		160,500	95,666	84,000	79,575	
January.	3,862,000	42,370	973.000	067	2,737,500	256,250	12,275	193,400	13,750	41,462	89,000	128,100	10.620	239,750
SOURCE OF WATER.	Sanitary Canal	laines River	Sanitary Canal. Desplaines River.	Kankakee River	ois R	Fox River		Illinois and Michigan Canal.	Big Vermilion River.	3 3 3	3 3 3	3 3 3	Sangamon River	Illinois River.
Location.	Kedzie Ave Bridgeport Lockport	3 3	Joliet.	Wilmington	Morris	Ottawa	La Salle	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Henry	Averyville	Wesley City	Havana	Chandlerville	Beardstown
YEAR.	1900 1899 1899	888 888 888 888 888 888 888 888 888 88	1899	1899	1900	1999	068	2888 2888 2888	1899 1900 1899	000000000000000000000000000000000000000	1899	1899	985 985 985	1800

Report of T. J. BURRILL, University of Illinois. STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-MONTHLY AVERAGES OF COLONY COUNTS. TABLE F-Continued

15, 183 31.650 15,912 18,330 27 9.816 9.911 19.087 посеппрет. 7.160 6.33.0 17,030 5,7:0 11,600 25.075 26.612 28,510 × 955 ZOVERBEE 1,717 2,715 2,112 5. 455 2,375 9,200 3.075 11,775 11.462 10,187 16.325 33.625 11,833 45,250 52,875 1.710 59,500 October, 12,125 1.990 10,125 60, 125 18.787 September, 45,712 63,125 1,950 95 0.200 1.806 1. 6,110 8,730 8,735 13,300 32,275 10,256 32,695 29.837 54,525 usuguy 12,050 3,161 1,850 2,628 3,911 915 5.902 29.275 20,000 13,400 5,737 851 1,650 96 27,045 Ame 55,350 13.25 14.15 14.15 15.25 June. 16,040 21,800 34,200 18,180 6,300 17,620 26.000 24.080 31.300 64,125 66,200 May. 48,500 25.500 48,333 66.000 85,500 60,125 58,375 57,500 54,666 15,825 14,550 46,625 17,500 77,000 55,333 18.125 750 April. 97,000 145,375 98 166 125,833 130.500 154,750 159,500 119,750 175,000 166,250 149,875 144,875 143,333 179,750 130,875 129,750 155,625 227,750 10,137 March, 86,500 202,666 191,500 88,500 66,800 79,750 69,625 98, 125 105, 125 104,100 113,100 95, 150 50,300 97,500 136,500 79,666 61.666 40,066 Rebruary, 64,925 46,600 13,750 18,940 84,960 23,500 23,350 19,040 18,620 24,460 35,087 70,075 76,525 4.865 January. M dlst. from No. Shore. 400 yds. from Mo. Shore Mississippl River.... Miss. River, 100 ft. from Ill. Shore... West of Center.... 400 yds. from Ill. Shore. West Shore.... 14 dist. from Ill. Shore. East of Center ... Center..... Center Inlet Tower.... Illinois River.... Center of Channel..... 100 ft. from Mo. Shore. Mississippi River, East Shore.... Missouri River..... SOURCE OF WATER. Jeff. Barracks Alton St. Louis Tap. West Alton ... Chlcago Tap .. Grafton Mitchell.... Kampsville. 282333 0.68.09 VIII 9888 888 899 LEVE

TABLE G.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS—PARTS PER MILLION. For Period June, July, August and September, 1899-1900.

	Illinois.
ARTHUR W. PALMER,	University of
eport of	

NITROGEN AS	Nitrates	. 881 1.089 July 12 June 14 June 25 .339	July 19 June 25 Aug. 2 June 13 June 13		June 27 July 3 Ang. 7, 28	(Sept. 4, 25 1.127 1.127 1.239 1.339 1.736 1.736 1.881 1.089
NIT	Nitrites	26.50 1.65 1.65 1.65 1.65 1.65 1.65 1.65 1.65	.02 .013 .001	.003	.001	0132 0132 0132 0064 029 0132 0132 0132 0132 0132 0132 0132 0132
S.N.	pepued ens-	.331 .31 .683 .552		2.144		
ORGANIC NITROGEN.	Dis- solved.	.615 .399 .399 .525 .389		304	:	.676 .485 .665 .443 .464 .3087 .3884 .3884 .3884 .3884 .389
5Z	Total.	.936 .609 .609 .1.208		755 2.448		.733 .662 .749 .749 .749 .7397 .609
NIA.	Sus- b'buq	410 80 8 90 90 90 90 90 90 90 90 90 90 90 90 90 9			:	
AS AMMONIA.	Albuminoid Dis- Dis-	.233 .185 June 14 Sept. 5 June 28 July 6 .209 .191	June 7 July 30 Aug. 2 Sept. 6 June 20	.1232 Aug 17 June 27 Sept. 21	July 10	143 143 143 169 169 185 185
NITROGEN AS	T'10'T	.339 274 208 208 457 7447	. 88 . 544 . 32 . 288	. 352	.324	2446 2733 2838 2932 2777 2774 2339
NITH	FreeAm- monia.	.079 .396 .16 .01 .038 .0882	.044	.0979 .028 .066	.016	.0601 .0502 .105 .105 .148 .098 .098 .0815 .079
SUMED.	Bysuspd	1.76 2.1 Sept. 27 Sept. 26 Aug. 9 July 6-20 4.18 6.4	June 14 Sept. 26 June 7 Aug. 21 June 6	15.4 16.16 Aug. 31 July 10 Sept. 28	June 5	6.1.9.1.9.1.9.9.9.9.1.9.1.9.1.9.1.9.1.9.
OXYGEN CONSUMED.	By Dis- solved	8.29 5.8 June 14 July 27 Sept. 20 Aug. 13 11.33		21.06 5.66 21.38 5.22 31.9 Aug. 2 28.7 June 27 11.55 Sept. 28	Aug. 14	07-01-7-7-4-4-8-7-07-07-07-7-7-1-1-2-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8
OX	Total.	10.05 7.9 11.9 8.4 6.6 6.6 16.51		21.06 21.38 31.9 28.7	13.4	13.5 13.07 8.68 7.81 6.18 8.74 9.74 10.05
.9n	Chlori	14.8 322. 22. 22. 12. 3.17 3.17	4 8	7.94 14.99 14.8 34.6 5.1	6.4	3.687 3.24 3.24 4.86 5.77 5.77 5.09 14.83 14.83 14.94
			Aug. 30 Sept. 6 July 9 June 7 Sept. 26	Sept. 28 Sept. 25 July 27	July 3	
	sng -sng	6.7		6 1. 6	:	7-10-00 1-00 0-4-11 0-1-00 1-1-1-1-1-1-1-1-1-1-1-1-1-1-1
ATION	Loss on Ignit'n b'vlos b'vlos b'vlos con l'10T	41.334.6 34.726.8 40.931.6 42.837.6		94.9 20.7	•	25. 41.5 34.05 46.92.38.3 33.2 82.85 58.8 49 15 23.08 46.8 39.6 38.1 74.8 65.6 45.9 50.7 44.3 106.8 41.3 28.2 73.8 41.3 28.2 73.8 41.3 28.2 85.4 34.7 26.8
APOR					:	25.08 48.034.700.444.00.808.808.74.00.114.
ON EV	-sng	73.8 85.4.2 146.4 146.4 28.3 28.3 171.	549.6 440.8 12.8 66.6	2 1984.6 9 3299.6 24 3917.2 28 644.4	738.4	
RESIDUE ON EVAPORATION	Dis-	268.05 239.8 July 19 Sept. 5 June 28 Aug. 6 156.1 159.9	June 7 July: 30 July 12 Sept. 12	223.7 2044.8 241.2 1984.6 Aug. 9 3299.6 July 24 3917.2 Sept. 28 644.4	Aug. 14	259.05 25
RE	T'otal.	331.9 320.2 320.2 J S 327.1 383.4				314.4 298.4 380.9 316.1 746.4 484. 379.5 341.9 320.2
	AVERAGE.	Average of 4 mos. Highest. Lowest Average of 4 mos.	: : : :~	of 3 mos. Aug. Average of 4 mos. Highest.	:	Average of 4 mos. """" """ """ """ """ """ """ """ """
	Year.	1899 1900 1900 1899 1900 1900	1899 1899 1899 1900	1899 1900 1899 1899	1900	1899 1899 1899 1899 1899 1899 1899 1899
200	NATER.	Illinois River —Grafton	Mississippi River —Grafton	Mississippi River W. Alton		Kankakee R. Wilmington. Fox River Ottawa Big Vermillion River, La Salle Sangamon R., Chandlerville Illinois River —Grafton

TABLE H.


STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. For Years 1899-1900.

JON. Report of ARTHUR W. PALMER,

University of Illinois.

A. ORGANIC NITRO- GEN AS	Total. Total. Dis-sign solved. Sus-shorted. Sus-shorted.	165 1.037 .609 .428 .651 1.07 198 .931 .478 .453 .024 1.362 1.886 Nov. 15 3.52 Feb. 14 .648 Nov. 1	251 1.142 .489 .653 .01 .295 2.45	373 1.378 .272 1.106 .006 .337 741 2.279 .325 1.954 .0072 .652 8.196 .4.96 .Mar. 7 4.56 .Dec. 28 .36 .Jan. 4	106 .886 .643 .245 .011 1.084 12 .892 .567 .325 .014 2.208 146 .91 .556 .384 .008 .373 148 .81 .008 .373 .416 085 .639 .406 .243 .022 .845 113 .596 .314 .281 .018 .522 108 .72 .38 .34 .043 .62 181 .783 .329 .454 .021 1.84 185 .037 .478 .453 .051 1.07
NITROGEN AS AMMONIA	Prochmannian. Interpretation of the process of the	221 .412 .347. .37 .403 .205. .88 .Nov. 29 2.24 .Jan. 4 .008 .June 7 .032 .June 7	8 9	. 175 . 875 . 875 . 134 . 13 . Nar. 13 . Sept. 28 . Apr. 16	3.4448 = = = = 2.28 3.4448 = = = = 2.28
OXYGEN CONSUMED.	Total. By Dissolved. By Suspd Matter.	7.8 2.1 6.6 4.3 May 31 Feb. 14 May 24 Aug. 13	89 10.8 6.3 1 8.8 6.3 9 May 3 6 April 9 9 Nov. 1	9 4.4 9 5 6 4.9 13.7 7 June 27 6 Nov. 30 9 Jan. 4	0.000
.9	Chlorin	Nov. 8 36.6 16.8 June 7 45 7.	June 28 15 15 15 15 15 15 15 15 15 15 15 15 15	Nov. 16 39. 31. Jan. 18 35.3 28. July 27 5.1 6. Ang. 1 7.1 3.	20 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
RESIDUE ON EVAPORATION.	Total Thorpoological forms Total T	349.5 274.7 74.8 38.7 29.7 9. 438.6 229.1 209.5 38.5 27.3 11.2 Peb. 14 1311.2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	342.4 153.1 189.3 36.1 26.7 9.4 365. May 24 568.8 April 23 1053.2 April 23 1053.2 July 12 12.8	Average of 23, 1284.6 285.2 869.4 50.3 17.6 32.7 1186.6 286.1 1600.5 67.5 19, 18.5 Highest, June 24 3917.2 Lowest, Dec. 28 180.4	300.77 905.5 30.4 37.8 29.7 5.6 331.4 259.5 71.9 38.4 31.8 6.6 341.8 29.7 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.6 31.8 6.7 7.4 4.0 7.4 4.0 8.3 7.9 4.6 31.8 8.1 2.8 3.1 2.8 3.1 2.8 3.1 2.8 3.1 3.1 3.1 3.1 3.1 3.1 3.1 3.1 3.1 3.1
	Average.	1899 Average of 34. 1900 39. 1900 lighest 1900 Lowest	1800 Average of 33 1900 Highest	899 Average of 23, 1284.6 1900 Highest. 1900 West.	856 Average of 20 1900
	SOURCE OF WATER.	Illinois River 190 190 190 190 190	Mississippi 198 River —Grafton 198	Missouri River 190 1	Kankakee R. 18. Wilmington 199 Fox River 198 Bigvermillion 188 River, La Salle 199 Sangamon R 188 Chandlerville 199 Illinois River 198





REPORT OF THE UNIVERSITY OF ILLINOIS.

BY

PROFESSOR T. J. BURRILL, Ph. D

THE EXTENT OF THE WORK.

The work herein reported covered the period from June 1, 1899, to October 15, 1900. During this time 2,800 samples of water were received from the collecting stations named below. The samples were usually taken once a week from each station, but from January 30 to April 30, 1900, daily collections were made from the Illinois River at Grafton, Ill., and three times a week from the Illinois and Mississippi rivers at Grafton, and from the latter near Mitchell, from July 9, to October 15, 1900. From each of these 2,800 samples at least duplicate cultural tests were made for the number of bacteria present, and frequently others were made for the purpose of comparison in different culture media or under conditions varying from the usual method, aggregating, all told, about 7,000 cultures for colony counts. Further tests were made in case of most of the samples to determine the presence or absence of Bacillus coli-communis or the group of organisms commonly associated under that name. In these from two to five series of tests were made for each sample, each usually having five distinct processes requiring skillful manipulation and discriminating judgment. Thus for the 2,800 samples there have been some 30,000 cultures and as many entries of the results.

THE PURPOSES OF THE INVESTIGATIONS.

The objects in all this were to ascertain the number of bacteria existing in the water at different times at the different collecting stations located at favorable points throughout the course of the drainage waters from Chicago to St. Louis, to determine as far as possible the facts in regard to the sources of contamination of these waters, and those in regard to natural purification, as well as to gather data showing the distribution in the waters of special kinds of bacteria associated with certain diseases of human beings. It was, in a word, to show what effects the sewage of Chicago has upon the stream throughout its course for 368 miles.

It should be noted that the work began in the early summer of 1899, eight months before the water was permitted to pass through the Sanitary Canal then in process of construction, and continued nine months after this great waterway was opened. Comparisons for the months of June, July and August for the two years, 1899 and 1900, before and after the opening of the Sanitary Canal, are especially noteworthy and are abundantly presented hereafter.

In order to determine in a general way the contaminations reaching the waters of the Illinois river besides such as came from the city of Chicago, samples were continuously collected from the Kankakee river at Wilmington, from the Fox river at Ottawa, from the Big Vermilion river near LaSalle, and from the Sangamon river near Chandlersville. All of these streams, with other tributaries of the Illinois river, receive the sewage of large aggregate populations and must have considerable effect upon the character of the water into which they discharge. For further comparative studies collections were also made from the Mississippi river above the entrance of the Illinois at Grafton and from the Missouri river at West Alton, and as incidental to the main work in hand, analyses are reported of regularly collected samples of Chicago and St. Louis tap waters.

COLLECTING STATIONS AND SHIPMENTS.

The collecting stations were selected and the methods of taking and shipping the samples for chemical and for bacteriological analyses were determined conjointly by Professor Arthur W. Palmer of the University of Illinois and Professor Edwin O. Jordan of the University of Chicago, as elsewhere described by them in this volume. For ready reference, those on the course of the stream, with their approximate distances from the pumping station at Bridgeport, Chicago, are here named in order of their occurrence:

Lockport, Michigan Canal, Desplaines river, and Sanitary Canal. 29 miles.

Joliet, Jackson street, Desplaines river, 33 miles.

Joliet, Rock Island Bridge, Desplaines river, 36 miles.

Morris, Illinois river (9¹2 miles below union of Kankakee and Desplaines), 57 miles.

Ottawa, Illinois river, 81 miles.

La Salle, Illinois river, and Illinois and Michigan Canal, 95 miles.

Henry, Illinois river, 123 miles.

Averyville (North Peoria), Illinois river, 159 miles.

Wesley City, Illinois river (4 miles below Peoria), 165 miles.

Pekin, Illinois river, 175 miles.

Havana, Illinois river, 199 miles.

Beardstown, Illinois river, 231 miles.

Kampsville, Illinois river, 288 miles.

Grafton, Illinois river, 318 miles.

Grafton, Mississippi river.

Alton, Mississippi river, 400 feet from Illinois shore, 333 miles.

Mton, Mississippi river, 4 width from Illinois shore.

Alton, Mississippi river, center of stream.

Alton, Mississippi river, ¼ width from Missouri shore.

Alton, Mississippi river, 400 feet from Missouri shore.

Mitchell, Mississippi river, 400 feet from Illinois shore, 348 miles.

Mitchell, Mississippi river, center of stream.

Mitchell, Mississippi river, St. Louis Inlet Tower.

Mitchell, Mississippi river, 400 feet from Missouri shore.

West Alton, Missouri river.

Jefferson Barracks, Mississippi river, 400 feet from Illinois shore, 368 miles. Jefferson Barracks, Mississippi river, ¼ width from Illinois shore.

Jefferson Barracks, Mississippi river, center of stream.

Jefferson Barracks, Mississippi river, ¼ width from Missouri shore. Jefferson Barracks, Mississippi river, 400 feet from Missouri shore. Chicago tap water.

St. Louis tap water.

The samples were taken at these stations according to strict instructions, packed in ice, and forwarded to the laboratory as speedily as possible. Further description of these processes is not given, because a full account has been made by others in this report. It should be said, however, that notwithstanding the great pains taken to prevent unnecessary delays, these sometimes occurred, and the samples were not received in good condition. At first, before collectors and express agents came to understand the necessities in the case, there was much trouble of this kind, though subsequently little complaint could be made. The figures for May, 1899, are not included on account of errors due to the causes just referred to.

Throughout the entire period, however, the method of sending samples for plating was, under the best possible conditions which could be devised, an unsatisfactory substitute for work upon the spot and at the instant collections were made. It is a fact that in case of most of the waters examined the numbers of bacteria had been at a maximum some time previous to the date of collecting, that the water had been in most cases undergoing the natural processes of purification, with an accompanying diminution of the number of living bacteria. When, therefore, the conditions were made unfavorable for the multiplication of bacteria, as by the want of aeration and low temperature, the rate of diminution was necessarily greater. Hence transported samples ordinarily gave a less number of colonies in the culture dishes than plating direct from the water at the collecting station would have shown. On the other hand, some special species, well adapted to the new conditions, undoubtedly did multiply rapidly in the water after it was enclosed in the collecting bottles. This sort of increase must have been more striking as the length of time was increased, because these special kinds were freed from the competition of the others. Any water rich in fresh organic matter would also offer an opportunity for notable increase in the bacteriological contents, especially under favorable conditions, during 24 or more hours in a collecting bottle. Thus, while our numbers are commonly less, they are sometimes presumably greater than place-plating would have shown.

For these reasons, as well as for variations always encountered in work of this kind, the figures herewith reported must be considered approximate ones only, but the errors are not believed to be sufficiently serious to affect the comparative results. In a very few instances counts widely deviating from those commonly obtained from given samples have been thrown out in preparing the tables, but in no case have numbers been otherwise modified. The figures are reported just as they were entered in the laboratory books.

PLATING FOR COLONY COUNTS.

Nutrient agar-agar was chosen instead of gelatin, more commonly used for colony counts, because upon the whole the former was believed to be the better adapted to the particular work in hand under the special conditions existing. It is true that more bacteria commonly develop in gelatin, but it is gravely questioned that better comparative results can ever be obtained by the use of this medium in the study of such waters as were here dealt with. The fact that liquefaction often takes place through the energetic action of some rapidly developing species before the slower growing kinds have grown sufficiently to be visible, and therefore before a proper count can be made, is in itself a very serious drawback to the use of gelatin. With it there can be less uniformity in the length of time allowed for the plates to develop before counting, hence there is a kind of irregularity introduced, due in part to the particular species or association of species one happens to have to deal with in a particular plate, which are not the same in another plate used for comparison. Anything which compels the laboratory employee to watch the progress of individual cultures and to modify his own action in accordance therewith tends to defeat the comparisons for which alone his work is valuable. It is impossible ordinarily, under the best conditions, to leave a gelatin plate more than four days before counting, often not more than half this length of time. The slower growing species may not appear at all within these limits, while others are destroying the plate by liquefaction. In the case of agar the growth is commonly less rapid, but after a colony is once formed it does not disappear except by indefinitely spreading, and even those overrun can often be seen at the end of a chosen uniform period of sufficient length to permit the best development for counting. As uniformity of results means in this work much more than a maximum growth from one or more samples, nutrient agar has claims for consideration aside from convenience of handling in hot weather.

The plating agar was made from butchers' meat with Witte's peptone, titrated against sodium hydroxide to the phenolphthalein neutral point, and brought by means of hydrochloric acid to 10 on Fuller's scale. Care was taken to have the medium several weeks old before use, as a means of avoiding spreading growths, and to secure better pouring quality. The prepared plates were kept in a basement room of a large building, where the temperature was practically constant at 20° C. This was found to be much more satisfactory than keeping the cultures in a box with a constant temperature maintained by running water. The plates were uniformly counted after a development period of ten days. When

					E /
t - Illinois and Michia	gan Canal				
Joliet - De	esplaine's Rive	r			
Morris —	Illinois River				
River					
River					
nois River					
	s River				
			repre	repre per c	(
	9		Length sents the	Lengi sents av	
	niversity	PL	. =	th of line erage nu imeter of	
			2	s shown mber of water	
Missouri River		0		thus: I bacteria for June	
			ctoria	1899.	
	6- 0,00	800,00		- 1	
	Joliet D Morris River River inois River	Morris — Illinois River River River ey City — Illinois River — Illinois River inois River inois River River River River River River River River	Joliet - Desplaine's River Morris - Illinois River River	Joliet Desplaine's River Morris - Illinois River River	Joliet - Desplaines River Morris - Illinois River River

helpful a lens was used in counting. Such dilutions were made of the water as were found by repeated trials necessary to secure from 50 to 100 colonies upon a Petrie dish 100 mm. in diameter.

BACILLUS COLI-COMMUNIS.


Much attention was given to the so-called colon group of bacteria. This investigation was undertaken with a realizing sense of the difficulties sure to be encountered in obtaining trustworthy results and of the care needful to avoid serious errors. In the first place, much variation is known to occur in cultures, and in the characteristics relied upon for identification, due to the special type in hand, whether or not these variations are considered to be of specific value when found reasonably permanent and sufficiently recognizable under variations of conditions. Then the previous history of the special organisms dealt with and the immediate conditions of growth lead to much difference in results. From inherent peculiarities and from external causes combined the investigator must anticipate considerable departure from any standard adopted for the characterization of this colon organism or organisms. It is, however, a matter of the greatest importance in such studies as this that the meaning expressed by figures in reports shall be clearly interpretable and open to critical review. With this in view, the processes made use of for the identifications reported are here described in considerable detail.

After some preliminary changes in methods a routine of procedure was adopted and afterward adhered to as follows:

(1) One cubic centimeter of the water to be examined was placed in each of at least two tubes containing a culture medium made up of:

5 grms. Liebig's meat extract.
10 grms. chemically pure lactose.
1 grm. pure carbolic acid crystals.
1,000 cc. distilled water.
Litmus, aqueous solution sufficient to color.

These were incubated at about 38° C. for 48 hours. If the color remained unchanged a negative result was at once interpreted, since the conditions were favorable for the growth of the organism in question and though very few were originally contained in the water, multiplication would rapidly take place, while the restraining influences of the carbolic acid and the temperature would keep in check the ordinary bacteria. In the presence of sugar, colon bacteria produce acid, and this is quickly indicated by the litmus. If one or both the tubes became red separation cultures were made in nutrient gelatin containing lactose and litmus as in the medium just described. Indications of the colonies of the organism were supposed to be given by the change of color in the immediate vicinity, and selections were then made by the other characteristics for pure cultures. These were made in sugar-free and in 2% glucose broth—the former for indol tests and the latter for gas in the fermentation tube and in litmus milk.



Sterile meat infusion was made sugar-free by inoculating with typical *Bacillus coli-communis* and incubating for 14 to 16 hours. It was then boiled, filtered, and sterilized in tubes. Cultures for indol tests were incubated 36 to 48 hours, after which about 1 cc. of .04% solution of sodium nitrite and 4 to 6 drops of concentrated sulphuric acid were added. If the red color did not immediately appear after shaking, the tubes were left for a time in the incubator.


In the fermentation tubes 30 to 60 per cent of gas was considered indicative of the colon bacillus, but in practice a positive record was made if all other tests were favorable, even though within somewhat varied limits a greater or less amount of gas appeared. The upper limit was, however, rarely surpassed when other characteristics agreed in indicating the colon bacillus. The lower limit proved more variable, due perhaps to the varietal peculiarity of the organism or to its immediately preceding history.

When all of the tests could be undoubtedly interpreted as showing the presence of the colon bacillus a positive record was entered; when some of them clearly failed and further trials resulted in the same way, a negative entry was made. If, however, as sometimes happened, the tests were contradictory or uncertain, the record was made to show this condition of things. In the tables following the signs + and — show the presence or absence respectively as thus ascertained of the organism or organisms in question, and an interrogation point indicates doubtful results.

No animals were inoculated to test the pathogenic properties of any isolated organisms. This was impracticable upon the scale upon which the work was done. Further, there was little to be gained by the process. The contaminations of these waters by germs pathogenic to man are presumably confined to those producing intestinal disorders, and these show very great variability in regard to virulence to animals widely different from their effects upon man. For instance, no added information would be probably gained in regard to the distribution of *Bacillus typhosus* by inoculating animals with cultures derived from the waters under examination.

RESULTS IN REGARD TO THE NUMBERS OF BACTERIA.

A glance at the tabular exhibits presented herewith, and better by the graphical showing, reveals at once the very striking differences which exist in the bacterial content of these waters in the different portions of the stream. There is everywhere much variation for which there are several well recognized causes. There are seasonal differences to which attention is further to be called; there are changes determined by rains and the wash from the shores of the main channel and of the entire tributary systems, and these, of course, are irregular in time and variable in extent. There are also changes brought about by artificial processes of such nature that the stream received at one time more than another decomposable matter, as the wastes from manufactories and



the offal from stockyards. But in spite of any such variations the numbers of bacteria are always enormous at Bridgeport and at other headwater stations. They decrease with marked regularity as one follows down stream to Peoria. The place called Averyville is just above this city, where the pumping works are located by which the place is supplied with water. This is 159 miles from Bridgeport, the other pumping station operated for the disposal of the sewage of the city of Chicago. Taking, for example, the number of bacteria per cubic centimeter in the water at these places during November, 1899, and April, 1900, our analyses gave the following figures:


Bridgeport		Nov. 15. 1,960,000 1,100	Nov. 21. 3,920,000 1,200	Nov. 29. 315,000 2,800	Average. 2,497,000 1,640
	April 3.	April 10.	April 17.	April 25.	Average.
Bridgeport	5,300,000	3,725,000	11,200,000	3,925,000	6,037,500
Averyville	28,000	17,000	5,500	1,850	13,087

The river receives great quantities of polluting matter as sewage and other refuse from the cities of Peoria and Pekin, and the numbers of bacteria again very greatly increase, followed as before by a notable decrease further down stream. Beardstown is 56 miles south of Pekin. A comparison of results obtained from these two places illustrates again the self-purification of the flowing water:

	Nov. 1.	Nov. 9.	Nov. 16.	Nov. 24.	Average.
Pekin	2,780,000	170,000	145,000	95,000	797,500
Beardstown		21,500	. 7,150	35,200	17,590
	July 6.	July 12.	July 19.	July 26.	Average.
Peklu	38,500	555,000	370,000	220.000	295.875
Beardstown	3,500	4,000	3,000	6,000	4.125

The dates vary by one day in some of these cases in regard to one of the parts of the couples from those on which the tests were made, but this does not affect the lesson gathered by the comparison. The averages in the last set of figures show respectively 45 and 71 times as many bacteria in the water at Pekin as at Beardstown, though the distance is but 56 miles and the time required for the water to flow from the former to the latter place is not more than 24 hours.

The kind of information gained from the figures just quoted can be much more satisfactorily obtained by a study of the graphic presentations in plates XXI to XXVIII, inclusive. These cover the collections made during June, July, August and September of the years 1899 and 1900, the corresponding portion of the two years upon which collections were made. The data upon which the plates are constructed come from the averages of all the counts made during the month indicated in each case. These particular months were chosen because they were the ones, and the only ones, during which the work was in progress both years covered by this report, and the comparative annual studies are important as will be shown later. The counts for other months of the year would have given similar exhibits though subject to seasonal variations. The two plates for each month are only different methods of presenting to the



eye the same information; but those with connected, oblique lines do not reach northward or up stream above Ottawa, because of the difficulty of representing the larger counts on a scale suitable for the smaller ones. This difficulty is met in the plates with vertical lines by doubling the long ones upon themselves; those representing the counts for 1899 being heavy or solid and those for 1900 of lighter and open construction. On these plates the distances in miles from one station to that next succeeding are given along the bottom.


The story graphically told upon the plates is the same for each of the four months for either year. It is for 1899 (the heavy or whole lines): (1) the enormous initial numbers represented by the long lines reaching two to four times the vertical width of the plates, (2) the phenomenal decrease between Morris and Ottawa and the surprisingly low showing through to Averyville, (3) the sudden and great increase at Wesley City and Pekin, and (4) the marvelous drop at Havana and onward. In all this the data for the four months give in effect the same exhibit in all of the cases; though September stands somewhat by itself. For this month, Lockport, Joliet, Wesley City and Pekin have much shorter or lower lines than usual, indicating that the numbers of bacteria in the samples taken at these places were much less during this month than usual or than were found for the other three months. In this connection the following figures, showing the counts for these places from August to December, 1899, inclusive, are of interest:

	Ill. and M. Canal	Joliet.	Wesley City.	Pekin.
August	3,215,000	3,220,000	2,355,000	1,928.000
September	669,600	217,000	484.500	391,250
October	1.359,375	1.145,000	2,373,750	1,047.500
November	582,000	1,023,000	705,000	812,000
December	746,250	1,026,250	20,600	69,000

The numbers are all very large. The variation usually is not more than is ordinarily anticipated, except those for December at Wesley City and at Pekin. These, compared with the others given and with those usual for these stations, are very exceptional and must have some special explanation. Upon the latter, however, it is idle to speculate. Let it be remembered that there is no connection between the first two sets of figures and the last two sets, for Joliet and Wesley City are separated by a long line of the waterway, from which the samples show very low counts.

The main lesson in this respect offered by the tables is so important that another set of figures may be given for comparison with those just presented. The Ottawa station is 48 miles from that of Joliet, and Averyville is 74 miles from Ottawa. Havana is 33 miles from Pekin and Kampsville is 81 miles from Havana. The average counts for these places for the same time as above were as herewith given:

Ottawa.	Averyville.	Havana.	Kampsville
August 7.500	4,000	5,700	6,200
September 40,100	3.100	6.137	3,480
October 33,050	3,770	12,430	1,717
November 24,750	2,800	164,000	6,850
December264,760	9,300	301,300	28,850



From Joliet to Ottawa in August the number drops from 3,220,000 to 7,500 and in November from 1,023,000 to 24,750. From Pekin to Havana in August the decrease is from 1,928,000 to 5,700, and in October from 1,047,000 to 12,430. In general the figures in this last set are decidedly less than those of the first set. And let it be noticed those from Averyville are less than those from Kampsville, the average of the former being 4,614, and of the latter 9,420. This is not always the case, as is shown by the graphical plates, but the fact is emphasized that the pollutions of the headwaters near Chicago have very largely disappeared at Averyville. The Kampsville water would certainly have been still better but for the new contaminations at Peoria and Pekin.

It is worth while to ask attention to the results indicated on these plates concerning the relative numbers of bacteria in the water of the Illinois river at its mouth and the Mississippi river before it receives the Illinois. Both these collecting stations are marked Grafton. Appealing to the charts with connecting oblique lines, XXI, XXIII, XXV and XXVII, it is seen that the Mississippi has a slightly less number for June of both years, and the same is true for July both years; for August it is the same for 1899, but the reverse for 1900, while for September the relation stated holds for 1900, but in 1899 the Illinois has fewer bacteria. In every case the numbers as found in the samples from West Alton on the Missouri river are considerably above those of either the Illinois or the Mississippi at Grafton. The zigzag lines of the plates usually tend upward for some cause to Alton, where the samples were taken from the mixed waters of the Mississippi and the Illinois rivers but above the mouth of the Missouri, and the upward slant of these lines from Alton to Mitchell not only always occurs, but is considerable for each month shown. At Mitchell the waters of the Mississippi and Missouri have become mingled, and our exhibit is based upon the average number of bacteria found in five samples taken at somewhat equal intervals on a line across the combined stream. The increase in number of bacteria from Alton to Mitchell is apparantly explained by the greater number in the Missouri, as shown by the samples from West Alton.


For the sake of further comparison in regard to the waters of the Illinois river at its mouth and those of the Missouri at West Alton, it may be well to bring together the monthly averages for the entire time over which this report extends. For 1899 they are as follows:

Jun	e. July.	August	September.	October,	November.	Dec.
Illinois 4,1	55 2,628	1,800	793	743	7,200	40,350
Missouri	20,000	32,700	23,400	16,400	28,500	10,500

For 1900 the records, when averaged for each month, give the following numbers:

	Jan. Fe	b. March.	April.	May.	June.	July.	Vug.	Sept.
Illinois	6,600 - 191.	500 159,500	11,550	6.850	6.850	3,900	2.700	5.270
MissouriS	4.960 - 40,	000 - 179,750	77,000	66,200	63,370	30,170	12,450	25,200

It will be seen that only in two cases are the numbers greater for the Illinois river—the numbers of D cember, 1800, and January,



1900. Otherwise the counts from the waters of the Missouri are largely in excess—from twice to more than ten times those from the Illinois. From what has gone before it is plain that about the same relation holds between the number of bacteria in the waters of the Missouri river at West Alton and those of the Mississippi at Grafton.

The waters of the tributaries examined do not, upon the whole, show so many bacteria per cubic centimeter as are found in the Illinois river near the meeting place of these waters. The Kankakee joins the Desplaines between Joliet and Morris. The samples taken at these two stations proved to have many times the number of bacteria of those from the Kankakee at Wilmington. This is not unexpected, for the waters at the former places are very highly polluted. From the Kankakee from 2,000 to 5,000 bacteria per cubic centimeter were usually found during the summer months and from 25,000 to 100,000, or above, from the Illinois at Morris.

The monthly averages for June, July and August of 1899, and for January, March and April, 1900 (no collections from the Fox in February) from the Fox and Illinois rivers at Ottawa are as follows:

	June.	July.	August.	January.	March.	April.
Fox	11,000	3,070	4,560	12,275	84,160	49,250
Illinois	20,550	9,300	7,500	256,250	116.750	68,500

The numbers from the Big Vermilion and the Illinois rivers at La Salle show essentially the same proportions as the above, and those from the Sangamon at Chandlersville and the Illinois at Beardstown are not far different.

VARIATION IN NUMBERS OF BACTERIA FOLLOWING THE SEASONS.

Any inspection of the figures presented must quickly make evident that the numbers of bacteria found in any given place vary greatly according to the season of the year. This is more pronounced at considerable distance from the source of pollution, and is readily explained, though at first it might be thought the natural tendency would be directly opposite the order shown in these studies. Consult, for instance, the last column in tables 57, 58 and 59, showing results from Chandlersville, Beardstown and Kampsville. Here the largest counts are for the winter months, and the difference is very great. The average of the counts as given on table 58 for Beardstown for January, February and March, is 268,600, while for June, July, August and September it is 8,100. Table 68, occupying four pages, shows the seasonal variation for the Illinois river at Grafton in a still more instructive manner with the largest numbers when the temperature of the water is the lowest.

What causes this remarkable difference? If we turn to tables r and 40, showing the numbers of bacteria in the Illinois and Michigan Canal at Bridgeport, no such variations will be found. At Jackson street, Joliet, table 4, the counts average for June, July, August and September


Lockport - Sanitary Canal Joliet - Desplaines River	Ilinois and Michia	gan Canal	
Ottawa - Illinois River La Salle - Illinois River Henry - Illinois River Averyville - Illinois River	River		
Wesley City — Illinois Pekin — Illinois River Havana — Illinois River Beardstown — Illinois River Kampsville — Illinois River Grafton — Illinois River	Bacteriological University of	for	represents average number of per cubic centimeter of water
Grafton — Miss. River West Alton — Missouri River Alton — Miss. — River Mitchell — Miss. River	XXVIII Laboratory ————————————————————————————————————	č o	shown thus:

2,459,400, but for November and December the average is 1,067.500, showing the greater numbers when the water temperatures are very high. This is no contradiction. The fact is the numbers of bacteria are small during warm weather at a distance from the source of pollution because the numbers are then large at this source. In other words, the ordinary purification of water contaminated with organic, fermentable matter, is largely due to the fermentive action of the bacteria themselves, and they are short lived. When the conditions are favorable for their rapid multiplication they more quickly dispose of the contaminating matter, and when their food is gone they perish. The lower waters contain fewer bacteria in warm weather because the organic substances are destroyed in the upper portions of the stream by the great numbers of bacteria there existing and working. When, however, this activity is less in the headwaters the fermentable material is carried farther along by the current, and the bacterial development similarly extends farther down stream. A barrel full of rain water becomes foul sooner in warm weather, but it also then purifies itself quicker. The bacteria multiply faster, hence become more numerous for a certain length of time, but they subsequently become fewer as a direct cause of this early development.

COMPARATIVE NUMBERS OF BACTERIA IN 1899 AND 1900.

The chief interest in this comparison lies in the fact that the Sanitary Canal was opened in the early part of the latter year. What effect has the larger intake of water from the lake had upon the bacterial content of the stream in any part of its course? We have just seen that an answer to this question cannot be had by comparing the counts of different seasons of the year. It cannot, indeed, be said that any difference noted for the same season in different years is due to any one cause, or in this case that the opening of the canal made all the variation observed between the results of the corresponding months of June. July, August and September of these two years. It is, however, plainly evident from a study of the figures that the increased volume of pure water had a decided influence upon the bacterial counts from the collecting stations of the upper part of the stream. Plates XXII, XXIV, XXVI and XXVIII bring this out in a striking manner. The solid lines (1800) for Lockport. Joliet and Morris are from seven to 75 times as long as the open lines (1900). The only exception is for September at Lockport, when the water was then practically stagnant at the time in the old Illinois and Michigan Canal.

From Ottawa onward the influence of the lake water is not so apparent. Except for the showing at Wesley City and Pekin it cannot be said that the difference between the two years as shown in plates XXI to XXVIII is anything beyond what might easily occur without aid from man. Neither is it certain that the difference at Wesley City and Pekin is principally due to the opening of the canal. Some changes in the disposal of the wastes from the distilleries and glucose works and of the wash



from the cattle sheds might easily account for the change, pronounced as it is. It is, however, true that the volume of water passing Peoria was largely greater for the months compared in 1900. The difference in average height of the water gauge as given for Averyville, tables 14 and 53, for the two years is over two feet. As the water is practically pure at this point during the summer, the sewage of Peoria is diluted somewhat in the same way as that of Chicago in the Sanitary Canal, and the tables clearly indicate this by the reported number of bacteria for June, July and August. This leaves unexplained, however, the different showing for September. Compare the heavy and open lines on plates XXVI and XXVIII for Pekin. The cause of the change of relation here shown is not apparent, though the great irregularities in the counts for this place render it not surprising. On table 16 the counts for September, 1899, are 40,000, 445,000, 570,000 and 510,000, though for August 2d and 9th they are 3,635,000 and 3,073,300, and for November 1st and 9th they are 2,100,000 and 2,780,000. Table 55 shows the counts for September, 1900, to be 1,320,000, 600,000, 47,500 and 2,280,000. It is scarcely worth while to try to look for any rule or law in such variations as these, but a reasonable explanation lies in the fact that the pollution all comes at Peoria from the west side and is not soon below uniformly mixed with the waters of the stream.


If there is any effect at all shown at Grafton, which is very doubtful, it is that the numbers of bacteria are slightly greater on account of the increased amount of water in the river. At all events the counts from the mouth of the Illinois were larger in 1900 than they were in 1899, as is clearly shown in plates XXI, XXIII, XXV and XXVII. It is to be noted, however, that the same is true for the Mississippi River at Grafton, except during June, as is shown on the same plates by position of the little squares (1900) and circles (1899). Surely the Sanitary Canal has nothing to do with the bacteria of the Mississippi above the mouth of the Illinois.

In a word, it may be said that these investigations give no demonstration of any change in the bacterial content of the drainage waters from Chicago traceable to the opening of the Sanitary Canal, except as applied to the stream above Ottawa. This refers, however, to the warm season of the year. In the winter the limiting point may be farther down stream.

BACILLUS COLI-COMMUNIS.

The methods used for the identification of this organism, or group of organisms, have heretofore been given. In next to the last column in tables I to 80 inclusive, the results for all the samples tested have been indicated by the signs +, —, and ?. The first means that the organism was present, the second that it was absent, and the third that the test was inconclusive.

With rare exceptions there are no negative results from samples taken above Ottawa. From the tests at Morris in 1899 there are 14 posi-



tive, no negative and 2 questionable, and in 1900 there are 23 positive, 3 negative and 11 questionable results. Thus, out of 53 analyses there were only two samples in which it could be affirmed the bacillus was absent. At Ottawa, however, a change begins. In 1899 the samples from the Illinois river at this place gave 10 positive, 7 negative and 1 doubtful tests, and in 1900 there were 11 of the first, 17 of the second and 5 of the third, or together 21 positive, 24 negative and 6 doubtful. At Henry there are 22, 30 and 5, in the same order; and at Averyville, the count shows 19, 21 and 2. Then at Wesley City and Pekin we see the same exhibit as at Bridgeport and at Lockport, to change again at Havana in such manner that we have 31, 12 and 14; at Beardstown, 37, 18 and 3; at Kampsville, 35, 18 and 2, and from the Illinois at Grafton, 31, 23 and 1.

If we compare with this the Mississippi at Grafton (above the Illinois) and the Missouri at West Alton, we have:

	+	-	?
Illinois	31	23	1
Mississippi	35	20	1
Missouri	36	7	7

As will be seen, the organism was present most often in the waters of the Missouri river, next in those of the Illinois, and least frequently in those of the Mississippi, if the whole time of the investigation is considered. The first seems to be generally true; the relation of the second and third changes at times.

An attempt is made to present graphically the results of the investigation on plate XXX. For this purpose the numbers were reduced to per cents of the whole number of tests made for each collecting station. The connecting lines show the results for 1899 and for 1900 separately and for the two years combined. It was not deemed important to include the stations above Joliet, because the per cent in almost every case was 100.

It will be noticed that the line for 1899 begins above that for 1900, but drops below at Henry and remains below to Havana, then crosses again at Beardstown, to remain above to Mitchell.

The explanation of this to Averyville is, no doubt, to be sought in the opening of the Sanitary Canal. The dilution of the sewage is effective throughout and controls everything to some point below La Salle, where other influences, like the greater current and the better conditions for the multiplication of this particular bacillus, overcome and make the showing of the chart.

The lines of the plate nowhere descend again so low as they are at Averyville. From this point on, local contaminations keep the percentages high, falling, however, from Kampsville to Grafton, then rising to Mitchell. This last seems clearly due to the influence of the Missouri, whose waters have, as has just been shown, a greater proportional number of the organisms.

SPECIES DETERMINATIONS.

The main efforts in this work were upon the matters above reported, though considerable attention was at one time or another given to the identification of the various species of bacteria procured in the cultures. It may be remarked that *Bacillus typhosus* was not recognized, though it is presumable that it was sometimes present in the waters examined. One thing was plainly evident, viz., that at a station like Grafton, considerably distant from a source of much contamination, the colonies of bacteria grown in a plating dish were of much fewer species than were those from water having fresher pollution. There had been a survival of the fittest, and, as a rule, only those best adapted to the changing conditions, or to the ultimate ones, survived. It would be contrary to all information on the subject to include theoretically among these any of the principal bacteria pathogenic to man.

CONCLUSION.

These studies in detail and as a whole tend strongly to show that neither before nor after the opening of the Sanitary Canal had the drainage waters from the city of Chicago for the period covered by this report any appreciable effect bacteriologically upon the waters of the Mississippi river. Even if the waters of the Illinois river at its mouth were proved to be a source of contamination to those of the Mississippi, the above statement would still seem to hold good, because of the conditions shown to exist at Averyville and Pekin. It is still more clearly apparent that Chicago sewage cannot be held responsible for the contaminations existing in the commingled waters of the Mississippi and Missouri Rivers.

T. J. BURRILL.

Professor of Botany in Charge of Bacteriology, University of Illinois.



APPENDIX



ERRATA.

TABLE 120.

Serial No. 1301; Jan. 28, No. Bacteria per cc. for 540,000 read 340,000.

Serial No. 1397; Feb. 19, Temperature of Air for o. read — .5.

Serial No. 1673; April 16, Temperature of Air for 16. read 17.

Serial No. 1712; April 23, Temperature of Air for 13. read 15.

TABLE 121.

Serial No. 1563; March 26, Temperature of Water for -5. read -.5.

TABLE 122.

Serial No. 1598; April 2, Albuminoid Am. Dissol'd for .364 read .264.

TABLE 123.

Serial No. 1788; May 8, Temperature of Water for 16.5 read 15.5. Serial No. 1896: May 30, Albuminoid Am. Susp'd for .08 read .04.

TABLE 126.

Serial No. 1305; Jan. 29, Temperature of Water for 2. read -2.

Serial No. 1306; Jan. 29, Temperature of Air for 8, read —8.

Serial No. 1399; Feb. 19. Temperature of Water for 1. read —1.

Serial No. 1674; April 16, Chlorine for 2.2 read 3.2.

Serial No. 1751; April 30, Res. on Evap. Suspend'd for 4. read 7.

TABLE 127.

Serial No. 1338; Feb. 5, Chlorine for 15. read 25.

Serial No. 1758; April 30, Color for .14 read .4.

Serial No. 1822: May 15, Nitrogen as Nitrates for .4 read .41.

TABLE 128.

Serial No. 1681; April 17, Albuminoid Am. Suspend'd for .189 read .184.

TABLE 129.

Serial No. 1250; Jan. 18, Temperature of Water for 2. read 3.

Serial No. 1490; March 12, Albuminoid Am. Suspen'd for .032 read .232.

Serial No. 1604; April 3, Oxygen Consumed by Dis'd for o.8 read 6.8.

Serial No. 1791; May 8. Nitrogen as Nitrites for .8 read .18.

TABLE 131.

Serial No. 1969; June 12, No. Bacteria per cc. for 3,900 read 9,900.

TABLE 132.

Serial No. 1726; April 24, Residue on Evap. Total for 344 read 394.

TABLE 133.

- Serial No. 1401; Feb. 19, Temperature of Air for 12. read -12.
- Serial No. 1433; Feb. 26, Temperature of Air for 13. read -13.
- Serial No. 1456; March 5, Temperature of Air for 11. read -11.
- Serial No. 1562; March 26, Temperature of Air for .3 read 3.

TABLE 134.

Serial No. 1275; Jan. 23, Nitrogen as Nitrites for .1 read -

TABLE 135.

Serial No. 1974; June 13, Color for .3 read .2.

TABLE 137.

- Serial No. 1314; Jan. 30, No. Bacteria per cc. for 63,000 read 63,400.
- Serial No. 1971; June 13, Nitrogen as Nitrates for .06 read .6.
- Serial No. 1971; June 13, Temperature of Air for 33. read 23.

TABLE 138.

- Serial No. 1384; Feb. 14, Oxygen Consumed, Total for 5. read 2.5.
- Serial No. 1731; April 25, Res. on Evap. Dissol'd for 274. read 276.
- Serial No. 1945; June 6, Albuminoid Am. Susp'd for .224 read .244.

TABLE 139.

- Serial No. 1428; Feb. 22, Nitrogen as Nitrates for 1.15 read 1.65.
- Serial No. 1440; March 1, Albuminoid Am. Diss'd for .252 read .232.

TABLE 140.

- Serial No. 1354: Feb. 7, Free Ammonia for 1.8 read 1.54.
- Serial No. 1354; Feb. 7, Albuminoid Am. Total for .154 read .432.
- Serial No. 1354; Feb. 7, Albuminoid Am. Diss'd for .432 read .216.

TABLE 142.

- Serial No. 1188; Jan. 4, Chlorine for 6.27 read 3.1.
- Serial No. 1218; Jan. 10, Chlorine for 4.67 read 2.3.

TABLE 143.

- Serial No. 1277; Jan. 24, Temperature of Air for 10. read —. Serial No. 1348; Feb. 7, Temperature of Air for 10. read —.
- Serial No. 1441; March 2, Height of Water for 3.4 read 3.
- Serial No. 1503; March 14, Nitrogen as Nitrites for .018 read .01.
- Serial No. 1539; March 21, Res. on Evap. Susp'd for 839, read 837.
- Serial No. 1773; May 2, Chlorine for 0.8 read 6.8.

TABLE 144.

- Serial No. 1278; Jan. 24, Res. on Evap. Susp'd for 328. read 238.
- Serial No. 1420; Feb. 21, Height of Water for 5.5 read 5.08.
- Serial No. 1432; March 2, Height of Water for 3.4 read 3.04.
- Serial No. 1466, March 9, Albuminoid Am. Total for .056 read .656.

TABLE 145.

Serial No. 1184; Jan. 4, Albuminoid Am. Susp'd for .096 read .094.

TABLE 146.

Serial No. 1422; Feb. 22, Res. on Evap. Total for 620. read 420. Serial No. 1422; Feb. 22, Res. on Evap. Susp'd for 466. read 266. Serial No. 1422; Feb. 22, Height of Water for 5.5 read 5.08. Serial No. 1444: March 1, Height of Water for 3.4 read 3.04.

TABLE 147.

Serial No. 1186; Jan. 4, Res. on Evap. Total for 216. read 266. Serial No. 1325; Feb. 2, Nitrogen as Nitrates for .7 read .9. Serial No. 1423; Feb. 22, Height of Water for 5.5 read 5.08. Serial No. 1445; March 2, Height of Water for 3.4 read 3.04.

TABLE 148.

Serial No. 1359; Feb. 8, Height of Water for 2.7 read 2.4. Serial No. 1700; April 19, Albuminoid Am. Susp'd for .154 read .104.

TABLE 149.

Serial No. 1360; Feb. 8, Height of Water for 2.7 read 2.4.

TABLE 150.

Serial No. 1361; Feb. 8, Height of Water for 2.7 read 2.4. Serial No. 1658; April 12, Albuminoid Am. Total for .848 read .846. Serial No. 1658; April 12, Albuminoid Am. Susp'd for .664 read .662.

TABLE 151.

Serial No. 1362; Feb. 8, Height of Water for 2.7 read 2.4. Serial No. 1918; May 31, Oxygen Cons'd by Diss'd for 9.8 read 3.8. Serial No. 1951; June 7, Albuminoid Am. Total for .63 read .68. Serial No. 1951; June 7, Albuminoid Am. Susp'd for .57 read .56.

TABLE 152.

Serial No. 1263; Jan. 18, Temperature of Water for 1. read 4. Serial No. 1766; May 1, Albuminoid Am. Diss'd for .166 read .176.



SANITARY DISTRICT OF CHICAGO

TABULATION FOR THE YEAR 1900

Of the average daily and weekly rate of flow, in cubic feet per minute, through the Illinois and Michigan Canal; through the Main Drainage Channel; through the Chicago River; through the Desplaines River at Riverside; through the Desplaines River at Joliet.

F THE	DAY OF THE WEEK.	Canal. Estimated from 3 daily readings reduced canal lock gauge near Bridgeport pumps.	Flow through Main Drainage Channel. Estimated from 48 half-hour readings daily of Controlling Works near Lockport, Illinois.	Being sum of (I) at Bridge- port and (L) at Lockport 29 miles below, without cor- rection for time intervals.	Flow of Desplaines River at Riverside. Estimated from 2 daily readings on the Riverside Gauge.	Flow at Dam No. 1, Jolict. Being sum of (I) 33 miles above, (L) 4 miles above, and (R) 28 miles above. Without corrections for time intervals.
1899		I,	L.	C.	R.	J.
Dec. 1900	31, Sunday	(44833)	000000	(44833)	(1320)	(46153)
Jan.	1, Monday 2, Tuesday 3, Wednesday 4, Thursday 5, Friday 6, Saturday	44833 45267 50000 46067 43267 35867	0 0 0 0 0	44833 45267 50000 46067 43267 35867	1320 1320 1320 1320 1320 1320	46153 46587 5 1320 47387 44587 37187
(1)	Weekly average	310134 44305	0 0	310134 44305	9240 1320	319374 45625
	7, Sunday 8, Monday 9, Tuesday 10, Wednesday 11, Thursday 12, Friday 13, Saturday	40033	0 0 0 0 0 0	40700 40700 40367 40033 39333 33767 29000	1320 2130 2820 3300 3300 3300 3300	42020 42830 43187 43333 42633 37067 32300
(2)	Weekly average	263900 37700	0	263900 37700	19470 2781	283370 40481
	14, Sunday 15, Monday 16, Tuesday 17, Wednesday 18, Thursday 19, Friday 20, Saturday	38667 38300	0 0 0 (58429) (90400) (107700) (132454)	32067 37300 39300 (97029) (129067) (146000) (167087)	3300 3300 3300 4650 8850 13980 20610	35367 40600 42600 (101679) (137917) (159980) (187697)
(3)	Weekly average	258867 36981	388983 55569	647850 92550	57990 8284	705840 100834
	21, Sunday 22, Monday 23, Tuesday 24, Wednesday 25, Thursday 26, Friday 27, Saturday	. 23167 . 24433 . 24167	(162150) (178081) 231667 220062 216400 234358 229337	(197350) (204481) 254600 243229 240833 258525 252270	30840 26160 2130 17040 17040 14880 14880	(228190) (230641) 256730 260269 257873 273405 267150
(4)	Weekly average	179233 25605	1472055 210293	1651288 235898	122970 175 6 7	1774258 253465
Feb.	28, Sunday 29, Monday 30, Tuesday 31, Wednesday 1, Thursday 2, Friday 3, Saturday	18900 17500 18367 23100 33067	225209 222363 205531 181094 106461 97392 95288	248142 241263 223031 199461 129561 130459 133521	14\$\$0 130\$0 11040 \$640 6660 4650 1980	263022 254343 234071 208101 136221 135109 135501
(5)	Weekly average	172100 . 24586	1133338 161905	1305438 186491	60930 8704	1366368 195195

4, Sunday 5, Monday 6, Tnesday 7, Wednesday	. 45100 . 44433 . 43267	L. 103173 93602 93504 98660	C. 146506 138702 139937 141927	R. 1320 11190 46680 79080	J. 147826 149892 186617 221007
S, Thursday 9, Friday 10, Saturday	. 51333	65033 86277 73548	122700 137610 114315	102360 129120 138690	225060 266730 253005
(6) Weekly average	325900 . 46557	615797 87971	941697 134528	508440 72634	1450137 207162
11, Sunday 12, Monday 13, Tuesday 14, Wednesday 15, Thursday 16, Friday 17, Saturday	38967 41100 33533 27600 28167	126759 147121 162766 186821 211542 149431 116194	168192 186088 203866 220354 239142 177598 152661	60000 97380 73590 58540 40800 28440 21300	228192 283468 277456 278914 279942 206038 173961
(7) Weckly average	247267 35324	1100634 157233	1347901 19 25 57	380070 54296	1727971 246853
18, Sunday 19, Monday 20, Tuesday 21, Wednesday 22, Thursday 23, Friday 24, Saturday	37667 35500 37167 37900 28100	201123 200867 197332 201329 253544 180356 124531	240090 238534 232832 238496 291444 208456 165998	17040 14880 14880 14880 14880 14880 14880	257130 253414 247712 253376 306324 223336 180878
(8) Weekly average	256768 36681	1359082 194155	1615850 230836	1063£0 15188	1722170 246024
25, Sunday 26, Monday 27, Tuesday 28, Wednesday March 1, Thursday 2, Friday 3, Saturday	37300 35167 28767 21467 28700	181839 148271 107675 75966 73333 105125 128671	225472 185571 142842 104733 91800 133825 169071	14880 14880 14880 14880 14880 14880 14880	240352 200451 157722 119613 109680 148705 183951
(9) Weekly average	235434 33633	820880 117269	1056314 150902	104160 14880	1160474 165782
4, Sunday 5, Monday 6, Tuesday 7, Wednesday 8, Thursday 9, Friday 10, Saturday	38233 45700 44000 40033 37200	140917 128577 128444 161869 159031 162511 165421	178084 166810 174144 205869 199064 199711 207254	13080 12180 12180 14880 17040 18300 19800	191164 178900 186324 220749 216104 218011 227054
(10) Weekly average	284166 40595	1046770 149539	1330936 190134	107460 15351	1438396 205485
11, Sunday 12, Monday 13, Tuesday 14, Wednesday 15, Thursday 16, Friday 17, Saturday	50800 46833 46067 -13267 -12167	144340 116371 84517 66725 79239 112114 138035	194073 167171 131350 112792 122506 154281 178402	140100 192870 292350 275820 215580 159360 120960	334173 360041 423700 388612 338086 313641 299362
(11) Weekly average	319234 45605	741341 105906	1060575 151511	139 7 040 199577	2457615 351088
18, Sunday 19, Monday 20, Tuesday 21, Wednesday 22, Thursday 23, Friday 24, Saturday	31333 32000 38967 37933 39000	158802 159696 161988 163596 154212 139488 75869	190702 194029 193988 202563 192145 178488 114202	112740 92580 88200 92580 74700 32220 64440	303442 286609 282188 295143 266845 210708 178642
(12) Weekly average	252466 36067	1013651 144807	1266117 180874	557460 79637	1823577 260511
25, Sunday 26, Monday 27, Tuesday 28, Wednesday 29, Thursday 30, Friday 31, Salurday	40733 34200 37900 38933 24367	75640 87085 102523 98506 156940 142190 132219	110173 127818 136723 136406 195873 166557 132219	131820 146160 159360 159260 129120 70740 58560	241993 273978 296082 289666 324993 237297 190779
(13) Weekly average	210666 30095	795103 113586	1005769 143681	849020 121289	1854789 261970

_						
A	or. 1, Sunday 2, Monday 3, Tuesday 4, Wednesday 5, Thursday 6, Friday 7, Saturday	37866 35800 37533 37867 32033	L. 103054 109310 124448 126400 118800 153479 212588	C. 132954 147176 160248 163933 156667 185512 237555	R. 58560 58560 61440 58560 51360 40300 34800	J. 191514 205736 221683 222493 208027 226312 272355
	(14) Weekly average	235966 33709	948079 135440	1184045 169149	364080 52012	154812 5 221161
•	8, Sunday 9, Mouday 10, Tuesday 11, Wednesday 12, Thursday 13, Friday 14, Saturday	21000 21000 25767	225688 231621 168150 109033 223323 225900 168606	248355 252621 189150 134800 263723 258500 200639	27240 22800 17040 13080 11040 11040 11040	275595 275421 206190 147880 274763 269540 211679
	(15) Weekly average	195467 27924	1352321 193189	1547788 221113	113280 16183	1661068 237296
	15, Sunday 16, Monday 17, Tuesday 18, Wednesday 19, Thursday 20, Friday 21, Saturday	27700 39667 44000 44367 37933 33233 33700	141471 141627 94363 94496 182763 147621 140067	169171 181294 138363 138863 220696 180854 173767	11040 11040 13080 19800 47760 36300 29640	180211 192334 151443 158663 268456 217154 203407
	(16) Weekly average	260600 37228	942408 134630	1203008 171858	168660 24094	1371668 195952
	22, Sunday 23, Monday 24, Tuesday 25, Wednesday 26, Thursday 27, Friday 28, Saturday	27300 32433 29633 28433 • 26100 22600 22200	192048 175069 226250 227883 230681 226869 117606	219348 207502 255883 256316 256781 249469 139806	25080 18300 14880 13980 12180 9840 6660	244428 225802 270763 270296 268961 259309 146466
	(17) Weekly average	188 6 99 26957	1396406 199487	1585105 226444	100920 14417	1686025 240861
Ma	29, Sunday 30, Monday 1, Tuesday 2, Wednesday 3, Thursday 4, Friday 5, Saturday	28333 38567 34500 42967 41533 39333 39667	66316 203792 181096 84421 145242 168375 180413	94649 242359 215596 127388 186775 207708 220080	5340 4380 3960 3960 3720 3000 2640	99989 246739 219556 131348 190495 210708 222720
	(18) Weekly average	264900 37843	1029655 147093	1294555 184936	27000 3857	1321555 188793
,	6, Sunday 7. Monday 8, Tuesday 9, Wednesday 10, Thursday 11, Friday 12, Saturday	35200 34067 39333 40733 34567 32767 32633	205498 222427 176381 111046 188527 193981 232131	240698 256494 215714 151779 223094 226748 264764	2640 2640 8850 18420 12180 6660 6660	243338 259134 224564 170199 235274 233408 271424
	(19) Weekly average	249300 35614	1329991 189999	1579291 225613	58050 8293	1637341 233906
	13. Sunday 14. Monday 15. Tuesday 16. Wednesday 17. Thursday 18. Friday 19. Saturday	24700	233127 213100 220517 205544 144252 125242 89008	261560 239267 248684 238511 180085 149942 113441	5340 4620 3300 1920 1320 360 360	266900 243 87 251984 24043 1 181405 150302 113801
	(20) Weekly average	200700 28672	1230790 175827	1431490 204499	17220 2460	1448710 206959
•	20, Sunday 21, Monday 22, Tuesday 23, Wednesday 24, Thursday 25, Friday 26, Saturday	23167 21233 19733 19267 19500 18433 19267	99144 212544 249867 240102 235671 250554 253910	122311 233777 269600 259369 255171 268987 273177	720 720 720 720 540 540 240 60	123031 234497 270320 259909 255711 269227 273237
-	(21) Weekly average	140600 20056	1541792 220256	1682392 240342	3540 506	16S5932 24084S

May	27. Sunday	1. 19767	L. 261448	C. 281215	R. 0	յ․ 2 812 1 5
mey	28, Monday	20933 21033	145433 265844	166366 286877	60	166366 286937
June	30, Wednesday 31, Thursday 1, Friday	20267	231069 237827 206550	251369 258094 228017	60 0 0	251429 258094 228017
June	2, Saturday		242467	263200	0	263200
(22) Weekly average	144500 20643	1590638 227234	1735138 247877	120 17	1735258 247894
	3, Sunday		249260 241475	269460 259175	0	269460 259175
1	5, Tuesday 6, Wednesday 7, Thursday	21433	250646 201088 170825	268579 222521 196325	1320 8850	268579 223841 205175
	8, Priday 9, Saturday	26400 27300	157029 154396	183429 181696	12960 7320	196389 189016
(23) Weekly average	156466 22352	1424719 203531	1581185 225883	30 450 43 50	1611635 230233
-	10, Sunday	27000 24967	162954 164571	189954 189538	8640 5340	198594 194878
	12, Tuesday 13, Wednesday 14, Thursday	24967 30533 28767	129329 120719 119279	154296 151252 148046	2640 1320 720	156936 152572 148766
	15, Friday	27900 24300	157941 287381	185841 311681	720 540	186561 312221
(24) Weekly average	188434 26919	1142174 163168	1330608 190087	19920 2846	1350528 192933
	17, Sunday	21467 18867 21467	250433 169608 159346	271900 188475	360 360	272260 188835
	19, Tuesday	17500 25067	163110 160248	180813 180610 185315	240 60 60	181053 180670 185375
	22, Friday 23, Saturday	19700 14567	213458 249454	233153 264021	60 6 0	233218 264081
(25)) Weekly average	138 6 35 19805	1365657 195094	1504292 214899	1200 171	1505492 215070
	24, Sunday 25, Monday	23700 23667	251833 238083	275533 261750	60	275593 261750
	26, Tuesday 27, Wednesday 28, Thursday	24167 24700 23933	205644 198083 190850	229811 222783 214783	0 0 0	229811 222783 214783
	29, Friday	24167 25567	198973 141223	223140 166790	0	223140 166790
(26)		169901 24272	1424689 203527	1594590 227799	60 8	1594650 227807
July	1, Sunday 2, Monday	23100 17333	184648 202896	207748 220229	0	207748 220219
	3, Tuesday	18100 17967 14800	200546 199025 205308	218646 216992 220108	0 0 0	218646 216992 220108
	6, Friday 7, Saturday	21967 30500	205019 200629	226986 231129	0	226986 231129
(27)		143767 20538	1398071 199725	1541838 220263	0	1541838 220263
	8, Sunday 9, Monday	30800 31400	190727 197927	221527 229327	60 240	221587 229567
	10, Tuesday	31100 29033 30200	213640 205915 205290	244740 234948 235490	360 360 240	245100 235308 235730
	13, Friday 14, Saturday	32300	130931 200256	163231 233456	60	163291 233456
(28)	Weekly average	218033 31148	1344686 192098	1562719 223246	1320 188	1564039 223434
	15, Sunday 16, Monday	30233	189883 201550	221283 231783	0 0	221283 231783
	17, Tuesday	28733 29333 30500	211560 206975 206252	240293 236308 236752	360 360 360	240658 236668
	20, Friday 21, Saturday	29600 27300	20832 208146 209871	237746 237171	360 360 360	237112 238106 237531
(20)	Weekly average	207099 29586	1434237 204891	1641336 234477	1800 257	1643136 234734
(20)	The transfer seems to the transfer to the tran		20.404.2	201111	201	

-						
Ju	ly 22, Sunday 23, Monday 24, Tuesday 25, Wednesday 26, Thursday 27, Friday 28, Saturday	24700 23467 23200 24700 22233 21933	L. 200935 123255 192008 178331 195755 237198 226275	C. 227335 147955 215475 201531 220455 259431 248208	R. 720 540 540 1680 3960 3300 3300	J. 228055 148495 216015 203211 224415 262731 251508
	(30) Weekly average	166633 23805	1353757 193 3 94	1520390 217199	14040 2005	1534430 219204
Au	29, Sunday 30, Monday 31, Tuesday g. 1, Wednesday 2, Thursday 3, Friday 4, Saturday	21200 21700 21233	306536 228425 241155 207880 240850 194119 237785	328269 249625 262855 229113 261850 215119 258251	2340 1920 1920 1680 1320 1080 720	330609 251545 264775 230793 263170 216199 258971
	(31) Weekly average	148332 21190	1656750 236679	1805082 257869	10980 1568	1816062 259437
	5, Sunday 6, Monday 7, Tuesday 8, Wednesday 9, Thursday 10, Friday 11, Saturday	19733 19733 19733 18167 21200 22667 25033	231208 231096 223982 210927 230473 224427 220169	250941 250829 248715 229094 251673 247094 245202	720 720 720 720 720 720 720 720 540	251661 251549 244435 229814 252393 247814 245742
	(32) Weekly average	146266 20895	157 2 282 224612	1718548 245507	4860 694	1723408 246201
	12, Sunday 13, Monday 14, Tuesday 15, Wednesday 16, Thursday 17, Friday 18, Saturday	21500 18567 22467 27600 24433 24200 25500	227357 236244 184307 222592 234198 224592 235450	248857. 254811 206774 250192 258631 248792 260950	540 720 1920 1320 2640 3000 3720	249397 255531 208694 251512 261271 251792 264670
	(33) Weekly average	164267 23467	1564740 223534	1729007 247001	12860 1837	1741867 248838
	19, Sunday 20, Monday 21, Tuesday 22, Wednesday 23, Thursday 24, Friday 25, Saturday	24500 26133 25833 26400 26700 27000 27000	129400 190541 143017 203079 202686 91113 210832	153900 216674 168850 229479 229386 118113 237832	5340 6660 6420 5700 6330 9240 12060	159240 223334 175270 235179 235716 127353 249892
	(34) Weekly average	183566 26224	1170668 167238	1354234 193462	51750 7393	1405984 200855
Ser	26, Sunday 27, Monday 28, Tuesday 29, Wednesday 30, Thursday 31, Friday ot. 1, Saturday	26400 26700 26400 26400 26400 25533 25500	232565 177879 264086 260223 262398 266485 265900	258965 204579 290486 286623 288798 292018 291400	39630 66130 84780 77220 62700 58740 56400	298595 270709 375 266 363843 351498 350758 347800
	(35) Weekly average	183333 26190	1729536 247077	19128 69 2732 67	445600 63657	2358469 336924
	2, Sunday 3, Monday 4, Tuesday 5, Wednesday 6, Thursday 7, Friday 8, Saturday	25500 24967 25933 27300 25933 28733 28167	252456 243612 125442 121229 82346 47725 68665	277956 268579 151375 148529 108279 76458 96832	45960 39300 33420 28440 23940 21300 18300	323916 307879 184795 176969 132219 97758 115132
	(36) Weekly average	186533 26648	941475 134496	1128008 161144	210660 30094	1338668 191238
	9, Sunday 10, Monday 11, Tuesday 12, Wednesday 13, Thursday 14, Friday 15, Saturday	27300 26667 28167 23333 26100 27000 27000	165942 71558 64819 57927 61302 107181 295962	193242 98225 92986 81160 87402 134181 322962	15840 13980 12180 11840 9360 8160 7080	209082 112203 105166 93000 96762 142341 330042
A	(37) Weekly average	185467 26495	824691 117813	1010158 144308	78440 11206	1088598 155514

Sept.	16, Sunday	1. 25500	L. 240781	C. 266281	R. 6420	J . 27 27 01
	17, Monday 18, Tuesday 19, Wednesday 20, Thursday 21, Friday 22, Saturday	26667 2296 7 23167 26100 25800 23700	1863 5375 7129 6069 31316 94832	28530 28342 30296 321 69 57116 118532	5700 5040 4380 3960 3720 3300	34230 33382 34676 36129 60836 121832
(38)	Weekly average	173901 24843	3873 6 5 55338	561266 80181	32520 4646	59378 6 84827
	23, Sunday 24, Mouday 25, Tuesday 26, Wednesday 27, Thursday 28, Friday 29, Saturday	25200 25333 26400 25800 24967 26733 28200	113663 243056 207265 326791 199540 194238 188071	136863 265389 233665 352591 224607 220971 216271	2640 2640 2640 2640 2340 2120 2120	139503 271029 236305 355231 226947 223091 218391
(39)	Weekly average	180633 25805	1472724 210389	1653357 236194	17140 2448	1670497 238642
Oct.	30, Sunday 1, Monday 2, Tuesday 3, Wednesday 4, Thursday 5, Friday 6, Saturday	27300 25800 26100 25500 25800 27000 27000	260363 257307 260763 262381 259600 258869 260198	287663 283107 286863 287881 285400 285869 287198	1680 1320 1320 1320 1320 1320 1320	289343 284427 288183 289201 286720 287189 288518
(40)	Weekly average	184500 26357	1S19481 259926	2003981 286283	9600 1371	2013581 287654
	7, Sunday 8, Monday 9, Tuesday 10, Wednesday 11, Thursday 12, Friday 13, Saturday	26100 25533 23167 22667 22933 22933 22667	263025 269436 231593 106391 183073 147767 150940	289125 291969 254760 129058 206006 170700 173607	1320 1320 1320 1320 1320 1320 1320	290445 296289 256080 130378 207326 172020 174927
(41)	Weekly average	166000 23714	1352225 193175	1518225 216889	9240 1320	1527465 218209
	14, Sunday 15, Monday 16, Tuesday 17, Wednesday 18, Thursday 19, Friday 20, Saturday	22667 23433 21967 23433 23433 25200 23433	218669 216726 189405 187373 179419 183173 181582	241336 240159 211372 210806 202852 206373 205015	1320 1320 1320 1320 1320 1320 1320	242656 241479 212692 212126 204172 207693 206335
(42)	Weekly average	161566 23081	1356347 193764	1517913 216845	9240 1320	15271 53 21816 5
	21, Sunday 22, Monday 23, Tuesday 24, Wednesday 25, Thursday 26, Friday 27, Saturday	23433 23433 23667 25033 24167 24200 23433	221129 183142 183592 182286 181712 180387 181021	244562 206575 207259 207319 205879 204587 204454	1320 1320 1320 1320 1320 1320 1320	245882 207895 208579 208639 207199 205907 205774
(43)	Weekly average	167366 23909	1313269 187610	1480635 211519	9240 1320	1489875 212839

TABLE I.

		TABLE	I.		
Serial Date. Number. 1899. 526 Aug. 29. 876 Oct. 31. 914 Nov. 7. 950 Nov. 14. 982 Nov. 21. 1011 Nov. 28. 1042 Dec. 5. 1072 Dec. 12. 1110 Dec. 19. 1146 Dec. 28. 1199 Jan. 9. 1239 Jan. 16. 1269 Jan. 23. 1303 Jan. 30. 1334 Feb. 6. 1457 Mar. 6. 1493 Mar. 13. 1522 Mar. 20. 1563 Mar. 28. 1597 Apr. 3. 1636 Apr. 10. 1675 Apr. 17. 1717 Apr. 24. 1753 May 2. 1786 May 8. 1817 May 15. 1856 May 22. 1895 May 29.	.00001 c.c. + - ?	and Michigan .0001 c.c. + - ? 4 0 1 0 1 1 0 1 4 0 2 1 1 1 0 1 2 1 1 3 0 1 4 0 2 2 3 1 2 0 4 0 1 3 2 1 1 1 1 2 3 0 1 2 1 1 1 1 0 1 1 1 1 1 1	I. Canal, Lockpon .001 c.c. + - ? 2 0 2 0 1 0 1 2 1 0 1 2 1 0 1 0 1 0 1 0	1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 c.c. ? 1 0
1923 June 5 1959 June 12 2000 June 19 2032 June 26	$\begin{smallmatrix}0&2&\dots\\0&2&\dots\\\dots&\dots&\dots\end{smallmatrix}$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccc} \ddots & \ddots & \ddots \\ 0 & 1 & \ddots \\ 0 & 2 & \dots \end{array}$	i ö	1 0
Serial Datc. Number. 1900. 1789 May 8 1816 May 15 1850 May 21 1889 May 28 1922 June 4 1955 June 11 1992 June 18 2027 June 25	(Drain .00001 + ? 0 1 0 1	TABLE I nage Canal, Ke .0001 + - ? 0 1 0 1 0 1 0 1 0 1 0 2		.01 c.c. + - ? 1 0 0 1 0 1 1 0 1 0 1 0 0 0 1	.1 c.c. + - ?
		TABLE I-			
Serial Date, Number, 1900. 1677 Apr. 17 1719 Apr. 24 1755 May 2 1818 May 15 1818 May 15 1858 May 22 1896 May 29 1995 June 5 1961 June 12 2002 June 19 2034 June 26	00001 + - ? 0 1 0 1	rainage Canal, .0001 + - ? 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1	.001 + - ? 0 1 1 1 1 0 2 1 0 1 0 1 0 1 1 0 2 0 0 1 1 0 0 1 0 0 1	.01 c.c. + - ? 1 0 0 1 1 0 0 0 1 0 1 1 0 1 0 1 0	1 c.c. + - ?
		TABLE I	ı.		
418 Aug. 8		splaines River,	Lockport.) .01 c.c. + - ? 0 3	$\begin{array}{c} .1 \text{ c.c.} \\ + ? \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 0 \\ 0 & 1 \\ 0 & 2 \\ \dots \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

TABLE III.

(Kankakee	River	Wilmington.)	
(ALUMINUL CC	TOTA CT.	ALTERNATIVE COMP.	,

Serial Date,	.1 c.c.	1 c.c.
Number, 1899.	+ - ?	+ - :
415 Aug. 7		1 0
496 Aug. 22		
528 Aug. 29		1 0
875 Oct. 30.	V 2 11	1 0
913 Nov. 6		0 1
981 Nov. 20		1 0

TABLE IV.

(Illinois River, Morris.)

Serial Date, Number, 1899.	.0001 e.c. + - ?	.001 c.c. + - ?	.01 c.c. + - ?	.1 c.c + - ?	1 e.c.
917 Nov. 7 962 Nov. 15 985 Nov. 21 1014 Nov. 28	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2 0 2 0 3 0 1 3 0 4 0		
1900 1196 Jan. 7. 1209 Jan. 10. 1248 Jan. 19. 1296 Jan. 27. 1307 Jan. 30. 1338 Feb. 6. 1373 Feb. 13. 1400 Feb. 20. 1461 Mar. 8. 1500 Mar. 15. 1526 Mar. 20. 1571 Mar. 29. 1602 Apr. 4. 1640 Apr. 10. 1680 Apr. 17. 1715 Apr. 23. 1758 May 2. 1784 May 8. 1822 May 16. 1854 May 22. 1893 May 29.		1 0 1 1 0 0 0 1 0 0 0 1 0 0 0 1 0 0 0 2 0 0 0 2 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0 1 0 0 0 0	3 0 1 1 0 2 0 2 2 0 4 1 3 1 1 1 1 1 1 1 1 2 0 0 1 1 0 1 0 1 0 1	2 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 1 2 0 1 0 1 0 1 0 1 0 1 1 2 0 1 1 2 0 1 1 2 0 1 1 1 0 1 0 1 1 2 0 1 1 2 0	
1928 June 5 1962 June 12 1996 June 19 2037 June 27		• • • •		2 0 2 0 2 0 2 0	0 1 1 0 1 0 1 0

TABLE V.

(Fox River, Ottawa.)

Seria				1 c.	.с.	1	C.C.		5	е.е	
Numb	er. 189	19.	+	-	- ?		-	?	+		0
384	Aug.	1	0	1		0	1 .				
420	Aug.	8	0	1		0	1 .				
500	Aug.		0	1							
529	Aug.	29	0	1		0	0	1			
881	Oct.	31	0	1		1	0.				
918	Nov.	7	0	1	٠.	0	1 .				
953	Nov.	14	0	1		0	1 .				
986	Nov.		0	1		0	1 .				
1015		28	0	1		0	1 .				
	19	000									
1282	Jan.	26	0	2	1	1		•			
1528	Mar.	21	0	3		0	3 .				
156 8	Mar.	28				0	2	1	2	0	
16 03	Apr.	4	1	0	* *	2	0.		1	0	
1642	Apr.	11	• •			2			1	0	
1681	Apr.	18	0	1		0	2 .	•	1	0	
1722	Apr.	25	0	1		0	1	1	()	0	1
1759	May	2	0	1		0	2 .		1	0	• •
1790	May	9	1	0		2		•	1	0	
1823	May	16	0	1	• •	0	2 .		1	0	
1859	May	29	0	1		0		•	0	1	
1930	June	6	0	1	• •	0	2 .	•	0	0	1
1963	June	12	0	1	• •	0	1	1	1	0	
1997	June	19	()	1	• •	0	2 .	•	1	0	
2035	June	26	0	1		1	1 .	•	1	0	

TABLE VI.

(Illinois River, Ottawa.)

Serial Date. Number, 1899.	.01 c.c. + - ?	.1 c.c. + - ?	1 c.c. + -1;?	15 c.c. + - ?
385 Aug. 1. 421 Aug. 8. 501 Aug. 22. 530 Aug. 29. 830 Oct. 31. 919 Nov. 7. 934 Nov. 14. 987 Nov. 21. 1016 Nov. 23. 1076 Dec. 13. 1130 Dec. 22. 1153 Dec. 29.		0 1 0 1 0 1 1 0 2 1 2 1 2 2 1 0 3 0 3 1 3 1 1 0	0 1 1 0 0 0 1 1 0 1 2 0 2 0 0 0 2 2 0 1 0 0 0 2	
1174 Jan. 5. 1221 Jan. 12. 1250 Jan. 19. 1283 Jan. 26. 1240 Feb. 6. 1402 Feb. 21. 1455 Mar. 6. 1490 Mar. 13. 1529 Mar. 21. 1569 Mar. 28. 1604 Apr. 4. 1643 Apr. 11. 1682 Apr. 18. 1723 Apr. 25. 1760 May 2. 1791 May 9. 1824 May 16. 1860 May 23. 1931 June 6. 1964 June 12. 1998 June 19. 2036 June 26.	1 0 0 1 0 2 0 0 2 0 0 2 0 1 1 0 2 0 1 1 0 2 0 1 1 0 1 0 1 0 1 1 0 0 0 1 1 0 1 0 1 1 1 0 0 1 1 1 0 0 1	3 0 1 2 0 3 0 1 4 0 1 3 0 2 1 1 0 2 1 1 0 2 1 0 1 0	2 0 1 0 2 0 1 0 2 0 1 0 2 1 0 19 1 2 0 1 1 0 1 0 0 2 1 1 0 0 0 2 0 0 0 2 0 0 0 2 0 0 0 2 1 1 1 1	
470.4	TABLE V			
Serial Date, Number. 1899. 388 Aug. 2. 423 Aug. 9. 502 Aug. 23. 533 Aug. 30. 882 Nov. 1. 920 Nov. 8. 959 Nov. 15. 988 Nov. 22. 1017 Nov. 28.			$\begin{array}{c} .1 \ c.c. \\ + \ - \ \end{array}$ $\begin{array}{c} .0 \ \ 5 \ \ldots$ $\begin{array}{c} .0 \ \ 1 \ \ldots$ $0 \ \ 1 \ \ldots$ $1 \ \ 0 \ \ldots$ $\vdots \ \ \vdots$ $0 \ \ 1 \ \ldots$ $0 \ \ 1 \ \ldots$ $0 \ \ 1 \ \ldots$	1 c.c. + - ? 0 1 0 0 1 0 0 1 0 1 1 0 1 1 0 1 0
	TABLE VI	III.		
Serial Date, Number. 1899. 389 Aug. 2. 424 Aug. 9. 503 Aug. 23. 534 Aug. 30. 883 Nov. 1. 960 Nov. 15. 989 Nov. 22. 1018 Nov. 28.			.1 c.c. + - ? 0 0 1 0 1 0 1 0 1 0 1 0 1	1 c.c. + - ? 0 0 1 0 0 1 0 0 1 1 0 1 0
, .	TABLE IX			
Serial Date, Number. 1899.	llinols River,	Henry.) .01 c.c. + - ?	.1 c.c. + - ?	1 c.c. + - ?
107 June 7. 134 June 14. 173 June 21. 212 June 28. 583 Sept. 7. 616 Sept. 13. 652 Sept. 20. 690 Sept. 27. 1116 Dec. 20. 1900 1170 Jan. 4. 1207 Jan. 10.		0 1	0 1 0 1 0 1 1 0 1 1 1 3	0 1 0 1 0 1 0 1 0 1 1 0 1 0 1 0 1 2 2 1 3 1
LEAST DULL. Transferred to the second	195			J 1

TABLE X.

		TABLE :	Χ.		
	(IIII)	nols River, A	veryville.)		
Serie	il Date, er. 1899.		.01 c.c.	.1 c.c.	1 c.c.
119	June 9		1		0 1
136	June 12				0 1
174 213	June 21		• • • • •	0 1	$\begin{smallmatrix}0&1&\ldots\\0&1&\ldots\end{smallmatrix}$
387	Aug. 2.				1 0
580 617	Sept. 6			$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
663	Sept. 13 Sept. 20			0 1	0 1
691	Sept. 27	• • • • • • • • • • • • • • • • • • • •		0 1	0 1
1275	1900 Jan. 24		0 4	0 2	
1312	Jan. 31			$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
1344 1380	Feb. 7			$\begin{array}{cccc} 0 & 2 & \dots \\ 0 & 0 & 2 \end{array}$	$\begin{smallmatrix}0&1&3\\3&0&1\end{smallmatrix}$
1406	Feb. 21			0 1 1	4 0
1437 1459	Feb. 28			$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2 2
1498	Mar. 14			1 0 1	0 2 1
1533 1570	Mar. 21		** **	$\begin{array}{cccc} 0 & 1 & 2 \\ 0 & 2 & 1 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
1608	Apr. 4.			0 1	1 0 1
1647	Apr. 11			0 0 1	$\begin{smallmatrix}0&0&2\\0&2&\ldots\end{smallmatrix}$
1687 1727	Apr. 18			$\begin{smallmatrix}0&2&\dots\\0&1&\dots\end{smallmatrix}$	$\begin{smallmatrix}0&2&\dots\\0&1&1\end{smallmatrix}$
1765	May 2			0 1	0 1 1
1795 1832	May 9			$\begin{smallmatrix}0&1&\dots\\0&0&1\end{smallmatrix}$	$\begin{array}{cccc} 0 & 1 & 1 \\ 1 & 0 & 1 \end{array}$
1864	May 23			0 1	0 0 2
1901 1935	May 29 June 6			$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
1969	June 13			0 î	0 2
2006 2041	June 20			0 1	2 0
2011	Julie 21	• • • • • • • • • • • • • • • • •	• • • • •	0 1	0
	(Hliv	TABLE Y			
Seria		.0001 e.e.	.01 e.e.	.1 e.e.	1 c.c.
	per. 1899.	+ - ?	- · · ·	+ - ?	1 ?
$\frac{120}{136}$	June 9 June 14			$\begin{smallmatrix}0&1&\dots\\0&1&\dots\end{smallmatrix}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
177	June 21			2	ĭ 0
214 581				1 0	
	June 28			1 0	î ŏ
618	June 28. Sept. 6. Sept. 13.		$\begin{array}{cccc} \vdots & \vdots & \vdots \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{array}$	$\begin{array}{cccc} 1 & 0 & \cdots \\ 0 & 0 & 1 \end{array}$	
	June 28. Sept. 6. Sept. 13. Sept. 27.		o o i	1 0	
618 692 1345	June 28. Sept. 6. Sept. 13. Sept. 27. 1300 Feb. 7.		0 0 1 0 0 1 0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 0
618 692 1345 1381	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14.		0 0 1 0 0 1 0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 0 0 0 2 2 0
618 692 1345 1381 1411 1460	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. May. 7.		0 0 1 0 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2	1 0
618 692 1345 1381 1411 1460 1534	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14 Feb. 23 May. 7. May. 22.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0	1 0
618 692 1345 1381 1411 1460	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. May. 7.		0 0 1 0 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 0 0 0 1	1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648	June 28. Sept. 6. Sept. 13. Sept. 27. 1300 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 1 0 2	1 0
618 692 1345 1381 1411 1460 1534 1581 1610	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 2	1 0 0 0 2 2 0 2 0 1 1 1 0 1 0 2 0 1 1 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 0 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 0 1
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 2 0 2 0 1 1 0 1	1 0 0 0 2 2 0 1 1 1 0 0 0 1 1 1 1 0 0 0 1 1 1
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23.	0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 1 0 1 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 0 1
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mav. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 18. Apr. 25. May 3. May 9. May 9. May 16. May 23. May 31.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 2 1 0 1 2 0 1 0 1 2 0 1 0 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 1 0 0 1 1 1 0 0 1 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 1 1 1 0 0 0 1 1 1 1 0 0 0 1 1 1
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 0 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 0 1 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 0 1 1 0 1 1 0 1 1 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21.		0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 0 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 0 1 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21.	0 1 0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 1 0 1 1 0 1 1 1 1 1 1 2	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 0 1 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21. June 28.	0 1 0 1 0 1 0 1	0 0 1 0 1 0 1 0 1 0 2 0 0 2 0 0 2 0 0 0 2 0 0 0 0	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 0 1 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21. June 28.	0 1 0 1 0 1 0 1	0 0 1 0 1 0 1 0 1 0 2 0 0 2 0 0 2 0 0 0 2 0 0 0 0	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 1 0 0 1 1 1 0 1 1 0 1 0 0 1 1 0 0 1 0 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 21. June 28.	0 1 0 1 0 1 0 1	0 0 1 0 1 0 1 0 1 0 2 0 0 2 0 0 2 0 0 0 2 0 0 0 0	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 2 2 0 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 2 0 1 1 1 0 0 0 1 1 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1829 1938 1973 2008 2042	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 9. May 16. May 23. May 31. June 7. June 14. June 28.	0 1 0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 1 1 1 1 2 1 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 0 0 1 1 0 1 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042	June 28. Sept. 6. Sept. 13. Sept. 27. 1300 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 9. May 16. June 7. June 14. June 21. June 28. June 7. June 7. June 7. June 14. June 7. June 7. June 7. June 7. June 7. June 14.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 0 1 0 1 1 0 1 0 1 1 1 1 2 1 3 1 4 1 5 1 6 1 7 1 8 1 9 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042 Seria Numb 115 138 178 221	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 28. June 7. June 29.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 0 2	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 2 0 0 1 0 2 0 2 0 0 1 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 0 0 1 1 0 1 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042 Seria Numb	June 28. Sept. 6. Sept. 13. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 28. June 7. June 28. June 7. June 28. (11 11 Date, er. 1899. June 21. June 29. Sept. 7.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 1 0 1 1 1 1 2 1 3 1 4 1 5 1 6 1 7 1 8 1 9 1 1 1 1 1 1 1 1 1 2 1 1 1 2 1 3 1 4 1 5 1 6 1 7 1 7 1 8 1 8 1 9 1 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 2 0 0 1 0 2 0 0 2 0 0 1 1 0	1 0 0 0 2 2 0 1 1 2 0 1 1 0 0 1 1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1902 1938 1973 2008 2042 Seria Numb 115 138 178 221	June 28. Sept. 6. Sept. 13. Sept. 27. 1900 Feb. 7. Feb. 14. Feb. 23. Mar. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 31. June 7. June 14. June 28. June 7. June 29.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 0 1 0 1 1 0 2 1 1 1 1 1 2 1 3 1 4 1 5 1 6 1 7 1 8 1 9 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 2 0 0 2 0 2 0 0 1 2 0 1 0	1 0
618 692 1345 1381 1411 1460 1534 1581 1610 1648 1689 1728 1767 1796 1829 1866 1829 1938 1973 2008 2042 Seria Numb 115 138 178 221 589 625	June 28. Sept. 6. Sept. 13. Sept. 27. 1300 Feb. 7. Feb. 14. Feb. 23. Mav. 7. Mar. 22. Mar. 30. Apr. 5. Apr. 11. Apr. 18. Apr. 25. May 3. May 9. May 16. May 23. May 91. June 7. June 14. June 21. June 21. June 29. Sept. 7. Sept. 14.	0 1 0 1 0 1	0 0 1 0 0 1 0 1 0 2 0 2 0 2 0 2 0 2 0 2 0 2 0 1 0 1 0 1 0 1 1 0 1 0 1 0 1 1 0 1 0 1 1 1 0 1 1 1 2 1 3 1 4 1 5 1 6 1 7 1 8 1 9 1 1 1 1	1 0 0 0 1 0 0 1 1 3 1 2 1 1 3 1 1 2 0 1 0 2 0 2 0 0 1 0 2 1 0 1 2 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 0 1	1 0 0 0 2 2 0 1 1 2 0 1 1 0 0 1 1 0

TABLE XIII.

(Sangamon River, Chandierville.)

Scrial Date, Number, 1899.	.01 c.c. + - ?	1 c.c.	1 c.c.
116 June 8. 182 June 22. 626 Sept. 14. 660 Sept. 21. 715 Sept. 29.		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0 1 1 0 0 0 1 1 0 1 0
1900 1276 Jan. 25 1315 Feb. 1 1355 Feb. 3 1384 Feb. 15 1438 Mar. 2 1470 Mar. 8 1508 Mar. 15 1544 Mar. 22 1572 Mar. 29 1613 Apr. 5 1652 Apr. 12 1692 Apr. 19 1731 Apr. 26 1770 May 3 1805 May 10 1839 May 17 1875 May 24 1911 May 31 1945 June 7 1981 June 14 2011 June 21 2051 June 28		1 3 0 4 2 0 2 1 1 0 1 3 3 1 1 1 1 1 3 0 0 2 1 1 0 0 2 1 0 2 0 1 0 2 0 1 0 2 0 1 0 0 2 1 0 0 1 1	4 0 0 4 1 0 1 3 1 2 0 3 3 0 0 1 1 1 1 2 0 2 0 1 0 1 0 1 0 1 0 1 0

TABLE XIV.

(Illinois River, Grafton.)

Serial Numbe	Date, er. 1899.	.01 c.c. + - ?	.1 c.c. + - ?	+ 1 c.c.	5 c.c. + - ?
117 184	June 8June 22			$\begin{smallmatrix}0&1&\dots\\0&1&\dots\end{smallmatrix}$	
222	June 30		** ** **	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	• • • • • •
591	Sept. 7		0 i	0 1	
628	Sept. 14		1 0	$\tilde{1}$ $\tilde{0}$	
662	Sept. 21		$1 0 \dots$	1 0	
701	Sept. 29		0 1	1 0	
996	Nov. 23	1 1	2 0	2 0	
1025	Dee. 1	0 1	1 0	1 0	
1057	Dec. 6	0 1	0 1	1 0	
1098	Dee. 15		0 1	0 1 1	• • • • • • • • • • • • • • • • • • • •
1127	Dec. 22	• • • • • •	$\begin{smallmatrix}0&0&1\\0&1&\ldots\end{smallmatrix}$	$\begin{array}{cccc} 1 & 0 & 1 \\ 0 & 0 & 2 \end{array}$	
1156	Dec. 29		0 1	0 0 2	
1187	Jan. 5		0 1	0 2	
1217	Jan. 10		0 1	$\begin{smallmatrix}0&&1&&1\\0&&2&&2\end{smallmatrix}$	
1261	Jan. 19		0 4	0 2 2	
1287	Jan. 26		1 1	2 1 1	
1317 1356	Feb. 1	0 2	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1386	Feb. 8	•• •• ••	$\begin{smallmatrix}0&1&1\\1&0&1\end{smallmatrix}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1408	Feb. 22	• • • • • •	0 2	3 0 1	
1484	Mar. 9.		, i i	4 0	•• •• ••
1509	Mar. 16		2 .0	4 0	
1536	Mar. 22		0 3	0 ĭ ż	
1579	Mar. 29		0 3	$2 \ 0 \ 1$	
1615	Apr. 5		2 0	1 0 1	
1654	Apr. 12		0 1	1 0 1	
1693	Apr. 19		0 1	1 0 1	
1732	Apr. 26	•• •• ••	0 1	1 1	
1771	May 3		1 0	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	•• •• ••
1806 1841	May 10	•• •• ••	1 0		
1841	May 24		1 0	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1912	May 31		0 1	0 2	
1946	June 8		0 1	0 0 i	0 0 i
1983	June 15		0 1	1 0	1 0
2019	June 22		0 1	2 0	$\bar{0}$ $\tilde{0}$ $\tilde{1}$
2053	June 29		1 0	2 0	1 0

TABLE XV.

(Mississippi River, Grafton.)

Serial Dat Number, 189		.01 c.c.	.1 c.c.	1 e.c.	5 c.c.
	8		0 1	0 1	
	30		0 1	1 0	
629 Sept. 663 Sept.	14 21		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{smallmatrix}0&0&1\\1&0&\ldots\end{smallmatrix}$	
702 Sept. 1026 Dec.	29	0 1	$\begin{array}{ccccc} 0 & 1 & \dots \\ 0 & 1 & \dots \end{array}$	1 0	
1058 Dcc. 1099 Dec. 1128 Dec.	6 15 22.	0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1157 Dec. 1900	29		ŏ i	ō ž	
1188 Jan. 1218 Jan.	5		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 1	
1262 Jan. 1286 Jan. 1318 Feb.	19		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
1357 Feb. 1387 Feb.	8 15		1 1	1 1 2 3 1	
1409 Feb. 1435 Mar.	22 9		0 2	$\begin{array}{cccc} 2 & 0 & 2 \\ 2 & 1 & \ddots \end{array}$	
1510 Mar. 1537 Mar. 1580 Mar.	22		$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
1616 Apr. 1655 Apr.	5		$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	0 1 1 1 1	
1694 Apr. 1733 Apr.	19 26		0 1 0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1772 May 1807 May	3		0 1 1 0	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1878 May 1913 May 1947 June	24		1 0 1 0 0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 0
1984 June 2019 June	14 22		0 1 0 1	$\begin{smallmatrix}0&0&\bar{1}\\0&2&\ldots\end{smallmatrix}$	1 0
2054 June	29		0 1	2 0	1 0

TABLE XVI.

(Mississlppi River, Cross-Section at Alton.)

	(Mississippi Mivet, Cross-acction at Anto					
Scrial Number.		. 1	e.e.	1	e.e.	
Munder.	East Bank—	-		†	_	
284 J 319 J 356 J 736 C 773 C	East Bank	0 0 0 0 0 0	1 1 1 1 1	0 0 0 0 0 0 1	1 0 1 1 1 0	1
285 J 320 J 357 J 737 C 774 C	East Center— [uly 6	0 0 0 1 0 0	1 0 1 1 0 1 1 1	0 0 0 1 0 0	1 0 1 0 1 0	·i ·· ·· 1
286 J 321 J 358 J 738 C 775 C	Center— [uly 6. [uly 13. [uly 20. [uly 27. [ult 5. [ult 5. [ult 26. [ult 12. [ult 26. [ult 26.	0 0 0 0 0 0	1 1 1 1 1 0	0 1 0 0 0 0	1 0 1 1 1 1 0	:: :: :: :: :: :: :: :: :: :: :: :: ::
287 J 322 J 359 J 739 C 776 C	West Center— July 6. July 13. July 20. July 27. Oct. 5. Oct. 12. Oct. 26.	0 1 0 0 0 0	1 0 1 1 1 0	1 0 0 0 0 0	0 1 1 1 1 0 0	· · · · · · · · · · · · · · · · · · ·
288 J 323 J 360 J	West Bank July 6. July 13. July 20. July 27. July 5.	0 0 0 0	1 1 1 1	0 0 1 1 0	1 1 0 5 1	• •

	TABLE X	VII.		
	(Missouri River,	West Aiton.	.)	
	al Date, er. 1899.	.01 c.c.	.1 c.c.	1 c.c
376 671 714 749 789 869	July 28. Sept. 23. Sept. 29. Oct. 6. Oct. 13. Oct. 27.		0 1 0 1 0 0 1 0 1 0 1	0 1 0 0 1 0 0 1 1 0 0 1 0 1
1189 1235 1263 1295 1320 1358 1388 1410 1439 1471 1499 1556 1594 1609 1650 1721 1766 1793 1831 1865 1910 1937 1970 2007 2050	1900 Jan. 5. Jan. 10. Jan. 19. Jan. 26. Feb. 2. Feb. 8. Feb. 15. Feb. 22. Mar. 2. Mar. 8. Mar. 14. Mar. 23. Mar. 30. Apr. 4. Apr. 11. Apr. 18. Apr. 18. Apr. 25. May 3. May 10. May 10. May 23. May 31. June 7. June 13. June 20. June 28.		2 0 2 0 1 0 2 0	0 4 1 2 1 3 1 0 4 2 2 2 0 3 1 0 4 2 1 1 2 1 2 1 1 2 2 0 2 0 2 0 2 0 1 1 0 1 0 2 0 2 0 1 0 1 0 2 0 1 0 1 0 2 0 2 0 2 0 1 0 2 0 1 0 2 0 1 0 2 0 2 0 2 0 1 0 2 0 2 0 2 0 3 0 3 0 3 0 4 0 2 0 3 0
	TABLE XV			
Serial Numb	(Mississippi River, Intake, Tower, St. Date, er. 1899. + - ?	. Louis W	aterworks. Mitchell 1 c.c. + - ?	.) + 5 c.c. + - ?
295 330 373 743 859	July 14. July 21. July 28. Oct. 6. Oct. 27.	1 0 0 1 1 0 0 1 0 1	1 0	
1299 1361 1426 1474 1547 1585	1900 Jan. 27 Feb. 9. Feb. 24 Mar. 9. Mar. 23. Mar. 30.	1 1 1 0 1 1 1 0 0 1 3 0 0 2	$\begin{array}{cccc} 1 & 0 & 3 \\ 1 & 0 & 1 \end{array}$	·· ·· ·· ·· ·· ·· ·· ·· ·· ·· ·· ·· ··

Number. 1899.	+ - :	+ - :	+ - ?	+ - ?
295 July 14		1 0	1 0	
330 July 21		0 1	1 0	
373 July 28		1 0	1 0	
743 Oct. 6		0 1	0 0 1	•• •• ••
859 Oct. 27 1900	• • • • • • • • • • • • • • • • • • • •	0 1	0 1	
1299 Jan. 27		1 1	1 1 2	
1361 Feb. 9		î d'i	4 0	
1426 Feb. 24		1 1	1 0 3	
1474 Mar. 9		0 0 1	1 0 1	
1547 Mar. 23		3 0	2 0 1	
1585 Mar. 30 1624 Apr. 6		0 2	1 1	1 0
1624 Apr. 6 1658 Apr. 13	• • • • • •	0 1	$\frac{1}{2}$ $\stackrel{1}{0}$ \dots	
1702 Apr. 20		ĭ i	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	
1810 May 11		$\hat{1}$ $\hat{0}$	2 0	0 0 1
1845 May 18		0 0 1	0 0 2	0 0 1
1885 May 25		1 0 6	0 0 2	
1917 June 1	• • • • • •	1 0	2 0	
1950 June 8	•• •• ••	0 0 1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
1987 June 15	1 0	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{smallmatrix}2&0&\ldots\\2&0&\ldots\end{smallmatrix}$	
2057 June 29				

TABLE XIX.

(St. Louis Tap Water.)

		(1000 2000 200)		/		
Seria Numb	al Dat er. 189		+ 1	e.e. ?	+ 1 e.e. ;	5 c.c. + - ?
265	July	7	0	1	1 0	
298 333	July July	15 22	0	1	1 0	
365	July	28	Ŏ	1	$\bar{0}$ 0 1	
781	Oct. 1900	13	0	1	0 1	
1289	Jan.	26	1	1	1 2 1	
1347			1	$\frac{1}{2}$	0 3 1	
1364 1376	Feb. Feb.	10 13	0	2	0 3 1	
1383	Feb.	15	0	4	0 4	1 0
1395	Feb.	17	0	$\begin{array}{ccc} 2 & \dots \\ 2 & \dots \end{array}$	$\begin{smallmatrix}0&3&1\\0&4&\ldots\end{smallmatrix}$	1 1
1413	Feb.	23	U	٠. د	0 1	1 1

1448 Mar. 5. 1476 Mar. 9. 1514 Mar. 17. 1549 Mar. 23. 1587 Mar. 30. 1626 Apr. 6. 1660 Apr. 13. 1704 Apr. 27. 1778 May 4. 1812 May 11. 1847 May 18. 1879 May 25. 1914 June 1. 1952 June 8. 1989 June 23. 2060 June 29.			0 0 0 0 1 0 0 0 0 0 0 0 0	2 1 3 2 1		0 0 2 2 1 0 0 1 0 1 2 0 0 1 1 1 1 1	3 1 1 0 0 2 2 1 1 1 1 1 1 1 0 0 1			
I. PRINCIPAL STATIONS	s o	N TH	Œ I	LLIN	OIS	RIVI	ER.			
).	0000)1 c.c.	.000)1 c.c.	.001	l c.c.	.01	c.c.	.1 C	e.c.
Collecting Stations.	water exam'd.	No. of days B. coli found.	No. of days water exam'd.	No. of days B. eoli found.	No. of days water exam'd.	No. of days B. coli found.	No. of days water exam'd.	No. of days B. coli found.	No. of days water exam'd.	No. of days B. coli found.
Illinois and Michigan Canal, Lockport Illinois River, Morris Illinois River, Ottawa Illinois River, Averyville Illinois River, Wesley City Illinois River, Grafton	• • •	7	32 3	28 1	11 20 	8 11 	4 30 22 1 22 4	4 20 6 0 3 1	2 23 34 27 26 35	2 20 19 4 13 13
	I.									
Illinois River at Averyville and Grafton Mississippi (Grafton) and	n, C Mis	Compa sourl	red (W	with est A	Tri (Alton	butar Rlv	ies a	nd	with	the
			.01	c.c.	.1	c.c.	1	c.c.	5	c.c.
Collecting Stations.			No. of days water exam'd.	No. of days B. coli found.	No. of days water exam'd.	No. of days B. coli found.	No. of days water exam'd.	No. of days B. coll found.	No. of days water exam'd.	No. of days B. eoli found.
Illinols River, Averyville Illinols River, Grafton Mississippi River, Grafton Desplaines River Kankakee River Fox River Big Vermillon River Sangamon River Missouri River			13	0 1 0 4 3	27 35 34 8 6 22 5 25 32	4 13 10 1 3 2 1 14 13	31 38 35 5 5 23 9 27 31	13 26 23 2 4 6 3 21 21	13 	2 3 10
	11.									
Total, Illinois River (Averyville and Grafte Total, tributaries of Illinois River				• •	62 66 66	17 21 23	69 69 66	39 36 44	• •	



STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, BRIDGEPORT.

Report of Arthur W. Palmer, T. J. Burkell. University of Illinois.

No. of Bac.	per Cubic Centi- meter.	1.035.500 2.355.000 4.500.000 4.770.000 1.070.000 1.425.000 1.250.000 1.500.000 1.900.000 3.920.000 3.920.000 3.920.000 3.920.000 3.920.000 3.920.000	1,990,000
	Presenc	:::+:::+!++++:+++++	
ture tine(Tempera	:: : : : : : : : : : : : : : : : : : :	φ :
ture r, C.	srequest of Wate		6.5
1.13	dgiəII əte7/	: 4410 f	; ;
MILITARES.		######################################	œ :
NITRO- GEN	Nitrites.	100 100	
pended.		0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	တ လ :
ORGANIC NITROGEN	Dis- solved.		36 : :
OI	Total.		7.72
	i p,pud	85 95 95 85 85 85 85 85 85 85 85 85 85 85 85 85	—÷ 5::
N AS 1A.	pioi p, Alos	<u> </u>	<u> </u>
NITROGEN AS AMMONIA.	Albuminoid A line in	++8884+889±9=9+69298+896 ++8884+89±9=9+6228+886 804+691-696000000000000000000000000000000000	≎) ·
Nr	.sinom	A)	ro :
	Уветет. Free ли-	m & c & si c & si - S & + c & + si c & = 4 & o &	න :
ONYGEN	solved. By Suspd	<u></u>	<u> </u>
ONYGEN	Total.		∞ : ∞ :
Chlorine.		28. 28. 29. 29. 29. 29. 29. 29. 29. 29. 29. 29	%
	on Igni- tion.		<u>= :</u> : :
3	. n mud	± ± ∞ 0 ∞ ± 0000000 0000000000000000000	7
TON.		118.4.58.4.60.4.118.4.58.4.60.4.118.4.58.4.60.4.118.4.58.4.60.4.118.8.4.119.8.	0e 2.
ORAT	lo ss o	25.25.26.26.27.25.26.25.25.25.25.25.25.25.25.25.25.25.25.25.	GI 9.
Evar		2-1-4-2-2-4-1-4-4-4-4-4-4-4-4-4-4-4-4-4-	+ :
RESIDUE ON EVAPORATION.	-sus		£ .
SIDU	Dis-	771 4 615 6 610 8 510 6 610 8 510 6 610 8 510 6 610 8 510 6 610 4 511 6 610 4 511 6 610 4 511 7 610 138 7 610 13	541.6
RE	Total.	771.4.615.6 (201.6.1000. (510.8.510. (510.8.510. (510.8.510. (510.8.510. (510.4.511. (510.	621.
*.1	юрО	Cas: 111111111111111111111111111111111111	
-	<u>.</u>		-
NCE.	Color.	NAM JANAMANAMANAMANAMANAMANAMANAMANAMANAMANA	N. F.4
APPEARANCE	nt.		: ::
APP	Sedi- ment	Cons'd Much Cons'd Much Cons'd Wuch Wuch Wuch Much Cons'd Cons'd Cons'd	Much
	Turb'y.	% 1	* :
: 0F	1899 Exam nation		:
DATEOF	1899 1899 Offee- Exami- tion. nation.	April 27 April May 11 May 19 M	£ :
	10.) (10.)		: :
No.	Serial	66.00	657

Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown. Turbidity-*Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LOCKPORT.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

No. of	rac. per Cubic Centi- meter.	5.53.000 6.175.000 6.175.000 6.175.000 4.115.000 6.080.000 6.080.000 1.110.000 1.27.500 1.110.000 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500 1.27.500	490.000 875.000
oli.	Presence Abs, of C	<u> </u>	· · :
	Temperat o'), 'il A lo		→ { ~
	Temperat		5.1-
10	Height Water	<u>∞</u> 1 <u>~</u> 1	: :
NITRO-	Sitratia		근 (1) 임명
NIN	20 HatiX	+0.	5,5,
NIC	Sus- Sus- pended.	<u> </u>	20 00 20 01
ORGANIC	-sid beylos	2255 55 55 55 55 55 55 55 55 55 55 55 55	35.E
	1070[]	16 छ में 16 16 16 16 में हो में में में 16 16 में में में में में में में में 16 16 में छ में 16	51-
AS	m p,pud		5.5 - 3.5 -
NITROGEN AS	Meaning Pivios -sid		5. 5. 8. 2.
NITERO	1'10'T	क्षेत्रक को को को को को को 🕂 को	# #
	Free Am-	0.00 € 20 € 20 € 20 € 20 € 20 € 20 € 20	1.02 1.02 1.03 1.03 1.03 1.03 1.03 1.03 1.03 1.03
EN	Solved. By Suspd Matter.		6.52 5.52 5.00 5.00
ONYGEN	Total. By Dis-	I man in the contraction in the	= 8 <u>1</u>
·91	Chlorin	10 10 11 01 01 01 10 11 01 0	7.5 36
	on Igni- tion.		29
5		<u> </u>	- c)
T10N.	D MOS	6884186841868418484848484848484848484848	= 1 = 1 = 1 = 1
PORA	Toll Loss on		5.65
RESIDUE ON EVAPORATION	bended	68 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	20 E
LE O	solved.		e +
RESID	-ei(I	6.33 6.97 4.10 6.33 6.33 6.33 6.33 6.33 6.33 6.33 6.3	633.6 562.4
	Total.	-	888
	торО	N 00:::::::::::::::::::::::::::::::::::	: :
52	Color.	2277472747277777777 2777472777777777777	5.5
APPEABANCE			2.2
APPE	Sedi- ment	Much ('ons'd Much ('ons'd Much Much Much ('ons'd Much Much ('ons'd Much Much ('ons'd Much Muc	Much
	· v.qand,	* * * * * * * * * * * * * * * * * * * *	H-
Sat Co.	1599 1599 ('Ollee-Examition.		ति के । : :
Dateor	0 1. ng		<u> </u>
	Collection.		: :
.o.V	Serial		67.

Trubinity * Decided, § Very Decided, † Distinct, † Very Slight, | Slight.

Color on Lonition G., Gray. B., Brown. DB., Dark Brown. L.R., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. R., Reddish. W., White. Gh., Grayish. Blk., Brack. GB., Gray. Brown.

Color M., Muddy, VM., Very Muddy, T., Turbid, C., Cloudy,

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burnell, University of Illinois.

SOURCE OF WATER-DESPLAINES RIVER, LOCKPORT.

No. of Bac.	per Cubic Centi- meter.	2. 12.200 2. 12.	
e or Coli.	Presence Abs. of	11111 1111 1111	
.tuə'	Tempera of Air, U		0
1111'e r, (',	Tempera	::::84444888.:3449856798979900000000000000000000000000000000	G
·.I.:	нgiэН этв₩		
TIO-GEN AS	Sitrates	<u>444445554686665555555555566248889</u>	.73
NITHO-	Sitrities.	<u> </u>	.003
	bended.	5444482684888888888888888888888888888888	.192
ORGANIC NITROGEN	Dis- solved.		
ON	Total.	22225222528228255555555555555555555555	.776
- So	Pud's Sus-sus	200	
EN A	Nichtinoid I'10'T Dis- bivoid bivios	<u>*************************************</u>	.256
NITROGEN AS AMMONIA.	Tot'l	<u></u>	
Z	Free Am- monia.	2:	. 152
N ED.	By Suspd	## ## ## ## ## ## ## ## ## ## ## ## ##	0.0
OXYGEN Jonsumed	By Dis-	n+m+n+n+n+n+n+n+n+n+n+n+n+n+n+n+n+n+n+n	σ. ∞
Con	Total.	$\overline{u} : \overline{z} \overline{\omega} \overline{\omega} \overline{\omega} \overline{u} \overline{u} \overline{u} \overline{u} \overline{u} \overline{u} \overline{u} u$	6.∞ 8.0
.en	Chlori	$\frac{\omega - \omega \omega \omega \nu (c_F - \omega \omega + \nu \omega + r \omega + \omega \omega \omega \omega \nu + r - r - \omega \omega \omega \omega \nu (c_F - \omega \omega \omega \omega) (c_F - \omega \omega \omega \omega \nu + r - \omega \omega \omega \omega \nu) (c_F - \omega \omega \omega \omega \nu + r - \omega \omega \omega \omega \nu + r - \omega \omega \omega \omega \nu)}{(c_F - \omega \omega + \nu \omega \omega \nu + r - \omega \omega \omega \omega \nu + r - \omega \omega \omega \omega \nu)}$	5. To
Jolor	on igni- zion.		
	z p,pud	######################################	ος. Ος.
TION	e sid b'vios	<u> </u>	4
RESIDUE ON EVAPORATION	Tot'l	ο οι ορασισισια ο σι-σιο α αρα σ	ક્યું. જ
N Ev	bended.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	· ·
) L.E	solved.		
RESII	-sid	\bigcirc	10.
	T'otal.		25. 25. 25. 25.
•	торО	N 0:1:::::::::::::::::::::::::::::::::::	9
<u> </u>	Color.	######################################	¥.5
RANC	သိ	वंत्रवं वंत्रवंत्रं वंत्र वं वंत्रवंत्रं वं	
APPEARANCE.	Sedi- ment.	Little Consid	*
	Turb's.	+*++*++++++++++++++++++++++++++++++++++	÷
OF	1899 Exami- nation.	May May June 6 4 11 11 11 11 11 11 11 11 11 11 11 11 1	
DATEOF		<u> </u>	92
A	1899 Collec- tion.	May Aug	
.oV	Serial	50107 50	6582

C. Cloudy. RB. Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Blk., Black. GB., Gray-Brown. Colon-M., Muddy. VM., Very Muddy. T., Turbid. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. Rh., Reddish. W., White. Gh., Grayish. TURBIDITY -* Decided. § Very Decided. + Distinct. + Very Slight. Slight.

TABLE 4.

STREAMS ENAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, NORTH OF JACKSON ST., JOLIET.

Report of ARTHUR W. PALMER, T. J. BURRILD.. University of Illinois.

	1	:	- 1	8	28	33	38		8	3	8	8	:	8	8	8	8	3	:	8	9	Ş	9	00	00	00	3	0.)	9	3	1
No. of Bac.	per Cubic Centi- meter.			1.072.5	2,165,000	0.5250 0.5250 0.5500	1.320.00		1.700,000	5,525,000	5,300.0	4.300.000		2.550,000	7.300.000	350.000	216,000	S.C.	:	710.000	0.056.	680.000	830,000	1.380.00	1.980,000	345,000	000.821.1	2.210.0	310,000	370.0KK	
	Presenc	:	•	+-	+-	+-	-	•				+		+	:		+	+	•	+			+	+	-					-	1
dut.	Temper:), 'ii', 'io	:			: - 0			19.	19.	33	5.	10.5	<u>5</u>	- 9	×.	<u></u>		 	S	16.		ຈ≀	-6.	00	0.	Ý.	10.	=	9	15.	
ainte	remperation to	-:	•	:				: : : : : : : : : : : : : : : : : : :	21.	35.	 			: ::	::	연	_ _ _	<u>.</u>	17.	20.		0.	-	Ç	x.	0	23	<u>ان</u>	0.00	10.	
10 11	dgioH ota77			:	:	:		-		:		:		:	:	:	:	:	:	:	:	:	:	:	:		:			:	
TRO- GEN AS	Sitratics.		7.	98.	3) a	_		6:	1	÷;	5	3 5			<u>.</u>			2)			_	_	?				38. I			Ø.	
NITRO- GEN	Nitrites	210.	8	90.	33	100	-	000		9.	0					•	200 —	9 9 -	<u>=</u>	<u></u>	.00e	_	_	400.	9.	(8)	90.	00.	=	.00	
SIC EN.	Sus- bended.	3.4 3.4 3.4		1.65	9.0 0.0	0.0	10 i	÷	31.504	31 35	25.33	3.136			% ?!	x :	% ⊙: ∞:	9:3	5.	55.		. ?	5 22	?? ?!	32. I	1.96	71	=======================================	6.56	3.01	-
ORGANIC NITROGEN	-siG beyles	2.65	1.85	-	¥ 3	- °	19.5	£ -	11.010	1.56	7	F86:	- + + + + + + + + + + + + + + + + + + +	-	- -	£.	- S	G.	192	= ====================================	5	.33	E :	or. ≎1	- :	9	1.16	7.	1.96	3.75	- m-
0.7	Total.	5.13	5.05	1 12	88	7	13	25.	2.52	3.61	3.73	7	~	3 2	35. -	55	3	1- i	- 1	3.56	4.68	200	35 T	ı.	6.11	2 9	1.36	55 33	20.00	05 9	
SA.	p,pud	1.632	1.696	£+G.	92	1.40	2 00	61		•		- 25 - 25 - 25 - 25		•				203	-	_	=	1.56	: ::	1 55.6	差ので	15 CE 51	1 536	21	3.13	 30	
	Albuminoid Dis- Biologia Divid	.768	. X64	.608	%; 3 -	83	61.	Z	.496	198.	+	56.	13.	.64	.701	19%	× = .	£ .	.61	826.	??	19.	1.01	101	33.50	101.	101.	SH:	??	1.76	
NITROGEN AMMONIA	I'to'l'	4.5	2.56	1.153	E ?	2 - 2		55.	21.1	1.11	1.41	18.	9.1	9.1	36:1	1.0.1	<u>z</u>	31		??	33.	-	??	000	? ì	30 30	? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ?	38.00	36	3.01	
7	Бтее Ата. топпа.	00 00	10.8		25 2	15.0	16	5. X.	61	11.4	20.00	음 -	26.4	00 00 10 10	10.3	59.65	9	x	oc.	± 25		ه ا <u>د</u>	12. 6	x.	<u>x.</u>		0) 10	<u>- : : : : : : : : : : : : : : : : : : :</u>	or.	-	
EN IED.	By Suspd	10.3	6.11	1.0	7 - 11 1 - 11				1-	5.0	∞: ??	13.6	23.33	- - - - -	33		G.	0.3	6 1	X	- 1	21	- 0-	0.	G. 657	23 6	0.01	19	13.6	30.3	
OXYGEN CONSUMED,	By Dis-	3 23.	5 18.6		<u> </u>	6 6 6 6 6 0	37	15.	1 15.7	17.3	1 2 2	2 OG X	5.3	20	5 16.8	ار ا ا	1 1 1 1 1	ia : ⊈ ;	7. 2. 2.	1 11 6	= =	G. ≅:	6 17.9	<u> </u>	=:	=	5 10 6	6 11.1	5 11 5	7 19.4	
- 5	Total.	83	200	57	i= 3	070	3 23	5.5	÷	5	26.0		900	- 50.	÷ ;	X.	- To G	56	···	63		5 5 T	5 27	101	.00	13	**	5 5	5 5X	£:	
	irold')	69	35	000	6	8 3		110	Z	3	2	106	199	2	9	=	0	=======================================	£;	3.	=	?? 	£	=======================================	=	=	1003	. S. S.	??	=======================================	
	Jgn. Troni tion	m,			mi a	4:			13.	===	==	~=	GI),	<u>C</u> :	~			<u>-</u> :		=======================================			÷		===		.:	::	==	==	
ON.	in p.pud	80.	6 27. 3	- 6.1	20 c	200	C 2:	8 91 7 89	10.	00 00	30 Oct 30	5.21.1	37 6		9 9	17 6	6 1.6	6 6	1 9 1	x 27 x	er ic X	≎? ⊆: У.	6 16.	:	56	J. 23 €	16.	1.35.6	2) X	X = x	
RATI	Loss on 1'10'l' b'vlos	55.6 34.8	± € €	11.13%	6 45.	200000000000000000000000000000000000000	c 10		50.	6.78	6.54.	91 2 66 8 21.	91.654.	ος ος	9 33	17 6 30.	:Se 27	5. x c c	<u>x</u>	.6.26	??		53 6 37 6 16	1.15	<u> </u>	S 6 32 8 33	œ.	??	15. × 10 × 10	69 6 21.	
EVAP					-	- :	10		23	species.	species	::		-	??	-	32		200	-						13	15	23	33	8	-
RESIDUE ON EVAPORATION.	Sus-					i i			_	×.		_		33				S : 5					17.6	-	216 1			55	=======================================		
SIDUE	-siQ .boyloa	511. 500.1	160.8	464 × 090.2	556 6	0.180	2 2 2 2	537.6	100	551.0	139.6	35	\$ CE	150	£.80	हार ६ महिल	192.× 150.1	5	535	(H)	9	170.1	- XGG	X (E)	585	13:	192	5 (30)	540 1	515.2	
RE	Total.	511.	500 1	× 191	<u>.</u>	616, 581,6 275 C cen c	516.0	587. 5 537	505 5 100	500	100	655 6 168	549.2	2000	(2) (1) ESC	517 6	101.X	515.5	583,6532	11 × 100	5(5)	521	6(46).		いった。一つない	611 6	1 223	5 THE	617.6540	633	
*.1	торО	None	:	• •	7. × E!)			9 9	*,	• •	* 9	0 9	-g -g	:	:	9 4	٠,	•,	•	.,	.,		-2	;	*	1 9	:	6.0	,.	4 4	
6	Color.	10.	23		φ: :-:	T) 18	0 G	210	F. 151	x.	F. 255	五.05	10.	 	13:5	는 1년 1년	-	1.0%	: S		E.55	~: <u>-</u>	F.6	5.5	- T	E.15	18	000	19	F. 5	
RANCI	9		N.		7.5		>		7	N.	7.	. I.	N.	7.	V.N.	7.	7.	7.	7.	7.	7.	-		7.						. 7.	
APPEARANCE.	Sedi- ment.	p.suo.	:	. 9	99	, ,	**		p.suo.	:	9.9	* *	Much	p.suo.	V. M.ch	Much	9.0	p.<00.)	Much	*	9 9	p.suo.	* *	:	C.M.ch	Much	:	:	p. <110.)	Mach	1
	T,q.m,T,	*	.0.	-+ .0	* ·	+ 1	* **		*	*	*	*	-16-	4:	.0.	46			H- 2	*	*	*	#	*	.0	¥:	46	*	*	*	
OF	Collec- Examition,	May 3	: 65	5 June 6	<u>ee</u>	₹ è	in le	: :	:		1112.		<u>(1</u>),	37	.: SE		:	:)/::-	:	97	160 1.	.YOY.	;	· · ·	× 0	Dec. 5		:	:	
DATEOF	Tend Collection.			. 9	27 9	5. 5		=	-	?	50		-	÷ 2 ≥ 2	\$	•	=	P	2 .	16	<u>:3</u>	(F.	9	13	ā	3	,		X	- 2	
	4 5 5	7 May		5153 Jun	:	. :	1 111	0	.,	;	.,	1119	:	;		7.015		:	(Oct.	: :		., 65	SNOV	.,		:) 1)6.C.		:		
.oV	Serlal	716	910	515	5000	300	1000	300	5431	5177	5541	52.00	50000	5660	27.70	27.6	54.50	5933	C121	6,076	613	6183	(2002)	6.27	6223	(1907)	6119	1013	651	0579	

Conon M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Trubibity *Posided, \$Very Decided, * Distruct, \$Very Slight, Slight.

Color M. Muddy, VM. Very Muddy. T. Turbid. C.,

Color on leminon G. Gray, B. Brown, DB. Light Brown, RB. Reddish Brown, BG. Brownish Gray, Bh., Brownish, R., Red.

Rb., Reddish, W., White, Gh., Gray, GB., Gray, Brown.

TABLE

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-DESPLAINES RIVER, SOUTH OF TOWN, JOLIET.

Report of ARTHUR W. PALMER, T. J. BURKILL, University of Illinois.

1,715,000 1.330,000 1.295,000 2.050,000 1,750,000 1,137,500 1.955.000 1.560,000 6.000.000 3.266,660 6.800,000 6.300,000 No. of Bac. per Cubic Centimeter. Presence or Abs. of Coli. 10. Zitrates.

| Height of | Water. | Temperature of Water. | Pemperature of Water, C. | Pemperature of Marer, C. | Pemperature of Marer, C. | Pemperature of Marer, C. | Pemperature of Marer of Ma ळ छुं गंग <u>तृं धार्थ घ</u>र्द्ध घर्छ छुं गंग ह
 72
 1.92
 5.53
 2.49
 3.04

 82
 1.72
 6.69
 1.61
 4.08

 96
 1.66
 2.09
 1.4
 69

 96
 1.12
 3.96
 1.88
 2.08

 72
 1.92
 3.56
 1.36
 3.04

 7.72
 1.92
 5.72
 1.38
 2.8

 7.72
 1.28
 4.6
 1.8
 2.8

 6.72
 1.08
 4.28
 1.36
 2.94

 7.73
 1.04
 4.28
 1.76
 1.24
 3.06

 7.84
 1.8
 1.8
 3.0
 1.4
 1.16
 3.36

 8.8
 3.
 1.76
 1.24
 3.36
 3.6
 3.36

 8.8
 3.
 1.76
 1.84
 4.76
 1.84
 3.92

 5.12
 1.76
 4.76
 1.84
 4.76
 1.84
 3.92

 5.12
 1.76
 4.28
 9.2
 3.36
 3.36
 3.34

 8.8< eng-sus-NITROGEN. ORGANIC paajos -sig Total. Albuminoid Am. p.pud -sng NITROGEN AS -sid b'vlo AMMONIA. ['10'] By Dis-solved. By Saspd Matter. Free Am-monia. OXYGEN CONSUMED. Total. Chlorine. Jolor Igni-tion. on p,pud -sng Loss on 1g'n. RESIDUE ON EVAPORATION. sid b'vlos 1,10,1 ->iq -oived. Total. Gas. .TobO Color. APPEARANCE 5108 ... 29 ... 30 § convd M. 5108 ... 30 § convd M. 5201 ... Sediment. Turby Collec-|Exami-1899 256 266 3 July 5 DATEOF 1899 5228 July 5379 ... 5430 ... 5728 ... 2 5786 Sept. 5830 ... 1 5573 Aug. 5573 Aug. 9.9

T., Turbid. C., Cloudy. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red Blk., Black. GB., Gray-Brown. V.M., Very Muddy. Colon-M., Muddy. Colou on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. Rb., Reddish. W., White. Gh., Grayish. TURBIDITY -* Decided. Nery Decided. + Distinct. + Very Slight. | Slight.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-KANNAKEE RIVER, WILMINGTON

Report of ARTHUR W. PALMER,	T. J. BURRELL,	University of Illinois.
		-

0. of	per Cubic Centi- neter.	8 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
,) Jo sqv	
	Presence	
	ot Water	
9.111.1	reinperarei	
10	।	86858888844589995555555556 8944444444
NITRO- GENA	Nitraties.	2
N. 1	pended.	#
NIC GEN.	-sns	
ORGANIC NITROGEN.	Dis-	21 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Total.	21 = 32 + 32 + 32 + 32 + 32 + 32 + 32 + 32
Sc.	E p,pud	86.25
EN A	Albuminoid Proff	<u> </u>
NITROGEN	PioT	B S E S E S E S E S E S E S E S E S E S
Z.	Free Am- monia.	<u> </u>
ED.	Matter.	
OXYGEN	By Dis.	$\frac{\mathbb{E} F \tilde{\sigma} \tilde{\sigma} + $
0.65	T'otal.	$\frac{\text{$\mathbb{Z} \otimes \mathbb{Z} \otimes \mathbb{Z} = \mathbb{Z} = \mathbb{Z} = \mathbb{Z} = \mathbb{Z} = \mathbb{Z} = \mathbb{Z}}{\text{$\mathbb{Z} \otimes \mathbb{Z} \otimes \mathbb{Z} = \mathbb{Z} = \mathbb{Z}}}$
,91	пітоІПЭ	သြား ရှာ ရှာ ရှာသွေးသွေးသွေးရန်သွေးနှင့် မြန်မာ မြန်မာနှင့် သွေးသွေးသွေးသွေးသွေးသွေးသွေးသွေးသွေးသွေး
Color	on Igni- tion.	
ż	E p,pnd	# 0 1 2 C C + 4 C C C C C C C C C C C C C C C C
ATIO	sid b'vios	8 4 4 4 4 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8
RESIDUE ON EVAPORATION.	1,10,I,	8 8 4 4 4 7 8 8 8 8 8 8 8 8 8 8 8 8 8 8
on E	benged gus-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
DUE	-sid	
RESI	.lato'l'	x + y x 4 4 4 4 4 6 6 x 6 - 4 y x 6 4 4 y 4 6 4 4 x 9 9 9
	134541	
	TobO	000 000 000 000 000 000 000 000 000 00
<u></u>	Color.	FERRETE FOR SERVICE SE
RANC	0,5	**********************
APPEARANCE	Sedi- ment.	ons'd Little .ons'd Little
7	7. E	
		enutical enutions of the contraction of the contrac
DATEOF	Examination.	No. 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Dan	1899 1899 'ol ec- Examition, nation.	==:::===:=============================
11 .0%	Lidas	60 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0

Trrbinity * Decided. § Very Decided. † Distinct. † Very Slight. Slight.

Coron M. Muddy, VM., Very Muddy, T., Turbid, C., Clondy, Color on Lanthon-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Cayish. Black. GB., Gray. Brown.

TABLE 7.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois,

SOURCE OF WATER-ILLINOIS RIVER, MORRIS, ILL.

No. of Bac.	per Cubic Centi- meter.		455.000	1,397,000	717.500	000.020.6	111 000	112,000	1 000,000	1 230,000	5 850,000	oostoosto.	5 890 000	6.550,000	4.890,000	1.285,000	383,300	380 000	115 000	1 090 000	000,000	2000.000.×	90E 000	955,000	245,000	170,000
	Presence Abs. of			+	<u>-</u>	-					+	- :			:	+	+	-1	+	+					+	
ture ent.	Tempera O'rik lo	17.	33	83	06	3 2	10	, oc	55	13	7	14.	13	17.	14.	16.	0	12.	10	16.	. 6	· -	· 1			, x
	rempera	21.	35.	26.	3 8		9.1	: 5	57	0.10	33	21.	35	26.	33.	30.	17.	10.	20.	22	2	. 2			n u	- =
10 1	Heigh otsW	3-	6.8	.6.		 					6.3	30	5.1	4.1	4.1	5.33	5.6	5.4	5.6	31.0	0 10			: c	0 2	2 :
TEO-	Nitrates.		39		•			_															-	-		5 ∞ • ⊕
NITEO	Nitrites.	035	99	90.	•																		_			
le EN.	Sus- bended.	15	16.	2												_	Ţ			7	9	•			5.5	35
ORGANIE NITROGEN	Dis- solved.	1.18	1.23	<u>장</u> .	•	1.144	-	•	-	4=		_	_	_	_			$\overline{}$					-		1 44	1.64
N.O.	Total.	1.93	2.17	1.73			-		0.4	_	_	_		1000		15000				-	_	1100	_		-46	283
00	p,pud -sng		256		16.	224	1.5.	191	288	388	12.	.416	.352	.416	. 192	.816	107.	.352	. 128	576	7.36	S.	200	33	3 × ×	3.52
EN AS	Albuminoid Pis- Pis- Solv'd	.544	544	.624	512	448	528	.336	-48	.448	.48	.448	.48	191:	.448	87.0.	.576	809	.384	.608	416	31.44	839	514	707	.98:
NITROGEN AMMONIA	Tot'l	88.	œ.	.736	672	672	7.68	σ ₀	.768	736	3.	1864	.832	88.	.64	1.344	86.	96:	513	1.184	152	2	5	P. C.	587	.312
Z	Free Am-	8.8	3.6	5.6	. x	7.52	∞ ∞	5.6	7.04	7.68	13.		<u>.</u>	<u>+</u>	10.4	8:2	13.2	13.6	∞ ∞	8.0	0	00	1. 5	· 00	0 7	5.6
ED.	By Suspd	€. €.	- i	3.0	. 7.	8:	1.7	3.9	1.7	3.1						_					-	_	_			
OXYGEN	By Dis- solved.	1 .	11.4	04	15.2	13.5	7	11.6	3	14.2	11.9	13.3	13.	13.7	13.9	14.3	8.11	1.1	9.1	8.6	10.9	10.5	8	0 0	- x	11.9
000	Total.	_	15.4		<u>1</u> -	15.3	5	5		1	16	16	5	7	7	1,	18		10	=	Ξ	3	Ξ	: =	13	13.3
ne.	Chlori	44.2	: :	43.5	∞	57.5	7.1	37	58.	83	70.9	79.	3	33	33	-38c	97.5	9	62.5	75.	73.4	33	51	50 3	36.5	49.5
Color	on Igni- tion.	B.	<u>ت</u>	m m	===	2	~	.: ::	Gb.	7	Gh.	Ch.	Cp.	Gh.	==	j.	<u>=</u>	GB.	≃:	3	Ö.	Ü	~	2		3
ż	i p,pud		es: .₹0		. 00	8.01	5- C5	3.6	©; ∞	10.4	3.6	:	1.6	37.6	Δì.	00	17.6	∞ →	©. ∞.	~~	~		33	70	5 TC	ાં
LAT10	Solvid b'vios p'vios	54.	53.	31.4	- F(2	∞ ?}	40.4	46.	55.652.8	T	16.8,43.2	:		0)	÷	39.3	17.6	31.2	18.	18.	16.	33	0	30	, T	46.
APOL	L'1oT	63.2	55.2 52.	39.68	51	33.6	47.640	49.646	55.6	64.454.	46.8	57	39.638.	96 8	14.4.42.	÷0.	35.217.617	36.	8.08	21.2	24.4	35.00	13.9.40	35.6	33.6	48.
RESIDUE ON EVAPORATION	Sus- pended.	36.	φ. 200	8.9	16.8	9.7	6.4	60.4	9.6	20.	φ. •		e; ∞	9										12	7	12.4
DUE	Dis-	19.3	8.45	2 420.4	355.6	395.6	67. G	312.8		01.2	₹.0c	:	395.6	10.4	133.5	361.2	7:4:	114.4	28. 28.	13.	.0.	%	.8 398.4	372.4	6 397 6	389.3
RESI	Total.	445.2 419.2	02.83	197.3	372.4	103.23	151.6 115	0	432. 4	121.2 401	425.2,420.4	391	308.43	446.4 110.4		444	153.6.144.4	448.1	331.8 328.8	390.4 372	383.6 370.	389.2 382.	404.83	384.43	401.63	401.63
	iono	None 4	*	-	: **	7	-	-	(2)	-	-	20:	00	·	-ji		र् ग '	ক 	ಣ —			ee .	-	**		
-	robO					_	•	3 Gas'y			-	2-1		· ~	•			-	~	-		•		~		None
CE.	Color.	F.4	. 는 당:		. ₹. ??	F.2				1.04		N F. O.	. F. LD	N. F.O.			. F.05						. F.06			. F. 3
ARAN		d N	2	=			7.		N.	d M		Z	7	7	7. 7	Z	Z	7	7			_	N.	N.	7	Z
APPEARANCE	Sedi- ment	Cons'd	7.	*	c.ons.d	9.0	:	:	Little	Cons'd	Little	:	:	Cons.	•	:	: :			:	Little	Cons'd	99	* *	:	:
	'Y'dan'l'	+ 0	10	*	:+	+	4	*	1 92	*	÷	+	-	* 0.0	*	50	198	*	*	+	+ =	+	15.		+ 88	
OF	1899 1899 Collee- Examition.	lay 3	une	- ÷	:	uly	:	: 1	s1 :	Ang.		:	:	•	Sept.	:	۲۱ :	Oct.	:	: 25	٠,	Nov.		:	3	Dec. 2
DATEOF	9 ее- Е п. п	200	e 6.June	es 0	2 67	3.1	=	1.7	કુટ	31.5	(-	7	25	호: -	· ;	2)		2		??	30	9	1-1	8	63	
I	1899 Collee tion.	May	June	::	:	July	9.9	:	:	:	Aug.				Sept.			Oct.	:	:	,,	Nor.	3.9	9.9	•	Dec.
,6M	Serial	5112	516	5209	5239	5332	5393	5428	5484	5529	5570	5624	5672	5738	57,65	5815	5930	3886	507.4	6143	6190	6235	6294	6350	6395	6590

Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Blk., Black. GB., Gray-Brown. Turbidity—* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. Rh.. Reddish. W., White. Gb., Grayish.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-FOX RIVER, OTTAWA, ILL.

IUR W. PALMER.	BURGILL.	niversity of Illinois
oning oning		
IL.	-	j
100		
-	-	
O Ĉ		
Report of		

No. of Bac	Cubic Centi Meter.	18	1.500
	jo salt	1111 : : : : : : : : : : : : : : : : :	
	Temper of Air.	तंत्रतंत्रतंत्रतंत्रतंत्रतंत्रतंत्रतंत्	0
ature er, C.	Temper		0
10 11	Heigh In'M	<u> </u>	
NITRO- GENAS	Sitratics	8.8.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2	
7	.bended.	※ 2 2 3 3 3 3 2 2 2 2 2 2 2 2 2 2 2 2 2	
	sofred.	に 20 20 20 20 20 20 20 20 20 20 20 20 20	
ORGANIC	-81(1	<u> </u>	
	Total,	8	
* . :	sns	######################################	
NITROGEN AMMONIA	ž sid	824	
NIT	monna.	1212	
	Free Am-	1-10 01 01 - 10 0 0 0 0 0 0 0 0 0 0 0 0	-
OXYGEN	By Dis- solved, By Suspd	$\text{adagagagragagraghragagragagragagragagrag$	6 7
Cox	Total.	$\log \log $	7.1
	то[пО	$-\omega \sin + i \pi \cos + i \pi \cos \cos \cos + i - i - \alpha \cos + \alpha \cos + i - \alpha$	5.
Color	Igni- tion.	東京戦略はおは発命の発行の発行のははのははのはは	::
o.x.	g p,pud sns	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
RATIC	Loss on Sid		86.38
1 2			
0N]	Sus-	- 20mm - 1mm - 2mm - 2	
SIDUE	Dis-	1060 4 20 20 20 20 20 20 20 20 20 20 20 20 20	<u>x</u>
RE	Total.	1066 4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	130.8
.1	opO	Non	:
53	or.		F.02
RANCI	Color.	zzz z zz zz zz z	
APPEARANCE	sedi- ment.	Cons'd Little Little Cons'd Little	:
	Turb).	8	183 183
DATEOF	1899 Exami- nation.	Second Se	:
DAT	1899 Collec- tion.	0	: 1
	5 ~	50113 Nay 50163 Juni 50113 Nay 50163 Juni 50	

Colon- M., Muddy, V.M., Very Muddy, T., Turbid, C., Cloudy, Trebbury-*Decided, § Very Decided, * Distinct, ‡ Very Slight, Slight, C., Co.or.-M., Muddy, V.M., Very Muddy, T., Turbid, C., Co.or. on Identics -G., Gray, B., Brown, D.B., Dark Brown, L.B., Light Brown, R.B., Reddish Brown, B.G. Brownish Gray, Bh., Brownish, R., Red, R., Black, G.B., Gray-Brown

TABLE 9.

STREAMS EXAMINATION SANITARY DISTRICT OF CHICAGO,

SANITARY WATER ANALYSIS-PARTS PER MILLION.

· SOURCE OF WATER-ILLINOIS RIVER, OTTAWA, ILL.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois...

No. of Isac.	per Cubic Centi- meter.	12,800	12,045		00000 000000 0000000000000000000000000	9.830	11 (500)	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	37,400	21.000	42,000	56,000	6.200 0.000	000.15	36.600	28,000	3.000	8.000	CH. 750	:56.CCC	212,000
ilo')	Presence Abs. of		: :				:		-	F .	+		1-	-+	- 1	+	-	-	+-	+-	+>-
	Tempera	8.8	:6 :6 :6 :6 :6 :6 :6 :6 :6 :6 :6 :6 :6 :	: :	S S	33.	<u> </u>	35.	88	· ~	35	·	કું કુ	38.	233	جَرَ	-0	=	x ;	_;	
9.111.1	Weinpera		3.57 19.27				÷ 10		~	9					9					2	9
	 Heigh	-					00 00 00 00 00 00		ئ د د د د				25 00 00 00						, , ,	2.5 25 52	ာ က
NITRO- GEN	Solutios.	.5 1.28 .6 .6	.575 1. .75 1.	re n	ີ່ຄຸ− ວິດ ວິດ	875 2	95 -	6 6	1255 0.35 0.35	375 1.	375 2.	e						2000	1	013 1.	05.1
	bended.	13.64							-												
ORGANIC NITROGEN.	Dis- solved.	.984	986.	1967	20 00 10 00	856	9.99	744	824	32.	.836	.836	5.5	89	.648	-12	.488	936	# S	205.	.936
QN	Total.	2.2.	1.21	라. 라.	17	1.16	300	1.56	.856	1.16	1.06	6.	3 -	.92	.744	1.06F	 %9: 	1.196	930	200	1.256
vo.	E p.pud	.032	₹ 8 8	80.	.16	308	9.69	272.	.013	208	.064	.032	888	0.83	.064	80.	960	. 138	010.	3	224
GEN AS	Tor'l Dis- solv'd	.512	÷ +:	304	1 65 1 65 1 65 1 65 1 65 1 65 1 65 1 65	301	.416	.416	355	288	448	.432	. 368 258	352	.33	.435	.352	4.	011.		304
NITROGEN AMMONIA	TotT	.544					2000	.688	.364	965	.512	164	+c. +	.384	385	.51 <u>2</u>	277	5258	700	2000	528
	Free Am- monia.	1.36	ಯ ಸಧ	က	7	· ·	ب من	33	8. 8 8. 8	7.68	6.4	7.07	x x		00		XX (24. F	0 1		5 or
EN IED.	By Suspd	.1					. es	ভঃ		ಞ	1.8	? ? ?	0	. n.	0.0	9.	7.	nj.	:	2.5	1.50
ONYGEN	By Dis-	15.8	<u> </u>	<u>ن</u> ق	323	= 3	2=	5	210.9	00	00	= "	JD 0.	00		(9	(-	: 5	ું ≎	9 10.7
55	Total.	15.9	##	<u>ক্র</u>	2 = =	<u> </u>	===	=	==	Ξ	10	= 9	ה ס	œ	3-	- :	9	:-:	: :	= 14	==
-	Chlori	30.	65.3	47.	3.4	<u>. 1</u>	69	75.5	20 20 20 20 20 20 20 20 20 20 20 20 20 20 20 20 20 20 20 2	30.5	82.5	6	6 G	69.5	61.5	දි	30.	<u>و</u> ي	11	96	24.5
Colo	on Igni- tion.	ಷಷ	ಪ್ರಜ	ರಲ	9 09	<u>ن</u> د	50	Ö.	<u>ئ</u> ئ	izi	G	<u>ن</u> د	<u>.</u>	Ö	<u>ت</u> :	ž,	E	<u>ن</u> د	i a	ے ≃	
	z p,pud	ॐ ः		→; c	0.0	17.2	:0)	∞ 7	2 × 0 ×	3.6	ė.				• 0	. s	δ. α α	9.8	3, - 1, C	- c	1.6
AT10	Loss on Sid b'vlos	53.4 23.6	26.8 46.8	5. 15 5. 15	64.4 64.4	55.6	49.3	₹ .∞	51.249.6	30	F. F.	43.640.4	N → 0	19.6		× 0	200	ο : : :	5.00	92.4	21.8
VAPOI	1.10T	9.1.6	43.6.26	39.639	64.4	5. S. S.	ت ت ن	47.6	21.08 25.24	33.6	30.4	43.6	57.0	20.	19.6	36.	× + ×	2. I. S.	0.12	0.00	100
RESIDUE ON EVAPORATION.	Sus- Sus-	9.6	17.6	54.		: -ري د زي		21.6		14.4	13.9	ر مورد د	2000	15.6	21.6	20.4	1.0	±.	0 10	0 9 0 8	12.4
IDUE	Dis- solved,	383.2	374.4	340.4	396.4	369.6	392.8	411.2	436.	414.	388.3	440.	319.6	356.4	319.2	309.0	508.	350.	965 6	239	327.2
RES	Total.	00 00		394.4	419.2	376.8	393.2 392.8	432.8 411.2		138.4		444.	366.8	37.9.	00	350°	304.4	368.4	0.100	361	.6
	tobO	Je		Gas'y		: :	:	*	19 -0	•	:	9 4	;	3,	3 7						:
CE.	Color.		도 본 연 양 양	F. 19) ?! : Eu	. F. 05	1.18		F.F.				E.E.	F. 1	F. 1	CO. 7	3.5	35			
APPEARANCE		e N		d N	5 7		ت م ده ت	d N	e		Z.		*)						7		7
APPE	Sedi- ment	Little	::	Cons'd	• 4	Little	Cons a Little	Cons'd	Littl	**	99	Consid	91717	9.9	* *	: :	: :	: :	" "	('ons'd	:
	Turb'y.	37 +	++	* +	3 21	00;		* 67	<u>ာ ္</u>	6:1	36 +	+ 4	2 00	:S	- 1		<u> </u>		=+	- + + - + - + - + - + - + - + - + -	* 65.
EOF	1899 Examination	June	July	: :	Aug.	33 1	: :		4 Sept.			Oet.	3	•	Nov.			a .a	Dog	Dec.	9 .
DATEOF	1899 1899 Collee- Examition.	d3	uly 3 J	17	. 331	Vug. 7	± 57	25	. —	18	3.5	:t:	91		စ္တ	0 :					200
70.7	Serial		<u> </u>	5432	5531		5677		5798 Sept	5881		5995 Oet	5084	6153	0.5	0241 NOV		3333		0504 Dec	5595
1		15.15	20.00	າຕໍ່ ເຕ	: 18	iĝ i	ನ ನ	2	ñ. ñ.	51.	50	i ai	5 3	6	9	0	0	9	5 5	3 E	3 3

Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Turbidity-* Decided. § Very Decided. † Distinct. ‡ Very Slight. Slight.

COLOR ON JGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-BIG VERMILLION RIVER, LA SALLE,

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

11.200 3.210 1.950 17,680 \$ 4 6 6 8 E 000 (F.X. 3,(80) per Carbic No. of meter. Bac. gentheraure ornheminio Changeria (*) Water. 21.9.2 to maist GENAS Salualix NITRO-888 200 200 310 310 310 130 850 Samanis pepued ORGANIC NITROGEN. solved. -SIC Triotal, Albuminoid Am. p.pad sns NITROGEN AS Pis: AMMONIA. Log. Free Am-monia. platter. 27 CONSUMED. ONYGEN sid ya bevlos Total, (hiorine. 'olor Igni-tion. on
 1994.8
 168... 26.8
 67.2.56.4
 10.8
 13.2.4
 10.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16
 13.2.3.2.16</t p_ipud -sng EVAPORATION. 50 sid b'vios on 350 1,10,1 рөрцөд RESIDUE ON -sns neartos ·8](I дидод, Tobo Color. APPEARANCE. Little * ('ons'd Little Little Sedi-Cons'd * Cons'd Much Little f,q.m.j. +== 45 Examination. 0681 Sept. VIIK. -《记载器中国函数 0031 5079 May 2 5115 :: 3 5159 June 5218 :: 1 5319 :: 2 6216 Nov. 6257VIIE. Will OFS 5999 Oct. SHEH Dec. 5440 5,410 6500 STOCK OF Serial No.

C. Cloudy. Brownish. R., T., Turbid. Reddish Brown. BG., Brownish Gray. Bh., Very Muddy. V.M.: Muddy. ('01.018 - M... DB. Dark Brown. LB. Light Brown. Reddish. W. White. Gh., Grayish. SHght. * Very Slight. + Distinct. Brown. Color on fanition-G., Gray. B., & Very Decided. Trreibity * Decided.

GB., Gray-Brown, Black.

TABLE 11.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LA SALLE, ILL.

Report of Arthur W. Palmer. T. J. Burrill. University of Illinois

No. of 13ac.	per Cubic Centi- meter,	# 1500 # 1500	138.500 123.500
Golf.	Presenc	<u> </u>	++
	Tempers, (
31, U.	rempere of Wate		
:16	dgi9H ota7/		==:
TRO- GEN AS	Nitrates.	0.00.00.00.00.00.00.00.00.00.00.00.00.0	3.12
NITRO- GEN	Nitrites.		
IC SN.	-sus bended.	854 - 146 -	5.0
ORGANIC NITROGEN	Dis- solved,	1000 20 20 20 20 20 20 20 20 20 20 20 20	1.128
O'N FIN	Total.	888844444488558558585858888888888888888	288
on a	i p,pud	- 8 + 8 4 8 4 8 8 5 ± 5 8 5 5 5 8 5 5 5 5 5 5 5 5 5 5	3.1.
< ;	Dis-b'vlos	<u> </u>	.352
NITROGEN	Albuminoid Pis- Pis- Solv'd	**************************************	.496
2	Free Am-	6.88.48.86.86.86.88.48.48.48.48.48.48.48.48.48.48.48.48.	ن د ن بر
Ν Ε10.	By Suspd	05034000 00003 -03-4-0040 055- 0 04 0505 +00000000000000000000000000000	3.57 3.52
OXYGEN	By Dis- solved.	@ Q @ @ @ E E G @ C C @ G @ G @ C C C C C C C C C C C	3. O.
Coy	Total.		13.4
ne.	Сиют	<u> </u>	33. 33. 33.
Jolor	on Igni- tion.		33
-	i p,pud	2012 25 24 28 24 28 26 26 26 25 25 25 25 25 25 25 25 25 25 25 25 25	25 41. 41.
AT103	ross on less o	19.19.29.29.29.29.29.29.29.29.29.29.29.29.29	53.631.2 62.8 62.4
APOR	Tot'l	2003年48年88年88年88年88年88年88年88年88年88年88年88年88年	62.8 62.8
RESIDUE ON EVAPORATION	Sus-	်ပ်သံသံသံသံ ပေးသံသံသည်။ သံပါသံသံသံလ်လံသပ်ပေးလံတ်တံ	× +-
DEE C	Dis- solved,	402 403 404 405 405 405 405 405 405 405	470.8
RESII	Total.	+68 +68 +68 +68 +68 +68 +68 +68	474.84.4
			र्व ∓
.7	obO	0 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	:
CE.	Color.		F.03
ARAN		ANNERS ESSENTIALES ESSENTIALES	
APPEARANCE	Sedi- ment	Consider Considering Much Considering Much Considering	Little
	Lucp, Z.	85-528-208-208-2088-2088-2088-2088-2088-	+= 0.89
OF	1899 Exami- nation.		
DATE OF	ec- F n. n	47	96
	1899 Collec- tion.	May. June Jully Sept Oct Nov Dec.	: ;
No.	Serial	200 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	6587

Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Jenition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bb., Brownish. R., Red. Grayish, Blk., Black. GB., Gray-Brown. TURBIDITY-*Decided. \$Very Decided. † Distinct. ‡ Very Slight. | Slight.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER--ILLINOIS RIVER, LA SALLE, ILL.

Report of Anthur W. Palmer, T. J. Burrill, University of Illinois.

No. of Bac.	per Cubic Centi- meter.	1	84,000 190,000
HoU	Presence		:++
nuə,	Tempera of Air, C		
	Pempera		
	Heigh ots7/		
NITRO- GENAS	Zitrites.	$\frac{165 \cdot 187 + \frac{16}{12} \cdot $	\$ 50 m
	pended.		138
ORGANIC NITROGEN.	Dis- solved,		S. C.
O PIN	Total.	<u> </u>	3 ×
w w	E p,pud	888 888 88	22 2
4 ;	T'10'T Dis- sid biological	· · · · · · · · · · · · · · · · · · ·	3, 33
NITROGEN	I'10T	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	648 1381
	Ртее Аш- топуя.		
EN MED.	By Suspd	<u> </u>	
OXYGEN	By Dis- solved,	$\frac{\sin (-\alpha \sin \alpha \omega)}{\pi \sin \pi \sin \pi \sin \pi} = \frac{\cos (-\alpha \cos \alpha)}{\sin \pi \sin \pi} = \frac{\cos (-\alpha \cos \alpha)}{\sin \pi} = \cos (-\alpha \cos \alpha)$	0.0
	Total.	<u> </u>	
	(,pjorh	\$\$\$\frac{1}{12}\$	<u> </u>
Cold	on Igni- tion.	ක්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්රා රාජ්ජාවේ ක්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත්ත	± ± ±
ox.	= p.pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 x =
RATI	For Paris	######################################	
VAPC	3 1'10T		15:54
ON F	bended.	84246583854583865858567848564658	= 33 = 23
RESIDUE ON EVAPORATION.	-Si(l		2.58 2.58 3.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50
Ray	Total.	### 19	1
	tobO	ou .	::
	or.		F.30 F.06
RANCE	Color.	NANANANANANANANANANANANANANANANANANANA	
APPEARANCE	Sedi- ment.	Cons.d	. :
	Turb'y.	++ W+ 0 0 + * + * + * + * + * + * + * + * +	+ %
OF	1899 1899 Collee Exami- tion, nation.		용취
DATEOF	1599 Tion.	the state of the s	28
		May New York Way N	:::
1.0%	Serlal	2	55.50

Conor - M., Muddy, VM., Very Muddy, T., Purbid, C., Cloudy, Color on Lanthon G., Gray, B., Brown, DB., Dark Brown, LB., Light Brown, RB., Reddish Brown, BG., Brownish Gray, Bh., Brownish, R., Red, Rh., Ch., Gray-Brown, Truendity * Decided, § Very Decided, * Distinct, ‡ Very Slight, Slight.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, HENRY, ILL.

T. J. Burrell, T. J. Burrell, University of Illinois.	
Jo	
Report of	

Nσ. of I3ac.	per Cubic Centi- meter.		8,850	158,000	10.800 10.800	15.500	00010	2,400	77,000	2,450		001.1	39,600	4.550	6.300	(SE)	0.150	00000 00000	6.000	10.00	3,100	1,000	2.000	10.800	15,900	51.000	113.000
	Presence Abs. of	:	8.		:	ļ	+	-	Ŋ.	2.	:	ı		+	+	1	, ,				1	ı	+	-	+		+-
ытиге Этите Этите	rempera Of Air, C	55	× 5		 	33 22	7 7	3.1.	.56		?} ?} ?		· ·	÷ ₹		ે જ	11.		16.	<u>.</u>	1-	1-	{-	6.	6 -		-11
	rempera		Si S	€ &	30,00	F 5	000	31.		26.	9.00	000	9	<u>ئۇ</u>	<u></u>	<u>∞</u>	. e						œ.	эō	٠ ٠ ٠	25.0	
101	Heigh otsW	ος σς	∞ o ∞ o	000		. j.					00 t					<u>; -</u> ;	- 7	- (-	. {-	, 20 100	1-	}	7.6	2.7.6	∞ -1 -3	6 7.6 5.0	zi.
NITRO- GEN AS	Nitrates.		% 	-		بياد			-	_						- 7		· -								S5 1.36	
NIN	Sitrites.		# -																								
NIC GEN.	-sus		##G: :9#	. D.		-															- 1						
ORGANIC NITROGEN.	Dis- solved.		986: -			- -	•				-	٠	_														
	Total.		1.53 1.53	. 20			•					-			_				4=							_:	_
SV.	my pind sugar		113		<u> </u>	•						_	• •			•										•	
	Albuminoid Pivior Divios		7.78					•				•				•	•				•						
NITROGEN AMMONIA	[1.10]	576	508 808	1.34	.67	D = 7	200	3++.	<u>6</u>	154	277		- T- C-	H.C.	.656	χ, δ	6.5	09.	56.	.56	.43	-448	. 433	.43	7	57.0	.00
	Free Am-	968.	.576	£.	o; 3	30.0	1.03	.38	1 9.	387	20.1	3.8	1.15	_	υž.	1.12	5 . o	· +	10	3.52	∞. ₩.	3.53	2.56	٠٠ ٠٠	3.68	÷ 50	7.97
EN MED.	By Suspd	-	را الم الم الم	_	00 € 00 €		33.		e5 €5	5 1.		÷ -	1 4	5.5	0.0	% - % -	0.00		1	6.	-	∞. ∞.	**	$\frac{6}{6}$	0.0		F:
OXYGEN	By Dis-	<u> </u>	2≧ ⊆	2	<u>ක් අ</u>	2 =	2	22	33	=	= 3															 	
	Total.		5 16.1						0				10														
	a.∓ = Chlori		0.12				· 53	<u>ئۆ</u>																			
_	Jgmi- tion.		0 to				_						<u>∞</u>	≃ ≃	ن 	<u>ن</u> د	5 C	9	S C.	<u>.</u>	6.		ن ت	0 8.	ۍ; ټ	¥: :	· .
ION.	p,pud -sng	33	4.21.6 6.9.6	3	_				6	2	9.5	0				9.0			6 3.	33	-	6.	-	0.8	-ji -	-; -; τ	· ·
OKAT	Loss on 1'10'l' as on b'vlos on b'vlos	.2. 84.	39.2.29.6	23.620	32.8 32.8 11.4 11.4	2 2 4 2 3	£	6 44	39.2.50	54.843.	0.000.41	4 39	76°	8 52	25. 25. 26.		2 7	87	8:	41.2 19.2	29.6 28.	41.633	88	48.8 48.8	36.432	5 5	0.0
Evar	pended.		88.6 60.6 89						56. × 07.		7 6						0.70				17.6 29		9.6 33			17.2 49.	
RESIDUE ON EVAPORATION.	-sns					96		~	우 i	1 2					_												* ·
ndisa	Dis- solved.	2 378.	316.	391	T. 64.7	129.6.402.	343.2,255.2	347.2 326.	105.6364.	2.374	301	388	123.2 386.8	1385.3	4.468	2555.12 27.67 3.12	441 9 435 6	139.2 122.4	389.6	3375.6	396.8 379.2	1368.4	372.	2 356.8		360	
RI	Total.	165.2	376.2	168	136. 286. 286.	1.001	343.	3. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	105.	129	410.4 306	113	<u></u>	4.38	.122		1	439	+08	107.6	396.8	388.4	381.6	357.2	384.4	25.1	. 100
1.1	орО	None	: ::	*	: :	;	:	:	:		: :	• •	:	;		: :	:	:	;	:	:	:	4	9 :	ag 11	:	
6	or.	F.05	프 다 ::	E.1	ਜੂਜ਼ ਜੂਜ਼ ਜੂਜ਼	1 F	£.3	F. 15	3} (E.04	5.5 5.5	5 03	F.03	F.05	20.75 20.75 20.75		F. 15	F.05	F.03	F.0.1	E.0.1	F.0.	.03	.03	F.03	라. 라.	(.1.
RANCI	Color	_; _;			===		=	Ę.	= :	= :	= =	-		Z.	, ,					N.	<u>,</u> ;	N.	7	7	7. 7		. 14.
APPEARANCE.	Sedi- ment.	Little	Cons.d	Much	Cons'd	*	:	:	:	•	: :	:	:	:	: :	: :	:	:	:	:	:	:	Little	:	3 3	f.one.d	n ciio
	Furb'y.	31 *	* *	*	÷ * ∞; °	*	* 61	* 17.	י א	÷	15.	- ÷	+	13 *	19	* *	+=	<u>~</u>	* 97	*	00	*	÷	÷.	_+ ; ;	- + - 0;	- 00
EOF	1899 Exami- nation.	30 May	6 June	:	ží Tuly	CIII :	;	13	.Vug.	9 3	: ;	7.7	Sept.	"	9 3	: 0	:::	"	:	Nov.	10				Dec.		j
DATEOF	1899 1899 Collec- Examition.			200	**	_		97	35:		<u> </u>		9	2			. –		33	-	-1	+		31 ∞ 00 1		2 0	
			65 une	99		: 10	: :		Sur le	26	500	;	II Sept	61	: :	-		:	69	12 Nov	:	:			49 Dec		
No.	Serial	5131	5105 5205	5256	5300 5300 5000	24046	5113	576	5551	200	5688	52.5	58	200	200	20.0	6042	8809	6919	6212	6260	63	63++	6407	6113	85.78 87.18	3

TURBIDITY -* Decided. § Very Decided. + Distinct. ‡ Very Slight. | Slight.

COLOR ON IGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red.

T., Turbid. C., Cloudy.

Color--M., Muddy. VM., Very Muddy.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-ILLINOIS RIVER, AVERYVILE, ILL.

Report of ARTHUR W. PALMER, T. J. BURRHLL, University of Illinois.

o of Bac.	per Cubic Centi- meter.		:		4.650	9,100	12,350	13.500	×.050.	160	7. ICE)	2000	21,000	1,950	<u> </u>	10000	2000	0.00.1	2,000	1.975	0000	0.600	2,556	550	1.100	1.2(x)	€£;		3.900	13,600	10,400
,ilo')	Presence BCC BCC		:			:		+.	٠.	:					,	1				1	1	-		1	1	1	4	:		+	-
:ent:	Tempers Tempers, (:	:	36.	100	∞ 201	- - - -		20.0%			:2:	:26:	÷ 13:	55	: : 13:	=	<u>.</u>	<u>;</u>	<u></u>	<u>x</u>	x.	x.	<u>.</u>	21	-	?	?}	e i	
ernn	Tempers		:	:	: :: :	56.	36.	92.5	÷ 15	3 %	18	ज	23.	.56	£ 1			: 2	.: ::	<u>5</u>	<u>×</u>		===		.	.01			5.	-	.0.
10 11	Heigh Dig/	5 32.	6.29.9	20 20 20 20 20 20 20 20 20 20 20 20 20 2	18 E	59.4		6.27.1				27.6		35.7	26.7	52.50	6 27 . 3	141	52.55	1.1 7.1 66	9. 17. e	57.6	α. 	× 55 0	7. X.	25. E.	11:38:1	2.75	25.28.33	28.5	3.05.
NITRO- GENA	Nitrates	6. 18	6: 15: 15: 15: 15: 15: 15: 15: 15: 15: 15	6. 1. 20	25. E.8.	ري 	9 <u>-</u> 92	6 - C	2		-	5.1.4	_	₹. ??	_ :	_	T. T. 60		5 3.2	1.1	3.6		33.0	3.0	-,	7. 1.6	8.1.8	31 %	35 25	- ×	1:
<i>Z</i> .	bended.				- 19 - 19 - 19 - 19 - 19 - 19 - 19 - 19																_	_	_							192	16
ORGANIC NITROGEN	solved.				986			_																_							
ORG	T'otal.				5 67		_																								<u>x</u>
	p.pud sns-		-		80	_	_	-				_	-	_		_				=	=	-							=	1810.	980
EN AS		.224	.416	3033	416	384	.432	.368	.336	2000	886	3.00	16:	2000	883	000	200	.116	.3:36	.3336	=	~!:	30.	:301	22.5	大学?	.176	.2565	-	. 435	3336
NITROGEN	Albuminoid Dissipation of the state of the s	 27:	210	416	524	.448	.513		544	0 10 2 0	200	35.	.416	114	.576	917	21 G	513	505	. C.S.S.	191	- 5	<u>x</u>	- X	911.	<u>x</u>	€	135	<u>x</u>	<u>x</u>	. ES
Z	Free Am- monia.	808.	£98.	96.	388	.088	.256	# :	202.	0.1	000	91.	.416	÷:	27.5	200	000	8 8 8 8 E	. 158	193	.356	27	#	23	27	3.55	1.92	1.6	-	-	3.36
EN MED.	By Suspd Matter.	3.2	4.6	4.0	- 1	01 02	1.3	≎i		000	01	_	5-1.3	35	0.00	3.5	-	2 00	_	× -	-	~	ig.				0.0	_	∞. ~	- 1	rc.
OXYGEN	By Dis-	.9.8	6	.6.10.		.3 11.1	12.7	<u>ි</u>	.5 3	e ≪ - ≪	101	2 10.8	.9 10.6	5,10.5	G 1	× .	€ 10°×	- ∞	. 6.	9.6	x. x.	x :))))	.5 6.1	.1 6.0	. 1	?	7. 6.7	 8.3	× ×	9.0
	Chlori Total.	112	ा ा	<u>*</u> =	<u>, 10</u>	<u></u>	_	_	يناده	31.		22	_	≘ .	= ;		E :		Ξ	E :0		.c	οc 	\$	1-	10	}-	1-	0.00	<u>ි</u>	15. C
	on Igni- tion.	3. 13	3.	5. HO	21-	3.	 ?!	 33							material		-	12, 38						5.	3. 51	3.		i. :30	i. 39	·:	.: ::
	e p,pud	-∞.			· · ·		00	61	- 5	200	S		2.0	3.3	_			9.6		0.0	oc.	oc.	:	_ ∝	- -	9.2	∞.	9.7	3 6 (0.0	∞ ∞:
ATION	Loss on Ig.	19.2			30.00		.6 28.8	62.4		- 00 - 00	59.2	<u>x</u>		% ∞. ∴ ∴	× .98	1.00		36.	2.36.8	31.	6 12.8 26	2≧ ∞,	36.8	00 63	∞. ‡‡ ?≀	35.	8.1	× 0	10.8	9.14	× × ×
VAPOR	1.10,1				34.43%.		53	တ		0.00 0.00 0.00 0.00			33.	200	36.8	.32		45.636.		3	330		क्ष		12		13	× ×	-	41.6	€.
ON E	pended.	15.2	55.6	5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5	155.6	333.	45.6		() () () ()	x x	0000	53 6	17.6	50.1	98	25	∞ - ?? ?	15 6	×	16.4	17.6	30.	13.6	15.6	<u>∞</u>	<u>×</u>	÷.	(C)	3.6	16.	÷
RESIDUE ON EVAPORATION	-SIG	305.8	37.4.8	354.		333.9	362.4	356.4		201	333.6	∞. 33. 33. 33. 33. 33. 33.	352.4	337.3	315.2	301.6	360.4 360.4		381.8	391.6	395.1	× = ×	9.70#	3.60.5	8.380.8	351.1	348.	361.1	376.4	338.1	372.
RES	Total.	318.	4:30.4	391.2 354	415.6 200	365.2	107.6 362	400.1356	306.x 374	315 2 201	380	305 +	100.	387.6	375.2	392.1367.6	036 5 366	383.6 368	392.8			× = = = = = = = = = = = = = = = = = = =	121.2 107.6	101.8	308 8	37.2.1	1.000	355.6361.			376.4
٠,	юрО	None	:	: :	:	:	:	•		:	:	:	:		1 :		: :		9.	:	:			:	:	:	4 -9	:	:	:	;
63	or.	F.04	야 도 도	5 -	. E.	<u>-</u>	<u>.</u>	٠ <u>٠</u>	10		??	90.	F. 15	?}	<u> </u>	0.		E. 2	??	·	F.03	× (-	÷.63		F.03	: 0:	F.04	F. 65	÷.0
RANCE	Color.		7.										N.	_	7.	7. 3	7. 7					7.	_			7.	Ξ.	Ξ.		7.	
APPEARANCE	Sedi- ment.	Cons'd	: :	: :	9 .	:	:	: :		:	:	:	:	:	:	:			:	:	:	p.suo.)	:	:	*	9.	Little	:	:	р.ѕпо.)	ř
	.v.qan,L	1:01	* *	* + ~~	+ - 6:	* 	+ 27	% ·	+ +	+ +	*	+ 6	16 +	*	30 *	*	+ +	*	+ -	*	<u>*</u>	4	+	+ x	+ 0	+ ???	20 +	+ 9	+ 52	+ (5)	+ 177
EOF	1899 Exami- nation.	May	:	Tube	211111	:	:	July.	: :	:	Ang.		:	9 0	9	<u> </u>	: :	:	Oct.	:			.70%	:	. ,	. ,	9 9	Dec.		9 9	*
DATEOF	1899 Collec- tion.	May 1	:		ine x	3 :	7.7	115 3	01 : :	01	ug.	x:	15	÷:	र्ह्य ।	l. 5	219	:	: :3	10	:	2.	~	1 :20%	:		× :	Энс. 5	:		97
.o.N	Serfal	_	050 1	5013	5155	5253	50 E	5333 Ju	. 1089	2007	-		96331	· 135					590P2 OC	HBT .	· LYX				. 000	, 6888		-			

Trubibity * Decided. § Very Decided. * Distinct. ‡ Very Slight. | Slight.

Color of Latinos G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddlsh Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Red. Blk., Blk., Blk., Blk., Blk., Brown.

Coron M., Muddy, V.M., Very Muddy, T., Turbid, C., Clondy.

TABLE 15.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER-ILLINOIS RIVER, WESLEY CITY, ILL.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

Color Constraint Constrai
Color Constance Chilorine Constance Constance Chilorine Constance Chilorine Chilorin
Color Constitute Constitu
Color Construent Construent Construent Color Construent Color Construent Color Construent Color Construent Co
Color Constraint Constrai
Color Color Constrain Color
Color Constrain Color Colo
Color Constrain. Color Constr
Color Constrain. Color Constr
Color Constrain. Color Constr
Color Constants
Color Constrain. Color
Color Linning Color Linning Color Linning Color Linning Color Linning
Color Linning State Stat
Constraint
6. 16. 6. 46. 16. 16. 16. 16. 16. 16. 16. 16. 16. 1
6. 16. 6. 46. 16. 16. 16. 16. 16. 16. 16. 16. 16. 1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$
APORATION. 1707 1.085 1.010 1.085 1.0
क् .b9bn9q अं मंथअंध्यम्धमंथ्यं कित्मधंष्यधं मंथ्येव्यं
8 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
RESIDE STATE OF STATE
80 5 5 6 5 6 5 6 5 6 5 6 5 6 5 6 5 6 5 6
Odor.
" "
Color.
Sedi- Colc ment. Sedi- Little M. F Cons d M. F Cons d M. F Cons M. F Cons d M
\$\frac{1}{2} \frac{1}{2} \frac
ATE OF 1899 1899 1990
T
Serial No. Decrial

C., Cloudy. Color on Jenition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rb., Reddish. W., White. Gb., Grayish. Blk., Black. GB., Gray-Brown. T., Turbid. VM., Very Muddy. COLOR-M., Muddy. TURBIDITY-*Decided. § Very Decided. + Distinct. + Very Slight. | Slight.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, PEKIN, ILL.

Report of Arthur W. Palmer, T. J. Burkhin. University of Illinois.
--

o. of 3ac.	per Centi meter.		307.000	097.500	010,500	300,000	510.500	30.04KD	.6:55,000	.073.333	000.000	55.000	40,000	500.000	510,000	000.056	550,000	000.000	900.00	170,000	115.000	000,00	970,000		15.000 2.000	7.(8)(0	
Colf.	to .ed /.	:		 	: :	· ·		, J.	· · · · · · · · · · · · · · · · · · ·	χ.	: 1				-	7		•	i ĝi		nigo.	-	».	:		1 :	_+
) , vi /. 10 onesorq	5. /	· · ·	y. =		.0	ر ان تر	-		20 C		0.			 - £) ,) ,	- 10			6.	.a	Ç.	:		_ e }	-
	ersW to																							:	:	-	
oann 'a	Tempera		12.56		3 45	₹ 6					0.66	क्रिक	\$.	हु है - -	100	21	2) 10		5 10	9	10	0 6.	0 .0.		:	10	_
A S	Meigh Heigh	က် <u>ဖွဲ့</u> က မ		ao s. ∞ s	200	213				?? =	H (SC)	76	#:				Ź			<u></u>	W 2	9		- P	K.	25	-
NITRO- GEN	Soliuliv	044		-	-	-		10101	-	<u> </u>		2	<u> </u>		-	-	- 5 - 5 - 5	200	10	<u>\$7</u>	_	2	(S)	200	200	(55)	
7.	рориад	÷ 689		_				-			_					_	201	68	5	256 1.	31.5	500			25.0	550	
ORGANIC NITROGEN.	solved,	68.													\leftarrow			-	25						012	(% () % (-
ORG,	-sid			-								_	-			_			_		_	_		_	_	7.7.	_
	Total.	3 1.13	-		1							****	-		. 35				**	_	=				-	8.0.	
AS.	p.pud -sus	80.																	-			_				. S.	-
TEOGEN	Albuminoid Dis-sid	84.																							O	. 16.	
NITROGEN	Tot'I	36.3	909	57.6	15	578	844	355	. 736	36.2	673	.752	900	679	1.13	. 752	£ 1.	1.0.	1.61	555	∞:	3 ?	X S	NA.		34.	
	Ргее Анг- пеонія.	8.8	.544 .544	5. 15. 5. 15.	8.	e; ₹	538	:8:	.864	15 E	∞ +	.432	रें	20.E	498	.336	Σ Σ	1.088	98:	1.76	16 16 16 16	xo d		0.10	200	300	
SN (ED.	By Suspd Matter.	65.00	9 00 1	es e ∞ ∞) 03 (0)		+ 9	.5.	- ;		. 9.	9.4	٠) : ن :	0.0	νc. ∞	යා ග	: 67	9.	6.	σĸ.	∞.	· ·	- c	Ç =	- 00	(e)	
OXTGEN	By Dis- solved.	===	10.6				0.0	6		<u> </u>	-		0.0	x m		œ'	: 10	<u>.</u> _	2		17.1	6.7	و د	20	i c	5.5	
000	Total.	13.3 13.3	, 20 + 	7 7 7	15.6			 	5.1		13.7	1.3	5 14.1	7 G	30.	∞: =:	= =				٢- إ				- 0	20.00	
	Chlori	15.00 0.10	20	<u> </u>	? ??	36.	े क्	33	,	S. 18	8	43.	46.5	<u> </u>	56.	55.	90,00	35	8			50.					
Color	on Igni- tion.	0.0° ≃	i mi	mi ==	<u>: ~:</u>	≃ <u>.</u> a	Gp.	Gb.	Gb.	i e	Gh.	Gh.	:		£ 123	<u></u>	<u>:</u>	===	(i.		<u></u>	33	÷ ÷	.: .:	ju	: ::	
ż	e p,pud	00 co	5 25	0.08	-	6.4	<u> </u>	œ	සා : ක :	-	33.		5.6	= -		2i -		- 07	∞ —	1:5:1	30 / 20 1	20.0	- 2	0.0	10	1.6	
LATIO	Loss on L'10T	. 5 38.	00	00 4	00	330	∞	्	63.6	+ 10 20 20 20 20 20 20 20 20 20 20 20 20 20	32.0	37.2	3C. T. S	00 00 00 00 00 00 00 00 00 00 00 00 00	33.6	+:		300	38.8	30 ₹7	× .9	4.44	9.16	401	6 11) ;;;	
APOI	1'10T	7		ණ ∝ි	000	30	5	-	67.13	e; :	55.6	 	30.5	3, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8,						3.00	19.6	9.00	. 0			39.6	
RESIDUE ON EVAPORATION.	popuad- gus-	0.10	108.	#6.8 20.5	31.6	χ χ, ς	9 =	31.6	88. 6	연화	=	₹}	\propto	2000	76.	13.6	2 2 2	-	34.4	∞ ∞	?? ???	2.1	1 1 1	0.1.0	23.5	3 01 3 02	
DI'E	solved.		o oc	× = ×	28.0	8.5	200	51.0	S	2000	364.8		, C 9	(0 (0		α, α,	÷ :	110.	6.4	0	9-1	2) 5	T 9		363 9	₩.0.±	
RESU	Trioff -sid	395.2 374.	136	374.8 328.	390, 358.4	5.63	379.6 335.2	6.8/2	:0	→ 00		1.8 352	378. 4.370.	5.6 363. 5.6 3333.	.140. 364.	<u>~</u>	308.1.386.		160.8 126	118.8 110	398.8 377.6	0.000	200 a 2000	205 5 265 6	376 4 34	3.1 880	
	11110,11		- -		_		3,10	20	30.00	25 00	376	1700	5 6	8 16	Ŧ	£ £	2000	2	46	=	68	× :	£ 1.	30	100	200	1
1.1	юрО	None	:	: :	S.i.h	Non	:	:	:	: :	:	:	: :	: :	:	: :	:	:	:	:	:		;	:	:	40	
<u>ن</u>	lor.	F.03	12:	? ? ! ! ! !	F. 15	E. 6	<u> </u>	F.05	F :	<u> </u>	F. 15	I-'. I	3	x 3.	F.06	10° -	- SO	10.7	F.3	F.04	E.0.	20.7	20.3	10.7	10.	F.03	
RANC	Color	Z.Z		-:-		7.7	 	N.	= :							7 7											1
APPEARANCE	Sedi- ment.	Cons'd	p. suo,	: :	:	: :	:	:	:	: :	:	:	: :	: :	:	: :	:	:		:	:		: :	:	:	Little	
	. y'd'm'l'	* *	* (.	* *	+	* -	- ₩	+	* -	+ +	-	*	* -	- +	*	÷ 4	F 4F	*	+	+	₩ •	<u></u>	+ +	-+	+	+ 1,	1
		100		= ?	क्ष	1-0	2 27	36	31 3	s. '9	£3;	98.		₩ 57 ₩ 17	177	- 3	2 0			5.	92		10		04	5. 5.	
DATEOP	- Exe	May	6 June	: :	:	5 Jul	:	:	7.112	: :		:	o Sept			30ct.	:		1.				Dec		:	:	
DA	1899 1899 Collec-Exami- tion, nation.	May 24	June (: :		17		. F.	112.	X 10	??	X;		2 00	96 .			- 51								· ·	
ON		5087 Mg	5162 Ju	0 00	5310	5355 July	5446	5496		50 %	- 02GG	. 01	57 SSTG	T &	. 91	00 00		() () () () ()		53 NO.	313	5	24 1166			333	
O.V.	[Black	ाँद रि	10	01 GI	500	10.10	516	T.C.	101	G 10	36	50	101	X X 01 0	5046	0000	(3(5))	6120	6196	6253	£ 3	6.36)	200	1140	0000	65/5	1

Colon M., Muddy, VM., Very Muddy, T., Turbid, C., Chondy, Color on Louring G., Gray, B., Brown, DB., Dark Brown, LB., Llight Brown, RB., Reddish Brown, BG., Brownish Gray, Bh., Brownish, R., Red, Black, GB., Gray-Brown. Terributy *Decided, §Very Decided, † Distinct, † Very Slight, | Slight,

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATTER ANALYSIS-PARTS PER MILLION.

-
7.
-
HA
_
FER,
62
-
\vdash
2
RIV
Q.
8
NOIS
-
ILLI
hand
0.0
TER
E
5
24
WA
OF
0
CE
P.
9

Report of ARTHER W. PALMER, T. J. BURRIEL, University of Illinois.

		* ***	
No. of Bac.	Cabic Centi- meter.	6.200 12.200 13.200 14.000 15.200 15.200 16.000	574.000
	Presen Abs. of] [[[[[[[[[[[[[[[[[[[1 +
sture Cent.	Temper of Air,		1
	Temper	######################################	;; <u> </u>
ht of	Helgl BW		4.0
TRO- GENAS	Nitrates	で 5 元 8 2 5 5 7 5 7 7 7 8 2 5 4 7 5 2 8 8 8 8 8 9 9 9 9 9 8 8 8 9 8 8 8 8 8	2.4 1.76
NITRO- GEN	Nitrites	<u>ਫ਼ੵ੶ਸ਼ਫ਼ੵਸ਼ਜ਼ੑਖ਼ਜ਼ਜ਼ਫ਼ਖ਼੶ਖ਼ਫ਼ਜ਼ਖ਼ਖ਼ਜ਼ਫ਼ੵਫ਼ਫ਼ਫ਼ਖ਼ਜ਼੶ਫ਼ੵ੶ਫ਼ੵਖ਼ਫ਼ਫ਼</u> ਫ਼	80.00
	Depued Sus-	1444860584888888458458888588854488888888	.32
ORGANIC NITROGEN.	Dis-solved.	### ### ### ### ### ### #### #########	105. 105.
ON	Total.	82548245835 8288445348888888	
AS .	p.pud -sng	<u> </u>	80.
AMMONIA.	p'vlos	4. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	.432
NITHOGEN AMMONIA	1.10'T	8 + 8 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	.96. .513.
	Free Am	+ + + + + + + + + + + + + + + + + + +	2.56
ED.	By Susper		3.6
OXYGEN CONSUMED.	By Dis-	@5335@65=3a==5=5=6=@5@r-@@65=r-r-@er-r-r-@6	# 30. # 30.
0 3	Total.		13.4
.aui	Срјог		36.5
Color	Igni- tion.		
· .	p,pud -sng	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	7 7
ATION.	Pisi(I	<u> </u>	30.8 30.8
APOR	Tot'l	σορεί το ει πο ο ποσεμπηροσμοροσιμο	31.3
RESIDUE ON EVAPORATION	bended Sus-	######################################	9.0 3.0
DUE (Dis- solved.		371.2
RESI	Total.	40 0 000000 0 0 0000000000000000000000	373.23
.10		пе	70 60
	obO		
YCE.	Color.	MANAMANANANANANANANANANANANANANANANANAN	0.
ARAZ			
APPEARANCE	Sedi- ment	Cons'd Co	: :
•	T.qun,L	#++*+***********	- F
OF	I899 Exami- nation.	Nay	: : - ::
DATEOF		$\frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}}{\sqrt{2}$	25 25 26
A	1899 Collection.	May May Jully Sept. Sept. Nov.	: :
ON I	Serial	4990 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900 4900	(5528 (5538

Colon-M., Muddy. VM., Very Mnddy. T., Turbid. C., Cloudy. Color on leavings G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown, RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. K., Red. Rd., Black. GB., Gray-Brown. Terremery -* Decided. & Very Decided. * Distinct. * Very Slight. | Slight.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO, SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER-SANGAMON RIVER, CHANDLERVILLE, ILL.

Report of Authen W. Palmer,	T. J. BURRILL,	University of Illinois.	

of ac.	per Cubic Centi- meter.	. :	3,700	10,550	9.900	19,700	37.250		2, (0)	10,500	45,500	4.9(0)	1.950	4.900	4.200	12,300	2,100	4.300	11.200	5,300	2.300	16,100	12,100
	Jo sq v	-		-																			
6 01.	Presenc	-			1	•		- '											+	7	0	-	1
9ann	Tempera				%; ? %; ?																		+
omn	Tempera	33	36.5	17	72.7	7 6	13:	5.26.	137	350	18	-1	6.		.c. &	10.	10	133	33	ж —	~	. 4	° ≥ − 0
	rigioH ofis7/				: -					-													
TRO-	Sitratics	-	0.5		? ; = ? ; =																		
NITRO GEN	Zitrites.				(E)																		
EN.	Sus-	.890	386	.63	595		368	4+	704	±.	??	.072	35.	355	==	¥9.	ğ.	91.	· ·	.03	30.		***
ORGANIC NITROGEN.	Dis-	1-	.763	.376	25.50	000	305	30F.	3063	90 71	.376	\$ 000 mg	\$5.55 \$5.55	355	. 153		30.5	36	.36	.296	306	2065	22
ON	Total.	.37	.03		Z. 9	200	.76	.856	-	.76	.696	o:	#3. **	.68	元	.395	£.	555	188	XXX.	38	.52	.76
	g p.pud	.32	.096.1	. 192	213	960	122.	960.	1561	G.	164	11.	960	-08T	.156	.036	.033	800	.011	.013	X(X).	1001	.000
SN AS					80.5 80.5					Ť	Ť												
NITROGEN AMMONIA.	T'10'T pourminoid b'vlos				1925																		
N. V.	monia.				.033								Ľ										
	Matter.				_																	0. 0.	es
GEN	solved.	9	_	∞ :	- DI		C.	_	7	₹:	33	1-	<u>{</u> -	5	9	٠,	_	6	25	iG	0 6	C 22	50
OXYGEN	Total.	9	ıG.	1-	. 6. 	-	23	G;	10	00	200	_	1-	œ.	-	10	ıC	_	<u>c:</u>	oc.	S:	33	5.7
	Chlori			.c	රා ර 1 - ජ	15	\$1	x	_	100	~ ~	x				oc	~	::		ıc	35		.
			_		1C 1C		_							_		=						9	1-
11	on Igni- tion.				ei≃ ∞o			_															
O.N.	F sns	1-	0	=	00 00	0 00	2.28	2 10.8	2,21.	3.0	6'16.8	× ×	1 15.0	30		<u>∞</u>	36.	આ	~	1-	9		-
RATI	Foss on Base of Total	47.2 39.6	6.31.	£ 30	55.65.46.30 35.00.46.30	439 2	62. 33.2.28.	19	8,37.	.∵ .∵	1	200	31	1	13.2 12.	x ====================================	(元) (元)	x 17x	6,16.	35.	2.04	26.8 26.	=
VAPC	1.10.L	() ()	00 i	9.						28				-91		253	30	9 30.	61 19	× + ×	8 31.	3 26.	8 46
RESIDUE ON EVAPORATION	popuod sus-	217.	16.9	201	150.4	39.68	163.6	108	100.8	5.	104.8	0.0	E.	50	61.6	26.	30.3	35	43.6	16.8	10.8	_	30
DUE	Dis- beyled,		: · ·		9 - 9 : 5 6 : 5 6 : 5 6 : 5	3	248.1	270.2	285.6	280.3	62.	206.1	262.4	215.6	360.8	303.6	X 20	.86.	.00	307.6	×9.0%	140.x	276 1
RESI	Thotal.	398. 1	25	00 00 00	140. 200 n. u		17.		-	231. 2	~		-		37.		323.6 302.	323 0 208	333.6 200	321.43	300.4.5	3000	33
	(5,5())			0	→ ~	- 50		దేస	**	å:	<u>್</u>	হা	**	či	**	**	**	66	**	30	~~	Šì.	55
	Odor.		:	:		:	;	;	3	3	:	:	*	*	*	:	9 9	:	•	: 	3	9	:
E	NCE.				E ×			F.04	_		_	_	F.03	F.03	F.05	F.03	F. 03	F.03	E.03	F. 03	F. 63	F.02	F.03
RANC	00	N.	7.	7.		. 7	7	. I.	M.	7.	7.	7.	7.	7.	-	7.	7.	N.	7.	7.		7.	N.
APPEARANCE	Sedi- ment.	Much	Cons.d		. ,	• 9	9 9	9.9	- a - g	. ,	9.9	Much	p.suo.)	Little	p.suo.	ittle	:	7 7	p,suo,		ittle	3.7	99
17.	Z Z S		+ Co	33	沙 李	*	-	*	\$}.	+	25	I. *		<u>.</u>	0.0	1	de.	+	0.) +	+	T		+
		June 1	£	?? ??	9 :-	-	10	17		-	27	0.5		6.	· · · ·	??	C:	91	200	88	C)C	ic	27
DATEOF	DATE OF 1899 1899 1901. Collect Exami-			9 1	int.	7.11	;	:	12.7.	:	:	:	Oct.	94	79	70%	99	9.9	9 9	:	Dec.	9.	9
DAT	1899 olire- tion.	65	ne 7	?}	1		6.	16	3.1		200	5			77	1. 1	30	15	21	Ŷi	P. 65	1:3	20
		5131 May	111 C 9	5.	o July	20 A 1155		: 0		こうんる			M Oct	:		10% S	. 63	., 9	: X.	()	J-0 1)	:	*
No.	Serfal	513	G	50	0 12 Z	5555	DEPT.	5649	5770	5 4.5	5911	5:7:3	CANA.	61(5)	616	6213	5555	6316	6255X	6450	6170	6521	6561

Conon -M., Muddy, VM., Very Muddy, T., Turbid, C., Cloudy, TURBIDITY -* Decided. § Very Decided. † Distinct. ‡ Very Slight. Slight.

Color on Janifion—G., Gray. B., Brown. DB., Dark Brown. Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

SOURCE OF WATER-HILINOIS RIVER, BEARDSTOWN, JLL.

No. of Isac.	per Cubic Centi- meter.		1.100	0,50) (S	1.050	. 350 100 100 100 100 100 100 100 100 100 1	2005.00	.£ €	9.60	3.960	6965	0 0 0 0 0 1 1	050.5	3,300	9,750	4.000	6.500	90012	000.75	95.288 60.000	00.03.0	194,000	181 500	48.400
6 of .	Presence Abs. of	:					:		+			1		F ;	1	+	+-		+-	+	: .	-	ł i		:
11116 'ent.	Tempera of Alr, C	57.5		26.5	33.		· ·		·	37.	??	<u></u>		9	200	+	<u>∞</u> :	C :	:	2 2	· ·		: : :	~	= -
	Fempera	:	 	16.5			رة الا			30.	10.	5.5	- 12 - 12 - 12 - 12 - 12 - 12 - 12 - 12	· ·	16.	18.	.61	:i o	ć g	:i :	. v	. 10	?		0.
.1.	ИзіэН э≀в <i>∏</i>	6.01	x +	-0	0.00	8.9	ئ. ئ. ب	. ec	6.3	6.	6.3	♠ 1	- - 4	m 🚓)	6.3	9.9	6.5	9.0	0.0	0.1	2.0	0.0	+ α 5 ω	1	1-
TRO- GEN AS	Nitrates.	ej 3	: : : : : : : : : : : : : : : : : : :	£ :	2	08	3	3.5	9	.56	?}	7	÷ «		.96	8.	- .	26.	00	() = () =	2 =) oc	8	1.18
NITRO- GEN	Nitrites.	.065	3.99	<u></u>	??	33.5	٠ <u>;</u> ٥	<u> </u>	<u> </u>	<u>=</u> :	<u>.</u> 05	.6 3	080	3.8	0.	80.	: 13	٥١	- ?	? ?	5 6	300	100	.09	.07
o Z	-sns-	43	0000	368	90.	448	x 2	43.50	.256	272	448	.192	30.	288	.481	84.	. 5 8 8 8 8 8 8 8 8	2000	000	5.6	9.0	25.7	656	464	92:
ORGANIC NITROGEN	Dis- solved.	98:	132	712	.76	7.65	θ. (6.9.	6.00	.664	849	110	808:	.+4 506	.676	181	.52	.616	07.	100	010.	618	2 1	200	776	986
Q.N.	Total.	65.	15:	80.	.33	77	88	80	8	56.	. 16		. 0	96:	896		90.8	.032	5000	000	920	000	200	67	. 196
	E p,pud	.096	194	91.	128	.032	2000 2000 2000 2000 2000 2000 2000 200	98	212	.064	324	. 116	× = = = = = = = = = = = = = = = = = = =	176	. H3	.113	<u></u>	080.	+00.	000	? :	~110	336	176	.11:
EN AS	biodi b'vlos	288	38	356	200	88. 33.	-98: -98:	# 88	504	.256	<u>.19</u>	368	212	:0: :::	35	288	304	950	700	100.	3 8	886	000	368	304
NITROGEN AMMONIA	T'10T 1'10T sign	384	.768 .768	350	448		35.5	388	.416	.33	.516	.384	25.55	448	.432		416	.432	0.00	000	130	80.6	808	110	.416
Z	Free Am-	148	199	35.5		352	22 E	118	416	8	200	∞ <u>.</u>	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	133	176	ж Э	880	2) 0	9.5	7 3	5.5	11	-	89	.61
- A. G.	By Suspd Matter.	1.9	- - - - - -	\ \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{	· 1~	œ :	· 	3 63	→: :::::::::::::::::::::::::::::::::::	6.	<u>ن</u>	ر ص ص) OC	÷;	<u>x</u>		016	- - -	- œ	ो ल ० १०	-	9.6	2.9	1.5 3
OXYGEN	By Dis- solved.	6.03	2.0	∞ ∞ 3; 0	. x.	11.7	00 c	00 n oc	-	€. 50	 	§ § §	ۍ ∞ د د د	6.00	ا .	7.1		10	00	1-1-	+ 1·	1	7.3	9.3	8.
Con	Total.	5.3	0 1-	51 E	12.5	16.	25.5		9.5	?! ∞	0.0	ος s	၁ တ ဘ ∞	17.	9.5	3) 3)	0000	000	0 10 0 0	000	1 7	0	6.6	2.5	11.3
.əu	Chlori		? :	16.	32 35			3 23		<u></u>		7.	25.	10.	5.	<u>ः</u>	∞ ∞ <	.16	5 2	16	200	200		22	
olor	on Igni- tion.			~ ~		j.			-:		,														
	= p,pud	9:		es 00	30.00		. o	41.638.8	6.	9.	T.		0 -	_									3.6		7
AT10.	Por Property of Pr	99.4	. T.		8: #		11 7 25 7	00. 00. 00.	33.6	0.0		<u>.</u>			7.3	15.6		. 0)	9			36.4		8.4
APOIL	LosT 1'10'T	13.00	53.652.4	50°.	50.0	₹. %	25.0	41.6	49.6	59.8 45.9	1.00	36.	30 0 30 8	34.	18.8	00	19.6	7.0	000 0000 0000 0000 0000	0.00	19 6	7	10.	288	45.3
RESIDUE ON EVAPORATION.	Sus- pended.	9:0	- - -	oi 5	7:82	58.	20.30 100.00	130	61.2	⊕ ₹	30.6												37.6		
CE 0	solved.	162.8 275.2 18	0 + .	73 73 73 73				_		00		-)	. 00	00		٠ <u>٠</u>	<u>ن</u>	. •	: 0	; x		300		ं क	7.
RESID	Dis	000	§ <u>≈</u>	421.2 309.2 388 4 399	129.2 340	100 cm	0000	309.2 186.8	337.2 276	334	349.6 320		335.6304	362.8 336	374.8 312.4	364.4313	381.×3501	3	୍ଟ	373 9 34	36		357.6320	347	7
	Total.	e 463	5.5	주 주 중 중 중 중 7	₹	350	305		33,	346.	345		335.6	36.	37.	36-	20 10	190	302	520	280	38	357	370	36
•:	юрО	None	:	::	:	3 :	: :	:	:	:	:		:	:	:	:	: :	;	:	10	:	:	:	• •	:
Pi Pi	Color.	F.05	F.15	F.06	F.06	F. 6	7 E	F. 15	F.06	80.E	0.1	F.06	2 CO	F.03	五 3 3 3 3 3	_ ;	5 8 5 8	30.6	10	F. 03	E (S)	F.04	F.03	F.01	F.04
RANC	O)			ZZ		<u> </u>	<u>.</u> z	Z	M.	Z:		:		Z .	 	:					7		7.		
APPEARANCE	Sedi- ment.	Cons'd		Cons'd	:	;	Wuen ()	S S S S S S S S S S S S S S S S S S S	Much	Cons.d	: :	Man	Consid	:	:	:	: :	:	:	:	:	:	:	• ,	:
	Turb'y.	* 9%		* *	*	* 4	+ *	÷	* 9~	÷ ·	÷ ;	# # C ?	* 66	. 9	*	+ 02	* +	+	*	* 10	+	+	* 91	£33 ÷	+ 0%
OF	1899 Examination.	21 May	: : : : : : : : : : : : : : : : : : :	luly:	:			:	** **		:	: :	**	***	:					. 9.)ec.	:	*		**
DATEOF	1899 1899 Collec-Examition.	<u> </u>	19	ly 6	ි බ	37.0); 5 00 5 00		45		1: 1		7 %		≘	61	٠.		15:	0 00		1-		- -	
,		May May	:	July.	;	:	2	: 9	9		30 Se 1	: :	-10	5 Oct	: 9	: :	: 2	-	:	:	:	8 Dec		;	:
No.	Serial	5098	524	5354	5460	5516	5612	59996	5716	5769	28.50	5073	5975	6015	6066	6121	618 635	6961	6335	6373	6132	6488	6531	6573	6597

T. Turbid. C. Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Black. GB., Gray-Brown. Color-M., Muddy. VM., Very Muddy. Trunnury * Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

STREAMS ENAMINATION-SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-FARTS PER MILLION

SOURCE OF WATER--ILLINOIS RIVER, KAMPSVILLE, ILL.

Report of Arthur W. Farmer, T. J. Burnell. University of Himode.

	and Salat And	
No. C. Bac. Per Cribic Cent.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	TOTOKE
Presence or Ala.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Pemperature of Air, Cent.	သင့် ရှိကို နှစ်မှုရေးကို နှစ်မှုရေးကို မြောင်းမှုရှိသည်။ မြောင်းမှုရှိသည်။ မြောင်းမှုရှိသည်။	20
Temperature of Water, C.	19.	27
Height of Water.	\mathbb{R}^{2}	16.
Sitrates S	[中文] 後次二百日日報日二十日本文日第第四名中文建設 [卷] 日報日本	5
NITE OFFICE SERVING OFFICE NAME OF THE OFFICE NAME	हिस-इन्हें इड्डिंग-इन-इन-नियं च इच इड्डिंग किन महि	9.
bended, % ~	8-66×66×492556842884425528	0++0
Organic Nitritoden Nitritoden Sus- Sus- Sus-	# # # # # # # # # # # # # # # # # # #	, C.C.
Total.	न्त्रहेत्वत्त्रत्ते प्रथमित्रहेत्व द्वारा स्थापन	
i p.pud ≈ns	88 8 4 4 8 8 1 8 8 8 8 8 8 8 8 8 8 8 8 8	5
A A S	***************************************	
TTBOGEN AS AMMONIA. Albuminoid T'01'l Dissolv'd solv'd	**************************************	c
Free Am-	854878787878787878888888888888888888888	210
Harlor,	$\frac{1}{1} + x + x + y + y + y + y + y + y + y + y$	
Solved Signature of Solved	一	-
Total. Co.N.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
(apprime,	- 1	0.4
olor on geni- ion.	長年民族は民民民民民民政治(治) 日本民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民民	
2 E p,pud sns	# n x & & & o i i i i i i i i i i i i i i i i	
POT TOOP TOO TOO TOO TOO TOO TOO TOO TOO	88888888888888888888888888888888888888	
Torr Loss	12	.
RESIDUE ON EVAPORATION SOLVED TOTAL	高いまという。一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、一般には、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のいは、のい	
beyles	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$,
S S S S S S S S S S S S S S S S S S S	$\infty + + \times + + \times + \times + \times + \times \times \times \times \times \times \times \times $	c.
Total distort		260
.robO	None	
E. lor.	######################################	1.7.
RANCE.	**********************	
APPEARANCE. Sedi-	Much Ons'd Consider C	
Y. qana,	S	
	warting a war a wa	7
PATE OF 18 18 16 16 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	No. 1 = 1 = 1 = 1 = 1 = 1 = 1 = 1 = 1 = 1	
DATE OF 1899 Collec. Examition.	Nay	2
Serial No.		2015
	?要本格的的原则是我的证明的证明的证明的证明的证明的证明的证明的证明。	3 1

Dury *Decided. \$Very Decided. + Distinct. ‡ Very Slight. Slight.
Colon on Jonation. G., Gray. B., Brown. DB., Dark Brown. BB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red., Colon on Jonation. G., Gray. Bh., Reddish. W., White. Gh., Graylsh. Blk., Black. GB., Gray. Brown. Terringery *Decided. & Very Decided. * Olstluct. \$ Very Slight. Slight.

TABLE 21.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON, ILL.

ARTHUR W. PALMER,	T. J. BURRILL,	University of Chicago
of		
Report of		

of C.	r vic ti- er.	28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
No.	per Cubic Centi- meter.	ক ক
	Presence Abs. of	1
	Tempers of Air, C	***************************************
	Tempera of Wate	
10.1	Heigh Mate	:::FF554 : FE00r-0r-0400444 : www.en.4444
TRO- GEN AS	Nitrates.	65.888.88888888888888888888888888888888
NITRO- GEN	Zitrites.	86.86.96.96.96.96.96.96.96.96.96.96.96.96.96
ic sn.	-sus bended.	88.5
ORGANIC NITROGEN	Dis- solved.	E. F.
OI	Total	
	i p,pud	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
EN AS	Dio ii	<u> </u>
NITROGEN	Albuminoid Discolv'd	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
N N	Free Am-	85.5.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.
N ED.	By Suspd	6.41-40x7-2001-1-103 2001-101-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1
CATGEN	By Dls-	$\begin{matrix} F_{-}^{-} \alpha = \alpha_F + \alpha_F \alpha \alpha \alpha \alpha_F + \alpha_F \alpha \alpha \alpha_F + $
Cop	Total.	<u> </u>
.911	Chlori	\$
Jolor	on 1gni- tion.	ಪ್ರಹಾಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ
	g p,pud	ရေသာသာသည်တွင် ႏုိင်သိသောလည်လို့ ရေးသ လာသာည်မှုမှုတွေများက လာမှာ လောသာသာသမာတ်တွင်းတွေ ႏုိင်းသောလာသည် တန်ဘာသာသည် တစ်ပေးသည်သည် မြောင်းသော လောသာသာသည်တွင် ႏုိင်းသောလာသည် လောန်သာသာသည် တစ်ပေးသည်သည် မောင်းသည်
ATTO?	solv'd b'vlos	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
APOR	Torr	は
RESIDUE ON EVAPORATION	bended.	888 48 48 48 48 48 48 48 48 48 48 48 48
UE O	solved,	
RESID	-siQ	
J.4.	Total.	
:	торО	N
	or.	84444444444444444444444444444444444444
TANCE	Color	ZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ
APPEARANCE	Sedi- ment.	Cons'd Cons'd Cons'd Little Cons'd
	Turb'y.	でごうもおしゅう2000 あこりと 20 1 2 2 1 1 2 2 1 2 3 1 2 3 2 2 2 3 2 1 2 3 2 2 3 2 1 3 2 2 3 2 1 3 2 2 3 2 1 3 2 2 3 2 1 3 2 3 2
OF	1899 1899 Collec- Examition.	Nay Aligh Sept. 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
DATEOF	1899 Collection.	
		48 28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
No.	Serial	20000000000000000000000000000000000000

TURBIDITY -* Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

COLOR ON IGNITION—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Gray-Brown.

Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy.

TABLE 22.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON, ILL.

No. of 13ac.	per Cubic Centi: meter.	(2000) (2	7.300
	Presence Abs. of	· · · · · · · · · · · · · · · · · · ·	+
	Tempera of Air, C		9
	Tempera of Wate	:::::::::::::::::::::::::::::::::::::::	
JO 1	ияјен - Иејки	□ : 「ころらユ: 「子辺のたったのよいの444 」以びいじゅつでは44○ : 「じードー : つがたるーカーがは44%」 「近に44が、 ないが	***
TRO.	Sofrativ		91.
NITRO- GEN	Withtes	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	
o X	Sus-	8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.	.301
ORGANIC NITROGEN.	Dis.	4±466666666666666666666666666666666666	396
ON	Total.	<u> </u>	9.
	gi p,pud	## 1	.113
EN AS	Albuminoid Pistori'd b'vlos	<u>88888948869984455558558</u>	303
NITROGEN AMMONIA	I'10'T	84418882544755585888888888888888888888888888	£.
Z	Етее Ли-	2 8 8 2 8 2 9 2 9 8 8 9 9 9 9 8 9 9 9 9	230
N ED.	By Suspd	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$.;
OXYGEN	By Dis- solved.	$\frac{\infty \circ \overline{a} \overline{a} \overline{c} \overline{a} \overline{c} \overline{c} \overline{a} \overline$	G:
000	Total.	$8 \mp 88488927745677627557666888888888888888888888888888$	- -: -:
.en	СЫЗогія		3.6
Color	on Igni- tion.		<u>~</u>
ż.	z p, pud snS	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3.6
SATIO	sici b'vios b'vios b'vios sici		9.61
APOI	1.10.L	े हैं हैं के की की कि की	÷;
RESIDUE ON EVAPORATION	bended.	88488888888888888888888888888888888888	÷ ;
DUE (Dis- solved.	μοροκοριμιτικουμι οκ κα κοκορτ ομκα κ	2 }
RESI	Total.	ုပ္ ကြာလုံး 'မႈလုံထုံကုံ 'မုံသွဲတာ ' မေးမေးမြေးများဆိုသည် သို့ ဆိုများများများဆိုသည်။	
	1		=
•	robO	None	7
E E	Color,	######################################	
RANG		ZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ	7.
APPEARANCE.	Sedi- ment,	Much Consid Little foos d Little foos d Little foos d	:
	*X,Q4U,L	* * * * * * * * * * * * * * * * * * *	Ŧ
SOF	1899 Exami- nation.	Na	
DATEOF	1899 Collec- tion.	######################################	<u>0</u>
	Co.l		:
No.	Serial	20	+15/67

COLOR M., Muddy: VM., Very Muddy. T., Turbid. C., Cloudy. Color ox lexition G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown. TURBIDITY -* Decided, & Very Decided, + Distinct, + Very Slight, Slight.

TABLE 23.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER - MISSISSIPPI RIVER, 100 FEET FROM ILLINOIS SHORE, ALTON, ILL.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois,

No. of Bac.	per Cubic ('enti- meter.	2.8000 2.8000
	Presence Abs. of	
ture ture	Tempera of Air, C	
ture r, C.	rempera	- : : : : : : : : : : : : : : : : : : :
10 1	dgiəll əlaW	
TRO- GENAS	Nitrates.	でお客がい。となびばまままがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるといりないがあるというないがあるというないがあるというないがあるというないがあるというないがあるというないがあるといりないがあるというないがあるというないがあるというないがあるといりないがあるといりないがあるといりないがあるといりないがあるといりないがあるといりないがあるといりないがあるといりないがあるといがのののののののののののののののののののののののののののののののののののの
NITRO- GEN	Nitrites.	64.000 64.000 65.0000 65.000 65.000 65.000 65.000 65.000 65.000 65.000 65.00
o i	Sus-	2146.828.825.828.825.888.888
ORGANIC NITROGEN.	Dis-	\$\frac{1}{2}\frac{1}{2
Or	Total.	<u> </u>
	i p.pud	8
N AS		884844486544885888865558855844488488488
NITROGEN AMMONIA	Albuminoid Piori Divios	5-1-1-2-8-8-8-1-1-1-1-1-1-1-1-1-1-1-1-1-1
NEA	minom	0.00
- i	Matter.	φφε-Παφαμμαση αροσημμη οι - φμω-τόα ο
OXYGEN	By Dis- solved.	00000000000000000000000000000000000000
CONS	Total.	5 7
ne.	('hlori	$\frac{4+\pi}{12}\frac{1}{12}$
Color	on geni- cion.	
	l la mod	:
TION.	b'vlos h'vlos h'baq	883388383888388889988888888989888888888
PORA	Tor'l Loss	23.54.54.55.55.55.55.55.55.55.55.55.55.55.
RESIDUE ON EVAPORATION	pended.	क 'मधामधामळधाष्ठ क्मधा क्षां क्षां क्षां मक्ष्य मुक्क्रक्षिक
E ON	solved.	@ @ @ @ @ @ @ @ @ @ @ # # @ @ # @ @ @ @
ESIDU	-sid	2005 2005
<u> </u>	Total.	93.000
*.a	юрО	None
	į.	- 19
ANCE.	Color	- SANANANANANANANANANANANANANANANANANANAN
APPEARANCE	ii.ii	u p q c p q q c p q c p q c p q c p q c p q c p q c p q c p q c p q c p q c p q q c p q q c p q q q q
API	Sedi- ment.	Much Cons'd Mich Cons'd Con'd Con'
	Turb'y.	*
OF	1899 Examination.	May May Auno Aug Coet. Nov. Dec.
DATEOF	ė,	$\frac{2}{2} \frac{1}{2} \frac{1}$
	1899 Collection.	No.
.o.X	Serial	5000 5000

T., Turbid. C., Cloudy. Color on lention-G., Gray. B., Brown, DB., Dark Brown, LB., Light Brown, RB., Reddish Brown, RG., Brownish Gray. Bh., Brownish, R., Red. Reddish, W., White, Gh., Grayish, Blk., Black, GB., Gray-Brown. Colon -M. Muddy. VM., Very Muddy. Turbinger *Decided. §Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 24.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM ILLINOIS SHORE, ALTON, ILL.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

o. of	per Cubic Centi- meter.	2000 2000	62,400
(!oll.	Presence Abs. of	<u>: : : : : : : : : : : : : : : : : : : </u>	+
Juo?	Tempera of Air, C	20,20,20,20,20,20,20,20,20,20,20,20,20,2	
9r, C.	Tempera	: : : : : : : : : : : : : : : : : : :	0.
1.10	dgl9H otsW	$\frac{(i-\delta)(i-4)(i)(i-5)}{(i-6)(i-4)(i)(i-6)} = (i-6)(i-6)(i-6)(i-6)(i-6)(i-6)(i-6)(i-6)$.: :
PRO- BEN AS	Nitrates.	で	
NITRO- GEN	Nitrattes.	90.000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
IC EN.	sus- sus-	464444668848884864864868686868686868686	
ORGANIC NITROGEN.	Dis- solved.	644 9 9 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	
No.	Total.	200 200 200 200 200 200 200 200 200 200	1.08
So	g p,pud	1	.972
	l'10T pionimino b'vios	884448888448884488848888888888888888888	208
NITROGEN	1'10T	5-4-8-5-8-4-4-8-4-8-4-8-4-8-4-8-4-8-4-8-	<u>×</u>
Z	Free Am- monia.	885-00000000000000000000000000000000000	.059
ED.	By Suspd	<u>できるいでは、できない。 このい - これキー・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・</u>	-
OXYGEN	By Dis- solved.	© ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○ ○	10.7
000	Total.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	11.0
<u> </u>	СПОП	<u> </u>	
Colo	on Igni- tion.		
O.X.	i p,pud sns	90000000000000000000000000000000000000	
RATIO	्र होत	3.5%%5.7%%2.15.5%%2.8%%2.8%%2.8%%2.9%%2.9%%2.9%%2.9%%2.9	5 30°
VAPO	1.10T		6§
RESIDUE ON EVAPORATION	Sus- pended.	<u>\$68886568888888888888888888888888888888</u>	٠ ٠
IDUE	Dis. borlos		196.
RES	Total.	25	∞. 1 554 .∞
	торО	N. 00:::::::::::::::::::::::::::::::::::	;
l si	or.		F.0.3
RANCE	Color.	ZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ	N.
APPEARANCE	Sedi- ment.	Much William W	1,
	.gang,	********************	+-
Eu.	1899 1899 Collec- Exami- tion. nation.	Mar. : : : : : : : : : : : : : : : : : : :	• 1
DATEOF	c Es	$\frac{1}{2} \omega = 742 \times \pi \pi 33 \times \pi \pi \pi 22 \times \pi \pi \pi 22 \times \pi \pi 23 \times \pi \pi 23 \times \pi \pi 23 \times \pi \pi 23 \times \pi 23 \times$	·
D	1899 Collection.		:
No.	Serial		6554

COLOR-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition -G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown. Trubinity * Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 25.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

Report of ARTHUR W. PALMER, T.J. BURBILL, University of Illinois,

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, ALTON, ILL.

No. of Bac.	per Cubic Centi- meter.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	Presence Abs. of c	<u> </u>
ent.	Tempera of Air, C	
	rempera etsW lo	$ \begin{array}{c} \vdots \vdots \overline{\otimes} \overline{\cup} \overline{\otimes} \overline{\otimes} \overline{\otimes} \overline{\otimes} \overline{\otimes} \overline{\otimes} \overline{\otimes} \otimes$
10.1	Heigh otsW	::: ಸಾವರ್ಷದರ್ಪರವರ್ಷ-೯- ಕಟಲುಬಹುಕುಲುಲು – ಲುಟ್ಟಾಕುಬಟಟಟ
TRO- GEN AS	Zitrates.	# 55 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
NITEO- GEN	Nitrites.	20000000000000000000000000000000000000
IC SN.	-sns bended.	
ORGANIC NITROGEN	Dis.	**************************************
ON	Total.	\(\text{\tilde{R}}\) \(\
	g p.pud	25
EN AS	Tor'l	<u> </u>
NITROGEN AMMONIA	1.10.L	25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
N A	Free Am-	00 00 00 00 00 00 00 00 00 00 00 00 00
á	By Susper	န္းနည္းနည္း က ကိုလည္တည္တည္းႏိုင္ငံတည္းကို အတည္တည္တည္တည္က မတ္တည္တည္တည္က မတ္သည္က မတ္သည့္အသည္က မတ္သည္က မတ္သည္က မတ္သည္က မတ္သည္က မတ္သည့္အသည္က မတ္သည့္အသည္က မတ္သည္က မတ္သည္က မတ္သည္က မတ္သည့္အသည္က မတ္သည့္အသည္က မတ္သည့္အသည့္အသည္က မတ္သည့္အသည့္အသည့္အသည္က မတ္သည္က မတ္သည့္အသည့္အသည့္အသည့္အသည့္အသည္က မတ္သည့္အသည့္အသည့္အသည့္အသည့္အသည့္အသည့္အသည့္
OXYGEN CONSUMED.	By Dis- solved.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Con	Total.	$\frac{85498222277535758575825774}{48884577627777777777777777777777777777777777$
ne.	Chlori	$\frac{1}{2}$ $\frac{1}$
olor	on gni- ion.	
	p,pud	နိုင္မရိုင္သီးလိုတ္တရာတ္လည္တရာမွာ လုပ္သည္က မရိုက္သည္တရာမွာ လုပ္သည္က မရိုက္သည္က မရိုက္သည္က မရိုက္သည္က မရိုက္သည္ လုပ္သည္လလိုင္း သည္သည္မရိုက္သည္က မရိုက္သည္က မရိုက္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့
TION	e sid	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
APOR	Tot'l	88
RESIDUE ON EVAPORATION	pended.	838.888.888.888.888.888.888.888.888.888
TE 03	solved.	
ESID	-siCl	
	Total.	
•	торО	O
-:	or.	######################################
SANCE	Color.	RENERGE EN
APPEARANCE	Sedi- ment.	Much Consider Much Sons de Consider Con
ΑF	Turb'y.	
		######################################
DATEOF	1899 Exami- nation.	May
DAT	1899 Collection.	
.0.7		\$50.8 May
.oV	Serial	4998 1004

Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. Ch., Grayish. Blk., Black. GB., Gray-Brown. Turning + Very Steided. § Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 26.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER.
T. J. BURRILL.
University of Illinois.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM MISSOURI SHORE, ALTON, ILL.

No. of	Cubic Centi- meter.	2	22.000
	Presence 10 .sdA.	<u> </u>	+-
этия Эпэў.	Tempera of Air, C	古일곱임막왥띰뫶띋똮쁅뫇뙁왥왞뭑텧qaqqel	ο.
r, C.	eroqmə'l' əts7/ lo	: <u> </u>	0.
101	figioff ota?//	::::=================================	66
TRO- GEN AS	Sitrates.	1 4 4 8 8 6 6 8 4 4 6 6 8 4 5 4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	.16
NITRO- GEN	Nitrites.	900000000000000000000000000000000000000	
S.S.	Sus- Sus-	200	.461
ORGANIC NITROGEN.	Dis- solved,	4.6.6.6.4.8.4.6.6.6.6.4.4.8.4.8.4.8.4.8.	99:1-
Q.N.	Total.	\$87555555555555555555555555555555555555	.o.
	E p,pud	\$\frac{1}{2}\frac{1}\frac{1}{2}\f	91.
EN AS	Albuminoid Pistory Distriction of Pistory Dis	8448654448844455554255555544554445	921.
NITROGEN AMMONIA	Torr	825.37.57.57.57.54.48.48.48.48.49.48.48.48.48.48.48.48.48.48.48.48.48.48.	.336
\vec{z}	Гчее Ань- пюніа,		.035
ED.	By Suspd	<u> </u>	i.e.
OXYGEN CONSUMED.	By Dis- solved,	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Ţ. =
Co	Total.	$\frac{r-c\delta^2 R}{c^2} \approx \frac{8}{3} \approx \frac{8}{2} \approx \frac{8}{2} \approx \frac{8}{2} \approx \frac{1}{2} \approx \frac{1}{$	6:]
.ən	Chlori		- · · · ·
Color	on Igni- tion.	我我我我我我就就就让你你就是你你你你你你你你你你你你	~ <u>`</u>
	i pipud	$p_1p_2p_3p_3p_3p_3p_3p_3p_3p_3p_3p_3p_3p_3p_3p$	5
ATIO	n sol l'10'l' b'ylos b'ylos -sid -sid -sid	######################################	
APOF	l'io'l	######################################	
RESIDUE ON EVAPORATION.	hengeg.	$\frac{4\pi^2}{600} \frac{1}{100} \frac$	2.65
ore o	.bevled.	第23年8年3月8日8日8日8日8日8日8日8日8日8日8日8日8日8日8日8日8日8日8日	;
RESII	·si(1	e_1 $= \frac{1}{2}$	2.3
	Total.		<u>e</u>
".	10bO	None	
ei	Color.	and the contract of the contra	F.0.
RANC	S	*************	Z.
APPEARANCE	Sedi- ment.	Much Much Much Much Much Much Much Much	
	Turb's.	######################################	-
OP	1899 Exami- nation.		:
DATEOF		교급수원은 소리 등 교육 등 교	୍ର ଶ
	1899 Collec- tion.		* *
No.	Ishros	######################################	6556

Colon-M., Muddy, VM., Very Muddy, T., Turbid, C. Cloudy, TURRIDITY *Decided. \$Very Decided. + Distinct. ‡ Very Slight. | Slight. | Colon | M., Minddy, V.M., Very Minddy, T., Turbid. C. Colon on Learnest G., Gray, B., Brown, DB., Dark Brown, L.B., Light Brown, RB., Reddish Brown, BG., Brownish Gray, Bh., Brownish, R., Red, Red, Gray, Brown.

TABLE 27.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM MISSOURI SHORE, ALTON, ILL.

Report of Arthur W. Palmer, T. J. Burnell. University of Illinois.

No. of Isac.	per Cubic Centi- meter.	18.8.90 19.00	9,600
	Presence Abs. of	:::::::::::::::::::::::::::::::::::::::	++
ture ent.	Tempera of Air, C		0.
	Tempera of Wate	, ::: <u>%</u>	÷ 0
1.1	dgiəll ətsW	: : : : : : : : : : : : : : : : : : :	က်ကံ
TRO- GEN AS	Nitrates.		
NITRO- GEN	Nitrites.	8 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
IC EN.	Sus- pended.		
ORGANIC	Dis- solved.	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	84. 42. 43.
ON	Total.	88-14-68 928-24-893-895-895-895-895-895-895-895-895-895-895	 22
S.	E p,pud	8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	.192
EN AS	Albuminoid l'10T Pist Divios Divios Pist Pist	4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	.192
NITROGEN	Tot'l	### ### ### ### #### #################	368
Z	Free Am-	200 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	030
N ED.	By Suspd	<u> </u>	2.9 2.9
OXYGEN	By Dis-		2=
000	Total.	<u> </u>	= =
	Chlori	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	4.6
Color	on Igni- tion.	anananananananananananan hannanananan	
Ż.	i p,pud	06.98678868888888888888888888888888888888	
RATIO	Loss on Ig. Dis- Dis- Dis- Dis- Dis-	0.50 86 12 86 12 86 86 86 86 86 86 86 86 86 86 86 86 86	30.4
VAPO	1'1oT	888686448588844864 8446448564865888448864 8446448664864884 8446448664864884 8446448664864 84464864864 84464864864 84464864864 84464864864 84464864	31.6
RESIDUE ON EVAPORATION.	bended.	444-458-888-888-88-98-98-88-88-88-88-88-88-88-	14.8
DUE	Dis- solved,	888 47 86 688 63 4 2 688 4 688 8 6 6 6 6 7 4 8 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	53.2
RESI	Total.		81.6 168.1
	торО	N	
			9.78
NCE.	Color.	MNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	M. E.
APPEARANCE	.i. ;;	•	
АРР	Sedi- ment.	Much Cons'd Much Nuch Cons'd Cons'd Cons'd	: :
	Turb'y.		+
OF	1899 Exami- nation.	28 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	; ;
DATE OF	1899 Collection.		200
	Col	May 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
No.	Serial	4996 990 900 900 900 900 900 900	6557

TURBIDITY-* Decided. § Very Decided. + Distinct. ‡ Very Slight. | Slight.

Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Grayish. Blk., Black. GB., Gray-Brown.

Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy.

TABLE 28.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 400 FEET PROM ILLINOIS SHORE, AT CHAIN OF ROCKS. PUNITING STATION, ST. LOUIS WATER WORKS.

Report of ARTHUR W. PALMER, T. J. BURRILL. University of Illinois.

No. of Bac.	per Cubic Centi- meter.	6 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	16,000
	Presenc lo sdr		+
	Tempera of Air, C		5.
er.	ngion ots7/ sroqmoT ots7/ lo	: 833220093320577777777777777777777777777777777777	
NITRO- GENAS	Sitraties.	\$\frac{2}{2}\frac{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac	
N.	pended,	25	
ORGANIC NITROGEN,	solved.	E 1585 F F F F F F F F S S F F S S S S S S S	_
ORC	Total.	488784120446812044681204 488784120446812044681204 48878412044681204 488784104 4887841204 488784104 488784104 488784104 488784104 488784104 488784104 48	
oc.	Sus-sus	88.25.25.25.25.25.25.25.25.25.25.25.25.25.	
▼ ;	T'10T Bis- bivlos	古智慧民群組長行為古聖聖古吉王五王聖詩家民華民新世史朝華史	
NITROGEN AMMONIA	I'10T	<u>1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 </u>	.432
	Free Am- monia.	<u>448899839884999888888899888888888888888</u>	.
EN IED.	By Suspd	\$\frac{6}{24-9} \frac{6}{24} \f	1) 7
OXTGEN	By Dis- solved.	<u> </u>	2 10.
ŏ	Total.	888989898989898	11.
	Chlori	$\frac{4}{9} + 3 + 3 + 4 + 4 + 4 + 4 + 4 + 4 + 4 + 4$	
11	on Igni- tion.		
ox.	on late and self and	్రిప్ చే — 'ఉదే ఉదే - కుకు — చేశారు ' ఉదే తే తే తే తే తే తే తే ఉద్దేది. బిక్ శ్రీక్ మా కృశ్యాత్రిక్ క్రిక్ ఉద్దారు. ఈ ' ఈ జక్కు ఉ దారు కృశ్యాత్రిక్ మా కృశ్యాత్రికి కొంటే ఈ మా కృశ్యాత్రికి కొంటే ఈ మా కృశ్యాత్రికి కొంటే ఈ మా కృశ్యాత్రికి కొంటే	71 1-
ORATI	Loss on L'10T'		- 25
Evap			
RESIDUE ON EVAPORATION.	Sus-	2656. 4 208. 8 2447.6 896. 4 122.8 763.6 896. 4 122.8 763.6 896. 4 122.8 763.6 896. 2 182.6 618.6 825.6 618.6 618.7 199.6 153.4 4 1006. 8 106.8 1560.	.10
SIDU	Dis-	2656. 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 26 . 4 208 . 8 208 .	100.
RE	Potal.	2656.4 208.8 2447.6 286.4 1208.8 2447.6 286.4 120.8 763.6 286.1 120.8 763.6 286.1 620.3 6 1252.8 286.1 6217.5 2584.4 11066.8 106.8 1550. 22018. 250.8 210.4 11066.8 106.8 1550. 22018. 250.8 120.8 22018. 250.8 121.2 22018. 2	244.4
*,1	tobO	υ · · · · · · · · · · · · · · · · · · ·	:
ANCE,	Color.	V	M. F.04
APPEARANCE	Sedi- ment.	d b db	:
	Lucht.	82800588712941289-2589529625674-65	+ 000
DATEOF	1899 Exami- nation.	April May	7 7
Dac	rs99 ('ol.ec-	Napriles Napril	::
No.	lairos -	7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	8909

Colon M., Muddy. VM., Very Muddy. T., Turbid. C., Clondy. Color on Janthon G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Readish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Black. GB., Gray-Brown. Tribbinty-*Decided. § Very Decided. *Distinct. ‡ Very Slight. Slight.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER, T. J. BURRILL, CORKS. University of Illinois. SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

No. of Bac.	per Cubic Centi- meter.		10,500 12,350 15,800	6.100	4.220 10.050 7.450	34,000	000 (c)	17,500	10.950	6.600	36,200 15,800	22,800 7,000 15,750
	Presence Abs. of			: :	++	- : ++	-+-+	-5.+-	+++	- 1 1	++-	+++
	Tempera of Air, C	51 81	8, 5; 8; 8, 5; 8;	386	8 2 8	श्रं श्रं श	* 2 2	± 5.5 €	;∞ ≈	રાં <u>∓</u>	0.77	ຂໍາຕໍາຕໍ
	riempera	: : : : : : : : : : : : : : : : : : : :	383 383	26. 27. 5.	ន្តន្តន	22.22	중 중 중 중 중 중 중	55.5	1-1-	o. o.	= x:	, 0°0
	Heigh AgieH ataW		288 288 288	0.00 0.00 0.00 0.00 0.00	6 16.6 8 14.3 12.9	8 13.9	8.6.6 6.6 6.6	20 05 170 -14 0 100 10 10	9 m 1 m m		<u>ක්ත</u> කැල් නැල්	7 7 7 7 7 7 7 7 7
NITRO- GENA	Nitrates.	017 : 58 016 : 73 006 : 64	₩									
7.	pended.	0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.										
NIC GEN.	solved.	314 2.0 412 1.9 666 3.4 57 2.0 506 4.0	44 03 03	<u>ಯ ಅ −</u>								
ORGANIC NITROGEN	-siQ											
	Total.	66.00 66	83 - 32 70 00 03 4 - 1 - 80	% ∞ ∞ ∞ ∞ ∞ % ∞	20 01 - 20 20 01 - 20		원왕. 	2000 2000 2000 2000 2000 2000 2000 200	× × × × × × × × × × × × × × × × × × ×	9.00	<u>∞</u> ∞ ∞	8 8 4
AS.	my p.	25.05.05.05.05.05.05.05.05.05.05.05.05.05										
NITROGEN AMMONIA	Albuminoid Piot7 Dis- bivlos	王载云载郑										
NITR		8 8 5 5 4 8 9 7 5 7 4 8 7 7 7 8 8 7 7 7 8 8 7 7 8 8 7 7 8 8 7 7 8 8 7 8 7		<u> </u>								
	Free Am-	6 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			0.00	<u>== 5</u>	290	<u>- 633</u>	300	.03	= 0.4 8.2.6	398
GEN IMED.	By Suspd		9 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3 ×	1-∞ 10	क्ष भट का		21 1° − ∞	0 0 m	90	≈ -	<u> </u>	_∞ w o re æ
OXYGEN	Total.	<u>1-1110 4-∞</u> ∞∞ ⊙ 51∞	3.6	- 01 -	∞ 10	<u></u>	<u> </u>	0 = 1 0 × 1	2.7.	3.9	6.5	1 × × × × × × × × × × × × × × × × × × ×
ne.	Chlori	<u> </u>						ರು ಬೆ ಬ	997	oo ₹~		
olor	on Igni- tion.	e e e e e e e	ವಾ ವಾ ೧೯ ವೆ ಮೆ ಗ	~ ~ ~ ~	ng mg mg	m m m	minimi minimi		***		0. 50	
	p,pud	रा च ७ ० च	হা হা	9 00	ରା ରା ଡ	00 00	야 아 :	න . ද	1 7 .	4.0	क् न्य	C 01
TION	Loss on Lyor Sugar	4. 20.8 43 7.226.8 40 7.226.8 40 8. 27.260 8. 27.260 8. 27.260	32.4.51 92.4.51 9.2.39	32.4 16.8 20 39.2 66	21.2 28.4 28.4 28.4 43	± 80 5- 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	\circ	26.2.2. 0.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	7. UT	9. 7.	20.8 1.6 17	18. I5 15.6 4
APOR	1'10'l'	64. 20.8 43 67.2 26.8 40 77.2 29.6 47 88. 27.2 69 103.2 16.8 86		104. 32.4 71 126.8 16.8 80 106. 39.2 66	92.4.21.2.71. 73.6.28.4.45. 73. 28.4.45.	126.8 14. 82. 76.8 28.8 48. 30.4 17.6 12.	36.4 13.2 23 43.6 24.4 19		5.6 13.6	 양양		333.2
RESIDUE ON EVAPORATION	pəpuəd sns-							##0. #30.# \$30.#	209.2 334.		349.2	# · · · · ·
UE O	solved.	2404. 198. 8 2205.2 1794.4 230.8 1563.6 2027.6 190. 1837.6 1554.8 226.8 1328. 3104.4 253.6 2850.8	3802. 219.6 3552.4 3309.6 280.4 3029.2 2420.8 217.2 2203.6	2512. 205.2 2306.8 2665.6 208.8 2156.8 2323.2 200. 2123.2	2141.6 188.8 1952.8 2390.8 197.6 2193.2 1646.4 161.6 1484.8	१७ ०० छ	०० ०१ ०१	97.5		छ य	9.	_∞ ∞
RESIL	Total.	40004	9.6 9.8 9.8 9.8 9.8	2512. 205 2665.6 208 2323.2 200	2141.6 188 2390.8 197 1646.4 161	1580.8 1283.2.18 1008.4.19	829.2 202 752.4 215 407.2 299	∞ ∞	00 =	597.2 271 375.2 236	574.8 257 668 - 257	588.8 254 558.4 258
		None 2404 1794 1794 1554 1554	3802. 3309.6 2420.8	2865. 2865. 2865. 2865. 2865.	239	<u>888</u>	왕만물	259 250 250 250	\$ 33.8 \$ 33.8	3 2	27.27	3122
*.1	tobO			: : :		• • • •	:::	: : :		3 3 3		: :
CE.	Color.	アアアアア		. F. F. S.				7.5.7 7.8.8	₹.0.₹ 30.₹	F.04 F.06	7 E E	
ARAN		1 2 2 2 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3	222		XXX <u><<<</u> <		ZZZ	ZZZ	77		SZZ	
APPEARANCE	Sedi- ment	V. M'ch	:::	:::		:::	Mucb ::	c'ons'd		Much Cons'd	C'ons'd	Cons'd
	Turb'y.		5 % S m.m.m	mm.m				* * *	* *	の <u>干</u> 。	* * *	* *
DATEOF	1899 Examination.	April May June		July ::	Aug.	17 :: 24 :: 31 Sept.	: : :	Oct.		<u></u>	30 Dec.	: :
124	· ·	ळ गुरु − ∞		<u>ంబన్</u>	\$5 to 0	1242	- 王 京	\$ 10 E			<u> </u>	10 50 T
DAT	399 11ec	ri Se se		>	bio		=			2		;
	Serial 1899 Lion.			5359 July 5408 :: 5456 ::	5514 :: 5556 Aug. 5604 ::		Ĭ.	5970 :: 6012 Oct. 6062 ::		<u> </u>	6361 5421 ::	

Tribbury * Decided. & Very Decided. + Distinct. ; Very Slight. | Slight.

Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. COLOR ON IGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown.

TABLE 30.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, AT CHAIN OF ROCKS, INLET TOWER, ST. LOUIS WATER WORKS.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

No. of	per Cubic Centi- meter.	11.000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1).4.n.
	Presence Abs, of		-
	Tempera of Air, C		. (.
r. oann	solutizates. Zitrates. Heigh officialists stanformorr officialists stanformorr	00000000000000000000000000000000000000	
ORGANIC NITROGEN.	Dis- solved.	23. 25. 25. 25. 25. 25. 25. 25. 25. 25. 25	
Sc.	Sas- Brid'd Bridge	8.68.48.88.88.88.88.88.88.88.88.88.88.88.88	
	T'01'T Discolv'd	<u> </u>	1
NITROGEN	T'10'T	2345-682-88-88-88-88-88-88-88-88-88-88-88-88-8	_
	Free Am-	+85588844888888888888888888888888888888	
OXYGEN CONSUMED.	Total, By Dis solved, By Suspd	######################################	2
ne.	Сиоги	$\frac{1}{2} \frac{1}{2} \frac{1}$	
Jolor	on Igni- tion.		
	p.doc. l.	25.56. 65. 62.76. 88. 25.66. 10. 65. 21. 21. 25.87. 25.66. 10. 65. 21. 21. 25.87. 25.66. 10. 65. 21. 21. 25.87. 25.66. 10. 65. 21. 21. 21. 25.87. 25.	0
on Ev	bended.	2	
IDUE	Dis.	226.4.2173.6 2.236.4.2240 2.246256.8242.8. 2.246256.8242.8. 6.256.8242.8. 4.192.2230.4. 1.84.82371.3. 6.182.2230.4. 1.192.82371.3. 6.183.2.230.4. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.183.2.123.6. 6.248356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356 8.256356	
RES	Trotal	8.000 8.000	
1.1	tobO	N- 	
APPEARANCE.	Color.	N	
APPEA	Sedi- ment.	V.N.ch Cons'd Much	
DATEOF	1999 1890 Collece Exami-	58 - 76 88 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	
DA	1599 ('olsection.		
No.	Infras	50.00 Nay 50.00	

Conor M., Muddy, VM., Very Muddy, T., Turbid, C., Clondy, Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Red. Blk., Black. GB., Gray.-Brown. TURBIDITY - * Decided. § Very Decided. * Distinct. ‡ Very Slight. | Slight.

TABLE 31.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

Source of Water-Mississippi River, 400 Feet from Missouri Shore, at Chain of Rocks, Pumping Station St. Louis Water Works.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois...

J.	. <u>0.:</u> :	1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	999
No.	Cubic Centi- meter.		1892 1892
	Presence 10 Presen		-+-+
	Tempera of Air, C		တ် က် က်
	rempera of Wate	. : : : : : : : : : : : : : : : : : : :	m 0 0
101	Heigh otaW	- :431 84 9 9 81 81 9 6 7 7 11 12 9 1- 0 0 0 10 7 4 10 10 10 10 10 10 10 10 10 10 10 10 10	4444
0-0 N AS	Vitrates.	<u> </u>	6. <u>6. 6.</u>
NITRO-GEN	Nitrites.	999 999 999 999 999 999 999 999 999 99	800. 400. 800.
IC EN.	sns- pepuad	888 15 15 15 15 15 15 15 15 15 15 15 15 15	
ORGANIC NITROGEN.	Dis- solved.	81-47° 644944988888888888888888888888888888888	36 296 296
ON	Total.	200	52.6
70	i p,pud	10.13	26 44 44 44
EN AS	Albuminoid Dis- Divios	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	976 113
NITROGEN AMMONIA	T'ot'T	1408448855888888888888888888888888888888	336 256
Z	Free Am- monia.	4	90.0.00
ED.	By Suspd	0.83	41-0 540
OXYGEN	By Dis- solved.	\$\int\operatorname{\sigma} \colon \text{\$\sigma} \colon \$\si	80 4 80 . 5-
CoO	Total.	8888888888888888488555555000 664486 164566 165 165 165 165 165 165 165 165 165	
.9n	Chlori	$\frac{\pi \cos \alpha \cos \alpha + \cos \alpha + 4 + \cos \alpha + \infty \cos \alpha}{\sin \alpha \cos \alpha \cos \alpha} \cdot \sin \alpha \frac{1}{16} - \cos \alpha \sin \alpha \frac{1}{16} \frac{1}{16} \cos \alpha \cos \alpha - \sin \alpha \frac{1}{16} \frac{1}{16} \cos \alpha \cos \alpha - \cos \alpha \cos \alpha \cos \alpha$	\$ 50 00 5 00 0
Color	on Igni- tion.	以识识就说就就说说话说话说话说话说话说话说话说话说话就 就	
N.	i papud -sng	F 4888	
RATIC	Loss on Ig. h. sid b'ylos -sid -sus -sid -sus -sid -sus -sid -sus -sid -sus -sus -sid -sus -sus -sus -sus -sus -sus -sus -su	66. 88. 88. 88. 88. 88. 88. 88. 88. 88.	
7APO]	L'3oT	\$26.000000000000000000000000000000000000	
RESIDUE ON EVAPORATION.	Sus-	世元 世の 00 世 せらせののののの 00の世 00のの)サ	452. 465.6 401.2
DUE	Dis- solved.	3256.4 248. 3008. 3559.2 216.8 3549. 3559.2 216.8 3549. 3555.2 202.8 206.8 3716. 3559.2 216.8 3359. 3565.2 222.8 2380. 3569.2 2381.6 2581.6 2582.8 2380. 2586.8 191.8 2571.9 2586.8 191.8 2571.9 2586.8 191.8 2571.9 2586.8 188.4 2336. 2586.8 2586.	316.4 299.3 304.
RESI	Total.	3256.4.248.3556.4.248.3556.4.248.3556.28.206.8 3813.6.228.38813.6.2896.8 32813.6.2896.8 32896.	4000
٠,١	юрО		: : :
			998
INCE.	Color		五江江
APPEARANCE	ii. nt.		
AP	Scdi- ment	> 0 0000	Much
	Turb'y.	689900000000000000000000000000000000000	. 6. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0.
SOF	1899 1899 Collec-Examition.	May June June July Scpt. Scpt. Nov.	
DATEOF	1899 Collection.	584-868866886886886888888888888888888888	
	Col	Name of the second of the seco	7 Dec 3 :.
.oV	Serial	66218 66218 66218 66218 66218 66363 66363 66363 66363 66363 66363 66363 66363 66363 66363 66363 66363 66363 66363	6533 6569

Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rb., Grayish. Blk., Black. GB., Gray-Brown. Turbibity-*Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 32.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

SOURCE OF WATER-MISSOURI RIVER, FORT BELLEFONTAINE, WEST ALTON, MO.

	•	_				^^	en.				pe	_			_		_	,,,,,				
No. of Bac.	per Cubic Centi: meter.	18.500	20.000	36 (11)		70.333	16.00		11,100	10,00	31.500	16.00	33°3		25.40	6,65	60,00	38.00	15.00	17,40	00.1	8.00.
(,0]]°	Presence 10 sd A	0.	+	:-	:	+	+	:	+-	++	1	-	1 6	,	!	- 1	+	+	+		+	:
ture Jent.	Tempers of Air, (:8	33.	35.	6:	35	6 1 6	<u>} </u>		53	<u>6</u>	FF =	÷	19.	10.		100	10	с .	?
	Temperature of Water, C.		9; 8 9. 8																			
	Helgh Mate	-	13.3																	_	_	
0. N A S	Zitrates.	21.	.36												- Andrews				_			
NITRO- GEN	Mtrites		20.0	600	0.	-S-	300.	.003	900.	800.	.007	.00.	(S)	5.5	000	900.	.00.	. OUG	800	.005	.005	.00.
S. S.	bended.		2.352	10	1:3	0)	-	_														
ORGANIC NITROGEN.	Dis- solved.		328																			
ON	Total.		2.68	0 0	01	2.1	_	-		-	Special		_						_	_		
vo.	gus-sus		.736																			
EN AS	Piot'l		821.	0.0	-	-:		.65	9 20	01:	:00:	.095	0.0 23.00 23.00	8 8	<u> </u>	-	<u>5</u> 1.	31.	. 113	: 1	960.	se.
NITROGEN AMMONIA.	TotT		864	1.024	1.052	1.024	30 T.	544	.116	3 ∞	. 116	5X2.	00 0 00 0 01 0	X	333	208	.852	<u>e</u> :	333	.116	272	.176
7.	Free Am- monia,		610	200	910.	3	.058	30.	20.	900	.024	.012	1.00		850	0330	0	.035	50	10.	90.	80.
ED.	By Suspd		35.4	50.0	19.7	10.8		0.0	ص ن ن	6.1	6.7	ت. ن	3? -	- ×	000	22	6.1	-	6.4	(-) (-)	6.3	<u>-</u> -
OXYGEN	By Dis-		₹- C		-10	10	÷		ಲಾ ಅ		÷	200	ec c		77	~	-	33	~	٠: ص:	-	च <u>ैं</u> च
င်္သ	Total.		4.0	3	133	16	0	2		3 ==	2	9	2	- 3	-1	9	19	9	1-	=	=	30) 10
.əu	Chlori	10.	10.10	. c	3.00	6.3	 ∞	3.0	6.01	14.8	16.3	<u>-</u>	<u>∞</u> 0	ت 20 ع	5 6	£9.9	16.15	0:2	<u>∞</u> .∞	6.91	67	
Color	on Igni- tion.		<u>=====</u>	<u> </u>	~	13.	::	~	=======================================		5.	<u>~</u> ;	<u></u>	<u>-</u> -		~~	<u></u>	~	~	~	15(5,	5
ż	e papud		118 8	. 9	95.6	57.2	30.8	24.	7:4		21		9	2 X	9			9.1	1.6	11.4	9.1	9.1
110	D 1108			-					25 -			00	<u>د</u> و									9
1 4	on sid	:	7.5	000	61.4	25.6	77 77	16.	24.0		9.6	-		x; =	00	13.6	16.8	15.2	<u>~</u>	18.	13 6	≘
FAPORA'	l'10T		133.6 14.8 118	-	_		43.2.22.4	10. 16.	22.4.24.0	23.2 11.	22.4 9.6	-	16.4	13.6 TO.8	00 00 00 00	.8 13.6	16.8	16.8 15.2	16.4 11.8	32.4 18.	15.2 13 6	15.2 13
ON EVAPORA	088			-	_		53	16.	031.6 48.4 24. E	23.211.		.8 33.6	. i .	221.6 13.6 10.8 286 61 6 11 1	3.86.1 18.8 8.8	.1 18.8 13.6	16.8	117.2 16.8 15.2	511.8 16.4 11.8	535.2 32.4 18.	180.4 15.2 13 6	180.4 15.2 13.
DUE ON EVAPORA'	Solved.		2 3044 8 1	1 2790 8	2133.21	.1,1730.	6 1279.2	903.6 10. 16.	931.6	641.1 23.2 11.	1 488 + 23.4	2 502.8 33.6	405.2	521.6 286	336.1	.8 316.1 18.8 13.6	568.8 22.8 16.8	8 117.2	2 511.8 16.4	535.2 32.4	180.4	36.8 180.4 15.2 13.
RESIDUE ON EVAPORATION.	Sus- pended.		201.2 3044 8 1	8 0020 1 901 0	2133.21	196.1,1730.	6 1279.2	1248.8 903.6 40. 16.	200.4 931.6	3277.3 641.1 23.9 11.	373.1 488 1 23.4	307.2 502.8 33.6	4 301.2 405.2	521.6 286	1336. 1336.1	.8 316.1 18.8 13.6	8 314. 568.8 22.8 16.8	318.8 117.2	317 2 514.8 16.4	535.2 32.4	180.4	17.2 336.8 180.4 15.2 13.
	Potal. Dis- sultanted. Sus- pended. I'tol'	one	3246. 201.2 3044 8 1	8 0020 1 901 0	.6 2133.2 1	.1,1730.	221.6 1279.2	.4248.8 903.6 40. 16.	931.6	3277.3 641.1 23.9 11.	373.1 488 1 23.4	810. 307.2 502.8 33.6	405.2	831.2 309.6 321.6	1336. 1336.1	(87.2 310.8 346.4 18.8 13.6	8 314. 568.8 22.8 16.8	8 117.2	2 511.8 16.4	535.2 32.4	180.4	517.2 336.8 180.4 15.2 13.
	Odol Lond. Loures Loures Loures Loures Loures	None	3246. 201.2 3044 8 1	0.000 1.000 0.000 0.000	. 2178.8345.62133.21	196.1,1730.	6 1279.2	1248.8 903.6 40. 16.	1192. 330.4 931.6	921.6 277.2 641.1 23.2 11.	: 860.8 372.1 488 1 23.4	810, 307.2 502.8 33.6	706 4 301.2 405.2	531.2.309.6 521.5 501.2.309. 502	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	* 832, 317 2, 514.8 16.4	535.2 32.4	180.4	2
	Potal. Dis- sultanted. Sus- pended. I'tol'	I. None	3246. 201.2 3044 8 1	1 0.00 1 4 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 1 1	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	F.04 : 1500.8 221.6 1279.2	F.02 · · · 1152.4248.8 903.6 40. 16.	200.4 931.6	921.6 277.2 641.1 23.2 11.	: 860.8 372.1 488 1 23.4	810, 307.2 502.8 33.6	706 4 301.2 405.2	831.2 309.6 321.6	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	317 2 514.8 16.4	535.2 32.4	180.4	f. F.02 · 517.2 336.8 186.4 15.2 13.
	Olon Total Total Total Total Sus- Sus- Bonded.	M.	3246. 201.2 3044 8 1	0.000 1.000 0.000 0.000	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	F.04 : 1500.8 221.6 1279.2	1248.8 903.6 40. 16.	1192. 330.4 931.6	921.6 277.2 641.1 23.2 11.	: 860.8 372.1 488 1 23.4	810, 307.2 502.8 33.6	706 4 301.2 405.2	531.2.309.6 521.5 501.2.309. 503	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	M. F.C. : 832, 317 2, 514.8 16.4	M. F.03 * 845 2 310. 535.2 32.4	N. F.03 - 517.2 336.8 180.4	M. F.02 . 517.2 336.8 180.4 15.2 13.
	Nedi. Color. Total. Dis-solved. Sus-pended. Sus-pended.	Consid M. None	3246. 201.2 3044 8 1	1 0.00 1 4 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 1 1	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	F.04 : 1500.8 221.6 1279.2	F.02 · · · 1152.4248.8 903.6 40. 16.	1192. 330.4 931.6	921.6 277.2 641.1 23.2 11.	: 860.8 372.1 488 1 23.4	810, 307.2 502.8 33.6	706 4 301.2 405.2	531.2.309.6 521.5 501.2.309. 503	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	* 832, 317 2, 514.8 16.4	M. F.03 * 845 2 310. 535.2 32.4	N. F.03 - 517.2 336.8 180.4	M. F.02 * 517.2 336.8 186.4 15.2 13.
	Turb'y. Sedi. Color. Total. Total. Dis- solved. Sus- pended.	M.	3246. 201.2 3044 8 1	1 0.00 1 4 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 1 1	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	F.04 : 1500.8 221.6 1279.2	F.02 · · · 1152.4248.8 903.6 40. 16.	1192. 330.4 931.6	921.6 277.2 641.1 23.2 11.	* M. F.02 860.8 372.1 488 1 22.4	810, 307.2 502.8 33.6	706 4 301.2 405.2	531.2.309.6 521.5 501.2.309. 503	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	M. F.C. : 832, 317 2, 514.8 16.4	M. F.03 * 845 2 310. 535.2 32.4	* N. F.03 - 517.3 336.8 180.4	* N. F.02 · 517.
APPEARANCE.	Turb'y. Sedi. Color. Total. Total. Dis- solved. Sus- pended.	10 * [Cons'd N.	28 N. Web M. F.08 . 3246, 201.2 3044 8 1	1 0.00 1 4 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 1 1	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	: 1 % · V.M. F.O. · 15(0).8 [331.6] 1279.2	8 * Much M. F.02 · 1152.4 248.8 903.6 40. 16.	1192. 330.4 931.6	* N. F.02 921.6 277.2 641.1 23.2 11.	* M. F.02 860.8 372.1 488 1 22.4	810, 307.2 502.8 33.6	20 * N. F.03 706 4 301.2 405.2	2. *	672.4 336. 336.1	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	736, 318.8 117.2	M. F.C. : 832, 317 2, 514.8 16.4	M. F.03 * 845 2 310. 535.2 32.4	N. F.03 - 517.2 336.8 180.4	* N. F.02 · 517.
APPEARANCE.	Turb'y. Sedi. Color. Total. Total. Dis- solved. Sus- pended.	10.July 10 * Cons'd M.	27 . 28 V. N.ch M. F.08 . 3246, 201.2 3044 8 1	0.000 + 12 + 13 + 140	7.1 2178.8 345.6 2133.2 1	F.05 1926.4 196.1,1730.	F.04 : 1500.8 221.6 1279.2	8 * Much M. F.02 · 1152.4 248.8 903.6 40. 16.	1192. 330.4 931.6	* N. F.02 921.6 277.2 641.1 23.2 11.	Oct. 6 * M. F.02 860.8 372.1 488 1 23.4	810, 307.2 502.8 33.6	20 * N. F.03 706 4 301.2 405.2	26 : 27 * N. F.03 :: 831.2.309.6 521.6	0 . 10 *	687.2 310.8 346.4 18.8 13.6	882 8 314. 568.8 22.8 16.8	. 1 * . N. F.01 . 736, 318.8 417.2	M. F.C. : 832, 317 2, 514.8 16.4	M. F.03 * 845 2 310. 535.2 32.4	* N. F.03 - 517.3 336.8 180.4	* N. F.02 · 517.
	Total. Total. Total. Total. Total. Total. Sus- pended. Sus- pended.	10 * [Cons'd N.	27 . 28 V. Nich M. F.08 . 3246, 201.2 3044 81	1 0.00 1 4 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 0.00 1 1 1 1	7.1 2178.8 345.6 2133.2 1	. 21 . 25 \$. VM.F.05 . 1926.4 196.1,1730.	: 1 % · V.M. F.O. · 15(0).8 [331.6] 1279.2	Nept. 7 8 * Much N. F.02 1152.4 248.8 903.6 10. 16.	14 . 15 8 . N. F.03 . 1192, 280,4 931.6	28 . 29 * N. F.02 . 921.6 27.2 644.4 23.2 14.	Oct. 5 Oct. 6 * " N. F.02 " 860.8 372.1 488 1 22.4	" 12 " 13 * " N. F.03 " 810, 307.2 502.8 33.6	. 19 · 20 *	20 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	10 * N. F.02 672,4336. 336.1	** 16 · 18 * · · N. F.02 · · (87.2 310.8 346.1 18.8 13.6	. 93 . 24 * 1 N. F.02 . 882 8 314. 568.8 22.8 16.8	" 30 Dec. 1 * " N. F.01 " 736, 318.8 417.2	Dec. 7 . 9 * Cons'd M. F. 02 . 832, 317 2 511.8 16.4	" 11 " 16 * Much M. F.03 " 845 2 310. 535.2 32.4	* N. F.03 - 517.3 336.8 180.4	· 28 · 29 * · 517.

Colon M., Middy. VM., Very Muddy. T., Thrbid. C., Cloudy. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Bik., Black. GB., Gray Brown. Color on Ignition—G., Gray. B. Brown. DB. Dark Brown. LB. Light Brown. Rh., Reddish. W., White. Gh., Grayish. TURBIDITY-*Decided. §Very Decided. * Distinct. ‡ Very Slight. | Slight.

333 TABLE DATEOF

6681 tion.

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

No. SOURCE OF WATER-MISSISSIPPI RIVER, 100 YARDS FROM ILLINOIS SHORE, JEFFERSON BARRACKS

Report of Arthur W. Palmer, T. J. Burnill. University of Illinois.

45.500 10,050 10,050 11,850 14,250 18,700 25,600 19.500 71,000 9.000 1.500 1,500 5.000 24.500 11.000 000.993 33,000 60.00 Cubic Centimeter. Presence or Abs. of Colf. Temperature of Air, Cent. l'emperature of Water, C. lleight of Water. GENAS Nitrates. NITIO-100 88 887 00.4 012 ORGANIC NITROGEN. solved. ·sid क्ष 🖭 Total. 15.00 Albuminoid Am. p,pud -sns AS AMMONIA. NITROGEN 1.101 Ргее Аш-тоніа. 7. 1. 2. 1. By Suspd ONYGEN CONSUMED. 3 -si(l ya solved, Total. Chlorine, Color tion. g on ne 9220.4 | 196. 2024.4 | 96. | 19.6 | 76.4 | 1519.6 | 187.6 | 133.2. | 47.6 | 13.6 | 34. | 1519.6 | 187.6 | 133.2. | 47.6 | 13.6 | 34. | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 | 13.0 p,pud -sng Loss on 1g'n. RESIDUE ON EVAPORATION. -siQ b'vlos 1,10,1 bended, -sns solved Odor. M. F.94 M. F.94 M. F.94 V.M. F.94 V.M. F.94 V.M. F.95 M. F.93 M. F.93 M. F.95 ('olor. APPEARANCE. V.M'ch Much Cons'd Cons'd Sedi-ment. Much Cons'd Cons'd Much Much Lauply 15 25 29 29 32893 Collec- Examination.

Sept.

5826 Sept.

5860

::::

Oct.

6018 Oct.

6058

: : :

Ang

: : :

33.77

: : :

6 July

5365 July

3223 Nov.

C. Cloudy. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Blk., Black. GB., Gray. Brown. T., Turbid. V.M., Very Muddy. Colon-M., Muddy. Color on Ignition—G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. Rh., Reddish. W., White. Gh., Grayish, Trubinity—*Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 34.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Source of Water-Mississippi River, East of Midstream, Jepferson Barracks, Mo.

ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.	
Report of	

No. of Bac.	per Cubie Centi- meter.	71.000	16,000	10.350	22,450	-1.80 -1.80	12,700	16,550	30,800		30.30 30.30	126 500	17,500	16.000	15,000	68.000	000	18,050	4.000	8.000	6.250	15,000	11.750	1350	21.000
Coll.	Presence Abs. of		:		:	:	:	1	+	:	ī	: +	+	+-	+-	- 1	ł	- +	+	+	+	+	٠	+-	
	Tempers (:	: .02 :03	:					:	:			:	:						:		:		
	risqmoT stall lo	:	9.3	189	26.		8 8	33	65		200		61	6	<u>-</u>	ਨੂੰ ਨੂੰ	19	6:	6	Ġ.	Ξ.	<u>ei</u> :	oi n	: :	. 0
	dgi9H Mata		2) ? ?} ?	20.5	60.00		0 0 0 0 0 0				T in	- 9	6.8	9	- 5 - 5	1 10		33	-	5.4	6.5	9:0	20	. 4	. vo
TRO. GEN AS	Sitrates.	.52	20.00	. 56	\$	٠.	7 :5	3	7	Ţ.,	7 -	= = = =	ક્	= :	. ÷	; e ;	6	3,1	33	.36	35	: 3	50.4	38	Ŧ.
NITRO-GEN	Nitrites.	200.	50.00	88	300	D.	300 300 300	800	3.	600.	90.	00.	300.	CO:	011	18	800.	.005	.013	.013	<u></u>		+10.	0.00	.0.
	-sns	- 19	3.776	18:00	384	£45.	959	80%	248	52.	2 x	38	986	97	2000 2000 2000 2000 2000 2000 2000 200	676	.695	.656	459	?? ??	.576		2	336	116.
ORGANIC NITROGEN	solved.	15	504	344.2	536 2	2 900	953	392	47.2.1	36.1	200	344	384	- 200			X 22.55	∞ ??	30 31	250	2 2 -		010.	7.	456
ORG	Total.																				06:1		108		
			168.4																				010	-	-
AS.	Albuminoid Am. Sus. Solv'd Sus. Solv'd Sus.			•															_	0.5	-				
	eid solv'dos	·	8 8																	307	•		578.		
NITROGEN AMMONIA	I ToT	9.1	1.04 38.1	.928	1.024	318.1 200€	5.10	.672	105.	9.6	102	355	38	916.	155	38	£ +.	.116	**	227	<u> </u>		200	43.5	<u>∞</u> .
Z	Егее Ата. топа.		9,00													035	10.	034	.01	3330	3			910.	.0.14
N ED.	By Suspd	55.5 65.5	28. c	19.4	1.81	£ 8	5.0	6.3	2.1	10 E	70 cc	2)	5.5	· ·	- - -		20.4	??		-:	6.6	: 0	3.0	3 53	ري. ري
OXYGEN	By Dis-	1 . 1	2 C	: -:	11.	اه ب	11. 11.		10		ر ا ا	73	F. C		200		{ - ; - ; - ; - ; - ; - ; - ; - ; - ; -	i i -	.c.	σ; ∝			# 0X		9.01
CON	Total.	5.5		35,50	30.03	7.7	20. T	{-	X.	:C 0		**	57	77 S	2 5	6.	∵	æ. ∞	9.6	<u>.</u>	-:		0 7	œ	13.1
ne.	Chlori		20 °C		00.0		-		10	x, 1		6.35	5.7	G: :	÷ 5	7	6.8	6.	c: ×				 22 23 O	- x	
Color	on gni- tion.		zi z						-:	~		~	=======================================	· -	<u></u>				~	~		:	<u> </u>		13,
	i p,pnd	- - - - - - - - - - - - - - - - - - -	x 00	9	5.65 5.60 5.60 5.60 5.60 5.60 5.60 5.60	0	- - 9	63	5.6	8.0	· ·		23.		c c	8.9		-:		0)			+ x		
TION	Loss on lg'n	15.2 66	110 129 250	31.634.	200	99.6 36.8 6.	(1) (1) (1) (1) (1) (1) (1) (1) (1) (1)	86. 28.8 57.2	5.6	116. 35.286.	K 0	1 50					15.2	18.91	-	00: 00:	:x:	:	00	??	9.1
PORA	1.10,L	82.	20.02		59.6 24.	M. C.	ता का के को इ. स	6. 2	÷	116.03	25.00 25.00	31.8	25.2.21	2012		\$? 	??	29.21	99	£.	S		93 6 18	×	13 6 1
RESIDUE ON EVAPORATION.	рерцеф	6.4	- T - S - S - S - S - S - S - S - S - S - S			_		:0	_		c :	-			.5	_		25	97		6.05			_	133.
E ON	-sus	8 253	X X X X X X X X X X X X X X X X X X X	6 183	201.2.2083.6		196, 8, 1626.	Mar.		2 Te									_		??	٠			
SIDU	-sitt solved.	. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.	34	219	30.		196.	196.	146.	2000	<u> </u>	197.6	200	27	13.	3.5	2(5)	2(E)	2000	300	161	0 001	300	8.	190.4
RE	Total	755.	3136.1 955. 3448.4 3148. 1964.8 3183.6	2051, 4 219, 6 1834, 8	20.584.8	21 22 0 00 0 1 2 1 2 1 2 1 2 1 2 1 2 1 2	1805 × 196 × 1626.	2207.6 196.	1306. 146.	300 0 108.8	991 8 913 6	610.	28. x. 182	451.0 E01.7	200	333.6 187.2	368	369, 6 260, 1	340	3350.6 206	210.1.161		50.00	261.6	322.1
1.1	орО	je j	: :	: 01	\$ 1					: :		:			:	:				d F	:		3110	:	:
		10	- - <u>10</u>	200	is.		- 10		윤 :	ĿG .	_ 22	33	103	Z 19	Q I	_).1	*	+	£.		-		7	<u>a</u>
VCE.	Color.	. F. S.	マンド・下、3 マンド・下、3 15	N. 1. 08		1217	V.N. F.01	V.N. F.04	V.N. E.03	N. F.05	- 12 K	M. F.03		20.7	E.0.7				F.04		. <u>F. 15</u>		. F. 5		
A II A		N		スシ		- 5		-	-			7.	7.	7. 7	7. 7	7.	7.	7.				•	Z		
APPEARANCE	Sedi- ment	Much		**	: :		:	:	:	: :	:	Muck	:		p.suo.)	:	:	9 0	:	MIICE	p.<(10.)	1,000	Much	p, suo.)	:
	∵v.q.m.L. ÷ ÷		0.0	-	(-)		0-00		27	200 V		.0.	15 ×	* *	*	* 2	* 00	* 1-2	*	*	* - :	* *	* - x	* 10	*
OF	Exami- nation.	6 June	: :		7[11]		9	Aug.	:				;				;		. :				: :	:	21
DATEOF			2 8		. 6.		2 27		11	- 1	8 E	1.	= :	27 0	0 5 0 5		19		· .	=	16		-		<u>=</u>
	collec- tion.	June	: :	:	July.		:	1112.	:	: :	: :	Verit.	:	: :	()()	:	*	ur e	.70%.	:	:	0 d 0 td] Jeef		:
No.	Bitros	5169.1	5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	531*	5363	1111	37.0		500	2655	2 10			250	5675	5000	6114			62/33	63320	6371	545	•	6562

TUREDITY * Decided. § Very Decided. † Distinct. ‡ Very Slight. ¶ Slight. (Onon. -M., Muddy. V.M., Very Muddy. T., Turbid. C., Cloudy. Colon on Ionition - G., Gray. B., Brown. DE., Dark Brown. LE., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of Arthur W. Palmer,	T. J. BURRELL,	University of Illinois.	

io. of	per Cubic Centi- meter.	1 1 1 1 1 1 1 1 1 1	11.400
.ilo!	yps of (-
.tns	Temperator of Air, Careence	0,000	
.O.,	remperaling of the state of the		3
10.	idgiəH isteM	::: : : : : : : : : : : : : : : : : :	500 EC
NITRO- GEN AS	Nitrates	2000 2000 2000 2000 2000 2000 2000 200	
NI	pended.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	-
ANIC OGEN.	solved.	## 6 4 4 4 4 5 6 8 8 8 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	_
ORGANIC NITROGEN.	Total.	0: 0:0:0:0	
	i p.pud	4 8 8 5 5 6 9 8 8 8 8 6 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8	
N AS IA.		200 200 200 200 200 200 200 200 200 200	
NITROGEN AMMONIA	I'Jo'T esid b'vlos	元元元	
NE	Free Am- monia.	99.88.89.89.89.89.99.99.99.99.88.89.89.8	_
ED.	By Suspd	$\frac{8508829226896565656588884849494948944}{26648668568868888484869494944}$	7
OXYGEN CONSUMED	By Dis- solved,	$\alpha \alpha \psi \circ C + \alpha \alpha \circ \alpha \alpha + \alpha \psi \circ \alpha \psi \circ \alpha \circ \alpha \circ \alpha + \alpha \circ \alpha \circ \alpha \circ \alpha \circ \alpha \circ \alpha \circ \alpha$	эc
	Total.	# 	<u> </u>
	Сріогі	သလယ္သန္နန္န္အန္နန္နန္နန္နန္နန္တန္တန္တန္တန္တန္တန္တန္တန	- 2
Cole	on Igni- tion.	<u>್ 'ಶಕ್ತ 'ದೆ ' ' ಚಿತ್ರಕಕ್ಕಕ್ಕೆ ' ಕಕ್ಶಶಕ್ಕರಾದರು ಪ್ರಾಪ್ತಪ್ತಪ್ರವನ್ನು ಪ್ರಾಪ್ತಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರಪ್ರ</u> ಪ್ರಪ್ರ -	×
FION.	i p,pud -sns	88 1 2 8 8 1 2 8 8 8 2 8 2 8 2 8 2 8 2 8	? <u>?</u>
PORA'	Tot'l	2.5. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	00 x x
RESIDUE ON EVAPORATION	Dengeg.	12-18-88-88-88-98-18-88-18-88-88-88-88-88-88-88-88-88-88	
ote o	solved.	<u> </u>	2)
RESIL	Total.	24.89. 6 22. 170. 8 33.31. 4 23.7. 6 33.31. 4 23.7. 6 33.31. 4 23.7. 6 33.31. 4 23.7. 6 33.31. 4 23.7. 6 33.31. 4 23.6. 4 23.2. 4 23.2. 4 23.2. 6 23.3. 2 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3. 2 23.3. 6 23.3	. 1
• 1		a a contract of the contract o	_
	robO		_
NCE.	Color	VAN. E.O.S. WAN. E	-
APPEARANCE	nt.		_
AP	Turb'y.	W.W.ch Water Cons.d Cons.d Cons.d Cons.d Cons.d Cons.d Cons.d Cons.d	
		<u> </u>	222
DATEOF	1899 rs- Exami nation	May 10 10 10 10 10 10 10 1	
DA	1899 Collection.	May June June June June Sept.	
No.	Serial	+ + + + + + + + + + + + + + + + + + +	Face

Tubbidity-* Decided. § Very Decided. + Distinct. ‡ Very Slight. | Slight.

COLOR ON IGNITION G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Gh., Grayish. Blk., Black. GB., Gray-Brown.

Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy.

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION

LNER, of Illinois.	No. of	per Cubic Centi- meter.	39,000 113,000	38.750 38.750 38.750 30.300
	(;0]]:	Presence Abs. of		-+++:
HCE W. PA. BURRILL.		Tempera of Air, (Q Q	
ARTHUR W. PA T. J. BURRILL, University	9TH H	riempera	क्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्रक्षेत्र 	<u>င်ရုံ</u> ထံကဲ့ရင်
	10.1	 figioH	& & & & & & & & & & & & & & & & & & &	: 0 0 kg 子 = 16 : 0 0 kg 子 = 16
Report of	NITEO- GENAS	Nitrates.		0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
		pended.	######################################	
RANALYSIS—PARTS PER MILLION RIVER, WEST OF MIDSTREAM, JEFFERSON BARRACKS, MO.	ORGANIC NITROGEN	Dis- solved,		5.55 5.55 5.55 5.55 5.55 5.55 5.55 5.5
HACK	NO.	Total,	& 2 1 2 2 2 2 2 3 4 5 5 5 6 5 6 6 6 6 6 6 7 6 7 6 7 6 7 6 7	28.53.5 2.53.5 3
N BAB	w.	g p.pud sns		8 2 2 2 8
TON	NITROGEN AS AMMONIA.	Torr Discosolvo Sus- Sus- Sus- Sus- Production		188 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1
MILI, JEF	NITROGEN	Tot7	2. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	
PER		Free Am-		2000 2000 2000 2000 2000 2000 2000 200
ARTS	OXYGEN CONSUMED.	solved. By Suspd	<u> </u>	-ressre :re
IS-P.	OXY	Total. By Dis-	3888822225884888475555527777777878888888888	
ANALYSIS-PARTS PER MILLION (VER, WEST OF MIDSTREAM, JEFFERS	ne.	Chlori	44464469999999999999999999999999999999	16 16 16 16 16 16 16 16 16 16 16 16 16 1
R AN.	Color	on Igni- tion.		
SANITARY WATER Vater-Mississippi R	o.x.	i p,pud	2. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	4.0 T 7.0 14.0 T 7.0
SANITARY WATE Water-Mississippi	RESIDUE ON EVAPORATION	no sid bylos	28.20.20.20.20.20.20.20.20.20.20.20.20.20.	20.2 20.2 20.8 20.8 20.8 20.8 20.8 20.8
NITA FER-	Evap	bended.	52522000000000000000000000000000000000	
to the same	E ON	solved.		क्ष्य विकास
RCE OF	ESIDI	-sid		
SOURCE		Tstoff'		281.6 5.80.4 5.34.6 5.34.6 5.35.6
,		TobO	None	
	CE.	Color.	M.W.W.W.W.W.W.W.W.W.W.W.W.W.W.W.W.W.W.W	
	APPEARANCE			
	APP.	Sedi- ment	V.W.ch Much Much Much Much	
		Turb'y.		=====================================
	DATEOF	1899 1899 Collec- Exami- tion, nation,	66June 66June 8887773 Aligher 6 July 88877 3 Aligher 88877 3 A	:::50 Dec:::
	DA	1599 Collection.	June 6 June 8 June 7 Ju	21,3336
	'ON	Inrios	N	

52531 ... 100 52

6021 Oct. 6056 ... 6056 ... 6256 ... 6256 ... 6256 ... 6256 ... 6256 ... 6256 ... 6256 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556 ... 6556

Colon M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Lentron G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rd., Gray-Brown. TURBIDITY -* Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 37

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISSUPY RIVER, 100 VARDS PROM MISSOURI SHORE, JEFFERSON BARRACKS, MO.

University of Illinois,

Report of ARTHUR W. PALMER,

T. J. BURRILL,

72.500 5.500 19.700 9.750 18.600 36.850 00.00 11,660 36,400 16.000 24,000 59,500 82,000 46,000 27,000 50,500 71,000 000.00 04.000 49.500 No. of Bac. per Cubic Centimeter. Presence or Abs. of Coll. Temperature of Air, Cent. QQC+8889446899 \(\alpha + \alpha + \alpha \alpha \alpha \)
\(\alpha + \alpha + \alpha \alpha \)
\(\alpha + \alpha + \alpha \alpha \)
\(\alpha + \alpha + \alpha + \alpha \)
\(\alpha + \alpha + \alpha + \alpha \)
\(\alpha + \alpha + \alpha + \alpha \)
\(\alpha + \alpha + \alpha + \alpha + \alpha \)
\(\alpha + \al lemperature of Water, C., 48 19.8 28 leight of Water, GENAS NITRO-Zitrites .472 3.968 36 2.72 .92 2.24 .392 1.368 .314 2.976 576 576 576 576 999 99 ORGANIC NITROGEN. -sns Dis-solved. 85.8 936 11.168 4.44 3.08 944 3.16 1.472 4.76 .928 3.32 .544 2.3 .648 2.76 .708 2.36 744 2.6 744 2.3 312 1.4 Total. .224 1.696 2.6 .192 1.168 4.4 .096 1. 3.0 248 Albuminoid Am. p.pud -sng NITROGEN AS 176 104 12 156 961. 184 12 88 52 AMMONIA 1.1.63 2.2.5 3.6 3.6 3.6 3.6 3.6 .768 [,10,] 016 032 028 04+ 036 CONSUMED OXYGEN 15.4 7.97.7 20.2 8.87.1 12.9 11.87.6 13.8 10.3 6.1 By Dis-solved. Total. Chitorine. gni-Color tion. on 医玻璃玻璃玻璃玻璃玻璃玻璃玻璃玻璃玻璃玻璃玻璃 p.pud -sng Loss on 1g'n. RESIDUE ON EVAPORATION. siti p,vios 1.10.T bended -sns solved. -sid Total. Odor. M F.03 Color APPEARANCE. V.M'ch V.M Consid Much Cons'd Cons.d Cons'd Sedi-ment. Much Much Much Lurby * - 22 8 23 0.10 85 8 5 Collec-|Examination. 1899 July Nov. 6 Oct. DATE OF 52220 833223 三部器 1899 tion. 5166 June 5362 July 5416 : 5165 : 5510 : 57718 5768 5822 Sept. 5862 5932 6963 6174 "... 6224 Nov. 5559 Aug. 5020 Oct. 5619 5319 0000 62.06 5656 Serial No.

C., Cloudy. n. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red Rh., Reddish. W., White. Gb., Grayish. Blk., Black. GB., Gray-Brown. T., Turbid. V.M., Very Muddy. COLOR-M. Muddy. DB., Dark Brown. LB., Light Brown. § Very Decided. + Distinct. + Very Slight. | Slight. COLOR ON IGNITION-G., Gray. B., Brown. TURBIDITY-*Decided.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ST. LOUIS, MO., TAP WATER.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

No. of Bac.	per Cubic Centi- meter.	910 910 910 910 910 1,180 910 1,180 1,320	4,050 1,450
	Presenc	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	+ 1
ent.	Tempers of Air, C	9	
	Tempers	6 5 5 5 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	-:
	Heigh JieW	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	£ 21
NITRO- GENA	Nitrates.	668988	
	pended.	88	
ORGANIC NITROGEN.	solved.	######################################	
ORC	Total.	888888	
	i p,pud	20000000000000000000000000000000000000	
EN AS		525 5 125 5 14 15 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	.080
Nitrogen A	Tot'l Dies-	48 88 88 88 88 88 88 88 88 88 88 88 88 8	5.5
Z	Бгее Ап-	00000000000000000000000000000000000000	© 9.
ED.	By Suspd Matter.	<u>∞ ∞ ∞ ∞ ∞ → ∪ ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;</u>	<u>- m</u>
OXTGEN	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	5.9 7.6.1
	Total.	三三級認義三三級與時代為為[2] 中世年高年年十五年[2] あ[2]	15 G.
	Chlori	404444 :4000000000000000000000000000000	_ <u>10</u>
	Igni- Igni- tion.		ا ت ت ت
	o siq b'vlos p sss hid'd	る	17.6
PORA	Torra on 1:10 T		15.6
RESIDUE ON EVAPORATION	Dended	88838 8888 8888 8888 8888 8888 8888 88	
CE O	solved.	· · · · · · · · · · · · · · · · ·	S)
RESII	Total.	408.25	318.8.280
•.	tobO	0:::::::::::::::::::::::::::::::::::::	::
			23 33 24
ANCE.	Color	MANAN MANANANANAN MANANAN MANA	M. F.03
APPEARANCE	sedl- ment.	Consid Consid Nuch Consid Consid Much Consid Much Consid Much Little Consid Much Little Consid Much Little Consid Much Little Consid Much Much Much Much Much Much Much Much	_
	Turb'y.	* 55.* 55.55.* * * * * * 55.* * * * * *	* + 4 6 6 7
OF	1899 Examination.	2886 2886 2886 2886 2886 2886 2886 2886	: : : :
DATEOF	Collect Examition, nation.		T 67
No.	Serial		රනවා රනවා

Conois M., Muddy, VM., Very Muddy, T., Turbid, C., Choudy, Trubibiry-*Decided. \$Very Decided. * Distinct. *Very Slight. Slight.

Color on Lowing Brown, B. Brown, D. Brown, L. Light Brown, R. Reddish Brown, BG., Brownish Gray. Bh., Brownish. R., Red. Color on Lowing Gray. Brownish. R., Red. R., Black. GB., Gray-Brown.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

AVENUE.
KEDZIE
CANAL,
WATER-SANITARY
OF
SOURCE

	1		
LMER. of Illinois.	No. of Bac.	Cubic Centi- meter.	29,025,000 2,000,000 2,000,000 2,000,000 2,000,000
	e or Coli.	Presence Abs. of	<u>[++++ </u>
HURRILL, University		Tempera of Air, C	∞ τ Ω ω τ τ τ α τ α τ α τ α τ α τ α τ α τ α τ
RTHUR W J. BURRI		Tempera	. සහ
of Ar		Heigh 9187/	
	O-N AS	Nitrates.	284-1844444455555405540554055555555555555555
Report	Nirro- GEN	Nitrites.	
	υż	bended.	888 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
i l	ORGANIC NITROGEN.	Dis- solved.	\$\frac{\pi}{8}\fra
RIC	Orr	Total.	83.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5
IST		i p.pud	0.00 0.00
Y D	SN AS		88888888888888888888888888888888888888
TARY MILLI AVENUE	NITROGEN	Albuminoid Pis-sid Solvid	### ### ### ### ### ### ### ### ### ##
	N	Free Am- monia,	22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1 53 P	- G	By Suspd	$\partial \psi \omega v_1 + \psi \psi - \psi \omega v_2 + \psi \psi \omega v_3 + \psi \omega v_4 + \psi \omega v_5 + \psi \omega \omega \omega \omega \omega \omega v_5 + \psi \omega \omega v_5 + \psi \omega \omega$
JAGO S - PARTS CANAL, KI	OXYGEN CONSUMED.	By Dis- solved.	
J - 6	OX	Total.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
- CHIC NALYSIS SANITARY	ne.	Chlori	######################################
ANA -SAN	Color	on Igni- tion.	ening in the second of the sec
VATER WATER		p,pud	10. 13.4400 10.01 13.4100 10.000
INATI WATE OF WATE	EVAPORATION	E P.Alos	84.65.78.48.08.64.64.64.64.64.64.64.64.64.64.64.64.64.
EXA NI NITARY SOURCE	APOR	Torr S	
SANITARY SOURCE	on Ev	bended.	88 88 88 88 88 88 88 88 88 88 88 88 88
STREAMS EXAMINATI SANITARY WATE SOURCE OF WATE		Dis-	### ### ### ### ### ### ### ### ### ##
SEA	RESIDUE	Total,	223
SE		[0+0,17]	
	u ——	obO	Gas'y Oils Oils Gas'y "" Gas'y "" "" "" "" "" "" "" "" "" "" "" "" ""
	ធំ	Color.	RESERVED
	RANC	- ప	zazazzzz ze ezetetetetetetetetezz z zazazo
	APPEARANCE	Sed1- ment.	Much Cons'd Much Little
		Turb'y.	*+*+++++++++++++++++++++++++++++++++++
		1900 Exami- nation.	3 - 6888 8 - 0 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2
ő	DATEOF	Exa nati	Man Man July Seples
CE 39	DAT	1900 Collec- tion,	
ABLE	1017	-	### Page 19 Pa
E)	1 OX	Serial Serial	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\

Turbidity—* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight. | Color — N., Muddy. V.M., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. 16., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White

TABLE 40.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, BRIDGEPORT.

Report of ARTHUR W. Palmer, T. J. Burnell, University of Illinois.

		100000 00000000000000000000000000000000
No. of	Per Per Cubic Centi- meter.	1.982,000 11.682,000 310,000 310,000 3.550,000 5.350,000 1.862,000 3.3550,000 1.260,000 3.3550,000 2.180,000 3.3550,000 3.3550,000 3.3550,000 3.3550,000 3.3500,000 3
	Presence Abs. of C	**+++
911. 311.	Temperat Of Air, Co	
orn;	Pemperator	
	idgi9H rotsW	4 10 4 4 4 4 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 4 6 4 6 4 6 4 6 4 6 4 6 4 6 4 6 4 6 4 6 4
0 - 0 X		ながは、ながない。 ながは、ながない。 ないまりをは、 ないない。 ないまれる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ない。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないまる。 ないな。 ないな。 ないる。 ないる。 ないる。 ないる。 ないる。 ないる。 な。 ないる。 ないる。 な。 な。 な。 な。 な。 な。 な。 な。 な。 な
NITRO-	Soffitti	<u> </u>
KIC .EN.	Sus- pended.	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
ORGANIC	-sid .boylos	
N.	Total.	**************************************
AS.	Sus-pod	21.52.42.82.82.82.82.82.82.82.82.82.82.82.82.82
	W p.pod -sus p.nouiungly p.nou	5+86686659888888888888888888888888888888
NITROGEN	1.10.L	8 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
<i>F</i> -4	Free Am- monia,	$\frac{\pi}{100000000000000000000000000000000000$
ED.	By Suspd	# 804700000000 1000000000000000000000000000
OXYGEN	By Dis-	<u> </u>
000	Total.	#
ne.	ПітоГПО	8.5.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8
Color	on Igni- tion.	後はははははればははははははははははははばばばばばば
·	j pag sus	$\frac{\infty}{\infty} = \frac{1}{2} = \frac{1}$
RATIO	and sid	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
VAPO	1,10,1,	
KESIDUE ON EVAPORATION.	bengeg gus-	ెక్కాని చిన్నాడు. మాడు జెక్కాని చెక్కాని చేశాని చేసిన చేశాని చేసిన చేశాని చేసిన చేసిన చేసిన చేసిన చేసిన చేసిన మాట్లు మండు మండు మండు మండు మండు మండు మండు మండ
IDLE	Dis-	24 458. 28 47 8 12 12 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
KES	Total	28. 1
•:	юрО	Gas.y.
	j.,	ではずれずれずれずれずれずれずれ ですれずれずれ 変さでは 本質 第9 8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
ANCE	Color.	NN NNHHAMANNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
APPEARANCE.	Sedi- ment.	Consid
1.	Z E	
		~===88-=84-=842======8-===8
H 0 B	foo fixami- nation.	Mar
DATEOF	Pad 1900 Collec-Exami- tion, nation,	의는 문항원 는 도로왕 등 다르윈 등 이 문항 는 조 등 다양 본 의 등 다 약 는
	E. 3.2	604 Jan. 604
No.	hiros	88 2 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8

Color on Jantion-G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish, W., White VM., Very Muddy, T., Turbid, C., Cloudy, Coton M., Muddy. Trubibiry—* Decided. § Very Decided. * Distinct. ‡ Very Slight - Slight.

TABLE 41.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT

SANITARY WATER ANALYSIS-PARTS PER MILLION

Report of Arthur W. Palmer.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LOCKPORT

Albuminoid Am.

sid b'vlos

['10']

Free Am By Suspd Matter.

By Dis-Total.

NITROGEN AS

AMMONIA.

CONSUMED. OXYGEN

Color

EVAPORATION

RESIDUE ON

APPEARANCE.

DATEOF

.0%

Chlorine.

0

Loss on 1g'n.

Igni-tion.

p.pud -sns

1,10,1

рәриәд

-sns

Dis. solved.

Total.

Color.

Sedi-ment.

Turb'y,

Examination. 1900

Collection. 1900

Serial

* Cons'd

6608 Jan.

.robO

T. J. BURRILL, University of Illinois,	iture r, C. ture Jent. e or Coli.	Dis- solved. Surates. Zitrates. Wate Tempers of Wate Of Wate Of Nate Of Nat	84 3.68 .002 .64 0 +	44 2.4 .001 .24 0	3.30 3.6 015 2 4.5 4.5 4.5 4.0 000 000 000 000 000 000 000 000 000	24 2.56 .002 .2	992 3.158 .017 .16 9.5	3.456 075 84 0 3.5 5 9	52 2.24 .15 .72 .	6 1.84 .17 .4 0 3,700,	32 1 99 003 19 0 835 H	368 2. 112. 004. 24	1.76 .004 .2 12. 21. + 3.	0.4 .003 .2 15.5 25. + 2,	1.64 42 15.0 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.	1.38 .003 .08 15.5 26.5 + 1	.96 .02 .4 19. 20.5 + 1,	1.04	64 12 28 +	2.34 .002 .2 22. 31. — 750.	.72 .002 .12 18 1.1	.00.4	100 F81	161	22.208.001.12	1.444 + 180.	370,	- 0	1010 - 10
		solved.	.84 3.68	4.2.4. 2.4.	68 3.6	.24 2.56	992 3. 168	184 3, 456	52 2.24	6 1.84	32 72 73	368 2, 112	1.76		- T - C -	1.38	96:	5:-	: : : : : : : : : : : : : : : : : : :	3.34	.73	.336	000	1	2 2.208	1.44	572 2.748	_ 0	OFO 121 E.
ORT.	AS OR	Sus-sugardardardardardardardardardardardardarda	525	- 1.0g	1.2 50.4	8.4	91.7	5 5	91.	4.44	2.00 2.00 2.00 2.00 3.00	33.48	90	928 3.14	in	3	.544 2.18 1	35 C	464 2.74 2	က	1.46	- 66. - 66.	% x	2.52	9	-		27.04	

M'sty Gas'y

Cons'd

: *

May ...

Gas'y

3 * ...
10 + Little
17 * Cons'd
24 * Much

9 9

7217 Apr. 7272 ... 7328 ... 7338 ... 7438 May

Apr.

* Cons'd

G.GGG.000.8 C., Cloudy. ++2. | | 2. T.,.Turbid. + 4 8 8 5 4 8 8 8 V.M., Very Muddy. 3.70f .496 1.104 2.14 .352 2.608 6.04 3 Coron-M., Muddy. 352 2 7 513.6 444.4 69.2 59.6 39.2 20.4 110.4 376.4 601. 13.6 28.8 114.8 535.6 468.8 67.6 60.4 21.6 38.8 114.8 515.2 230. 315.2 330. 315.2 TURBIDITY-*Decided. & Very Decided. + Distinct. + Very Slight. | Slight. M. F.2 Black Much 1 Oct. " 8594 Oct.

Color on Joniton-G., Gray. B., Brown. DB. Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White.

170,000

M'sty Gas'y

Much Littl

* Cons.d

* *

Sept

8368 Sept.

8436

6 Aug

1 1 1

8113 Aug. 8173 ... 8233 ...

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS PARTS PER MILLION.

SOURCE OF WATER-DES PLAINES RIVER, LOCKPORT.

Report of Arthur W. Palmer, T. J. Burnell, University of Illinois.

No. of	Cubic	meter.	1.500	30,000	156.000	18.200	90.5	000.031	120,000	95.000	62,000	15,000	6.000	2.500	4.000		10.600	066.4	3.400	138,700	17,550	0.000	2.900	10 01 20 01
G OL	jo 's	Pre-						-								:	_		7					: :
orn ii onto	npera Air, C	nell' Lio				œ	(C) (C)	10 42			33.57	10.	÷ .	÷.	ig.	:	÷;	2	25		??	15 15 10	33	
ture (!.	eregi 91871	Ten	0.					: c						SC.		: !	9					:	:	
10 1	dglə ətrW		-	: -										:	:	:	:	:	:	30.5	24.	20.5	30.5	्राः
TEO- GEN AS	ates.	ntiN	1.52	ਕ ਟ -	00	9.4	1.13	20.00	23	× +	35	.61	$\frac{\infty}{2}$.56	. 16		œ :	91.	X	÷	-	.16	80.	8 5
NITEO- GEN	.səti	Nitr	.007	8.8	015	.007	0.00	20.0	30.	.011	.022	.046	.015	+10.	X00.	2.3	÷.	00:	-	.003	000.	.003	30.	19 :
IC SN.	ged.																							.064
ORGANIC NITROGEN.	red.		ł																					5.5
0.7	fal.	o,L	1.48	36 25 25	00	96:	92.	5.50		1.176	1.2x	. Sec	186	186	33	3	1.16	- (x.	1961	oc.	.692	119	908.
ov.	< ₹	pud	.113	910	190	.032	.016	080	.176	.113	. (332)	.048	80.	. 192	200.		91.	100	OKW.	.03	-1	. 116	HH).	190
JEN AS	Albuminoid T	Dis	+	?} ?!	288	.384	30	586	200	108.	301	33	:33	333		.435	.33	C T	.32	-	282	33:	108:	25 E.
NITROGEN	T.	10.L	515.	256	352	.416		36.5	100	.416	3336	.368	+	513	512	976.	oc ;	27.0	9	1	133	136	368	± 55 55 55 55 55 55 55 55 55 55 55 55 55
Z	-mv e		.36	048 0.48	£15.	77	- 1	956	36	<u>∞</u>	- 1	80.	0.0	20	210.	X 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	200 200 200 200 200 200 200 200 200 200	1.	5	30%	<u>x</u>	:G)	.053	E. 8.
ED.	ter.	S VS Just	.:. .:.	7:	-	1-	-:	: 5	3.6	+	9] 20		0.0	93	1-0	250	20 1	-	**	-	-	~?	e;	نان
ONYGEN	-si(By I	<u>5</u>	(0 -1 01				0 1-	10.3	9.4	? ? G:	9.4	C: J:U	0.1		7 ? 50 ?	9	T. 0	5.	x.	œ.	? ? C:	<u>с</u> .	= G
0,0	tal.	oT		ना २) भा २-																				= œ. €.∞.
ou-	hlori	(a)	15.5	10.1	8.4	ن نن	G. 5	. .	÷ ??	∞ ≎1	23	-	1-		9.6	57 20 10	- 0	20.00	. C.	g; g	6.	10.3	с:	ر م م پ
Color	lgmi-		~	ಶಷ	В.	<u>~</u>		÷ ~	~	<u>~</u> :	<u></u>	~		<u></u>	::	<u>:</u> ;	-	- :	: ::		-	~		<u></u>
2.	[🖺]	pud	13	: 7						6.4	6.	9.1	6.4	*	7.0	ic o						}	36.8	-01
ATIO	on lg	Alos DIS	10.8	51.4	23.6	36.8	6.5	2 00	??	17.2	1.4	x.	.96	19.2	25.00	× + 50	7.00	27.0	0.6	36.	37.0	5.	17.6	8 18 8 18 9 18
TAPOI	Loss	10.T	62.8	+∞. ±∞.	26.4	÷ ; ;	£ :	1 2 2 2 2	17.3	23.6	30.4	34.4	32.+	53.6	57.6	5.5.0	. e	2.5	C .	200	×.	.00	-	33.55
RESIDUE ON EVAPORATION.	geg.	uəd 15	28.8	= °≥	æ. 33	5.2	9.6	13.4	300					-	5. S	1.2.	2 10	2.2	20.00	29.6	£	€ - -	ည သ	0. E
DCE	is-		396.8	######################################	379.2	242.	2.000.00	200	14:4	138.4	-13	238.±	306.4	368.1.354.4	100 TO	+09.6.581.2	\$49.6	1	11:5	- + + · · ·	340.4	356.4	2000 x	379.6
RESI	fal.	o'I	19. C	1 20 0 1 1 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1		257 2	100	0 20	170.1.142	258.8 138	97.6	25.5	533 6	208	10.0	13.9.6 10.00	250 C 250	2 6	F18.2 550	37 1	357.6 340	367.6 356	201.8	388.8
.1	tobO				:	:					:			:	:	:				:	:		. T. t.	
CE.	Color		1.3	F.F.		に 注:					?? 						05.1	1.7				٠) ا	<u></u>	
ARAN			()			1. X	7.	7.7	N	N.	7.	7.	7	*	-	7, 7	7. 7	7. 7	-	7.	7	7. 7	7.	F
APPEARANCE	Sedi-	nent	Littl	: :	9.9	p. suo.	11117	suo,	11	9.0	:	:	: .	Little		T tak	ויוננונ	n siio i			Little	9		: :
	· A, q.		+	7	+	*	+ +	- #	*	*	*	*	*	ļ	F- 31		4	F 9		+ -	+	+ -	++:	* *
OF	1900 1900	nation.	Jan. 3	: :	:	: · ·	dor c	: 13	08	÷ ;	vpril 3				Nay 1		91 ::			une 5	:	-	: '	uly .
DATEOF	00	9.0	\$15	ος <u>ι</u> Ω	8	£1:	0 1	9	19	56	1 2 1	S.	16			Ci	3	27 5			12	X	1	63 CP
pand	1900	tion.	Jan.	: :	*	: ;	Feb.		:	:	.April	:	*	9 .		VIA.	e 14	7.7		June	4 9	. 7	: '	ylaly:
.0%	.o.X Iniros			6223	67.37	元につ	CT3	3501	2112				0000	5000				0201	1022	1655	1011	1740	Q -	7895 7895 1

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. RG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White, VM., Very Muddy, T., Turbid, C., Cloudy, Colou -M., Muddy. TURBIDITY-* Decided. § Very Decided. * Distinct. ‡ Very Slight. | Slight.

TABLE 43.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PA

SOURCE OF WATER-DRAINAGE CHANNEL,

Color On

RESIDUE ON EVAPORATION.

APPEARANCE.

DATE OF

Igni-tion.

p,pud -sng

I'10T bended -sns

Dis-

Total.

Color.

Sedi-ment.

Larp, A:

Collec-Examination. 1900

Serial No.

tion. 1900

* +++=*

Odor.

Chlorine.

Loss on 1g'n.

ier, f Illinois.	No. of Bac.	per	Centi- meter.			335.000 680.000	695,000		520,000	1,890,000	900.000	2,620,000	80,000	340,000	250,000	415,000	965,000	000,000;	260.000	545,000	120,000	000	2,515,000 400,000	395,000	355,000	
PALN L, ty of	cor.	ogu JC	Prese					:		:	:		:	:		:	:	:			:	:	:		:	
Report of Arthur W. Palmer, T. J. Burrill, University of III	ture ent.	era () (,	qm9T ni A lo					:		:	:		:	:			:	:			:	:			:	
TILL J. I.	ture r, C.	era ete	qm9T					:	: :	:	:		:	:	: ::	23.	333 333	: : :	3 65	23.	24.		o 당하	15. 15.	10.	
of A1	lo 1 r.	en Ep	I9H SW		:			:	: :	:	:		:	:			:	:		:	:	:			:	
port o	FRO- GEN AS	°S6	Nitrati	8.1	5j.	20,	<u>21</u> .	ક્ <u>રં ત</u>	88	83	30.	01.63	82.	: : :	ં	.16	91.	ol .	.16	.16	5.	٠; <u>-</u>	0 :9			
Rej	NITRO-GEN	·se	Nitrite	8.8		S	00.	88				3 :		:3				3 8			.005	3.8	38		.01 <u>2</u>	
	IC SN.	.ba	sng	673	. 148	: 1	92.	.288		.33	3:	476	.496	≈. 16 419	1.116	£.	<u>.</u> .	.892	? ? }	₹.	.295	000	070	. 28-1	.332	
	ORGANIC NITROGEN		-siU 9vIos	54.	.452	.54	3	.392 6	.52	44.	217.	286	504	378	284	.22	رة د قري	2002	88	.22	328	904	.004		.33	
	Oy	1.	втоТ	_	9.			∞ ⊙ ∝	_		-	_	- 1	2.52	-			01.1 %	.9.	.63	33	1.4	3.1.212	.62	.652	
PORT.	on.	Am.	p.pud -sng	.504	960	.032	.128	91.	304	960.	000.	128	960	× × × × × × × × × × × × × × × × × × ×	512	272.	929	.652	199	80.	.224	432	528	.48	224	
Lock	EN A	Albuminoid	Dis-	.176	372	176	.144	000	176	192	100	176	.176	16	176	.112	<u>211.</u>	8 5	208	.128	.16	208	134	808	960.	
MILI DAM,	NITROGEN AS AMMONIA.	Albur	Tiot	89:5:	.368	208	272	36.5	<u>∞</u>	288	300	301	272	368	889	.38.1	.368	355	· -	308	.381	64	652	.688	.32	
IS-PARTS PER MILLION. NEL, BEAR TRAP DAM, LOCKPORT.	A		Free A sinom	2.08	1.53	3 0	33	1.36	1.2	9.1	200	1.93	1.36	2000	· 67	1.84	न हो हो हो	0 00	1.336	.112	82.1	200	52.00	00:	1.36	
TTS AR T	ED.	bq	By Sus	63	3.1		4.1	- 1	ા	ر دن دن	200	0.00	œ:	ာ လ ဘ	8.6	55	1-1	- i-	: =:	က	დ. დ.	Σ. :	5.6	05 05	1.7	
PAE BE	OXYGEN		By Dis	10 cc	ن -	·	5.0	2 5	5.4	ر ان ان	ب بر ي د	0 01	∞ . ∞ .	7 Y.	5.1	3.4	1	; 7	3.9	က	m ,	5.6	1.1.	∞.	+.+	
IS-	0.00	.[Tota	5.5	7.7	5 5	5.1	10 IC	5.6	100	0.3	6.0	4.6	_ · ∞	4.9	<u></u>	}- ?	715		6.	6.9	7:11	0 0	0.6	6.1	

272.4 255.2 1 176.8 1

20.5 20.0 20.0

HHHHHHHHHHHHHH

++**

Aug.

8082 Aug. 8114 ... 8136 ... 8155 ...

Much Little

8023 8058 8058

Foul Gas'y

2888888 2888888

<u>eurrururururaa 4-84-5-165-005-000</u>

COLOR ON IGHTION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. T., Turbid. C., Cloudy. VM., Very Muddy. Color-M., Muddy. Slight. Tunnintry-* Decided. § Very Decided. † Distinct. ‡ Very Slight.

2,480,000 360,600 2,070,000 620,000 120,000

70,000

∞<u>ೱ</u>ౙౙ౾౾౾

8888888

50 00 10 1 0 0 00 10

23.6 24. 1.6 24. 1.8 16. 1.3 2 16.4

199.2 168.4 168.4 168.8 177.2 186.8 178.8 168.8 168.8 168.8 163.6

Oct.

8615 Oct. 8627 ...

M'sty

VLittle

०० मा गुरुश म

6.1

198. 180. 200. 197. 216.4 215. 176.8 146.

9 9 9

Little Cons'd Little Cons'd

M'sty

F. 4 F. 01

Sept.

TABLE 44.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

SOURCE OF WATER-DESPLAINES RIVER, NORTH OF JACKSON ST., JOLIET.

No. of Isac.	Centi- meter.	1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.255.000 1.256.000
	Presenc	++++++++++++++++++++++++++++++++++++++
	Tempera of Air, C	e 0 c a a - u u - u u u u 5 5 5 2 2 2 5 2 5 2 5 2 5 2 5 2
	Tempera	88 - 35 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
1,16	Heigh oran	
A Sc	Zitrates.	$\frac{-6 \cdot 5 \cdot 6}{2} = \frac{1}{2} \cdot \frac{1}{$
NITRO-	Nitrites.	\$
	-sns	& 5655555
ORGANIC	solved.	85888888888888888888888888888888888888
ORG	.sid	
	Total.	$\frac{1}{2} \frac{1}{2} \frac{1}$
AS	id sug	
ITROGEN	Albumimoid fig. Dis-	8 8 2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
NITROGEN	Tot'l	8
A	Free Am- monia.	$\frac{16204}{1620} + \frac{1}{2} + \frac{1}{2}$
N SD.	By Suspd	$\frac{\partial F \otimes \mathcal{O} \otimes$
OXYGEN	By Dis- solved,	$\frac{499}{2} \overline{v} = \frac{1}{100} \frac{1}{10$
Cos	Total.	్షాప్యాన్
.əu	Chlori	5元 & 後 x 3 2 4 4 5 5 5 5 8 8 4 8 5 8 4 8 8 6 8 8 6 8 8 6 8 8 6 8 8 6 8 8 6 8 8 6 8 8 6 8
Color	on gni- ion.	最近我我我我就就就就就就就就就就就就就就就就就就就就就就就就就就就就就就就就就
ŏ	— 4	$\frac{1}{2}$
10N.	grad p,vlos	్యాండికి మాగా నాలా పాటా చెన్నారు. మార్జులు మార్
ORAT	Loss on 1g	1000 + 20 - 100 +
SYAP		
ON I	Sus	
RESIDUE ON EVAPORATION.	PI2-	2010 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
RESI	Total.	25. 2
	tobO	Gas y Cas y Navy Navy Cas y Navy Navy Cas y Navy
ri.	or.	######################################
APPEARANCE.	Color	******************
PEAR	Tit.	Cons'd Little Cons'd Little Cons'd Little Cons'd Little Little Cons'd Little Little Little Cons'd Little
	Sedi- ment.	
	<u> </u>	
10 P	1900 1900 Collect Exami- tion, nation.	THE TO SEE THE TENT OF THE TEN
DATE OF	lec-	กระเมลายนายสาย และสมอนาสาย และ
	Collection.	888 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
.0%	Serlal	<u> इ.स.च.च.च.च.च.च.च.च.च.च.च.च.च.च.च.च.च.च.</u>

Color on Lantion C. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown, BG., Brownish Gray. Bft., Brownish. R., Red. Rh., Reddish. W., White. Colon - M., Muddy, V.M., Very Muddy, P., Turbid, C., Cloudy, Terrinty * Decided. Nery Decided. + Distinct. + Very Slight. Slight.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-KANNAKEE RIVER, WILMINGTON.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

No. of Bac.	Cubic Centi- meter.	10000 10000
	Presenc	1+++ : ; +++++++++++++++++++++++++++++++
	Tempera of Air, to	
	Tempers	000-0 10-000000000000000000000000000000
	dgiəH ots7/	
TRO- GEN AS	Zitrates.	**************************************
NITRO- GEN	Zitrites.	
S. S.	Sus- pended.	\$\$\\ \frac{1}{2}\\ \frac{1}\\ \frac{1}{2}\\ \frac{1}\\ \frac{1}\\ \frac{1}\\ \frac{1}{2}\\ \frac{1}\\ \frac{1}\\ \frac{1}\\ \frac{1}\\ 1
ORGANIC NITROGEN	Pis- solved.	\$ \$ 2 \frac{1}{2} \infty \frac{1}{2} \infty \frac{1}{2} \frac{1}{2
OBO	Total,	2 2 5 2 3 2 3 2 3 5 5 5 7 5 8 2 5 2 5 2 5 2 5 5 5 5 7 5 5 7 5 7 5 7 5
	i p.pud	25
87 .	Dis- biold b'vlos	8 + 8 8 8 8 8 8 8 8 8 8 7 8 7 7 8 7 8 7
ITROGEN	Albuminoid Pistoria Divios	<u></u>
Nr	.sinom	91
D.	Matter.	<u> </u>
OXYGEN	solved.	$\frac{4-1}{4} + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + $
OX	Total.	$\frac{4}{4} \frac{1}{2} \frac{1}$
ne.	СЫОгі	နားသည္ကေတာ့ သည္တည္ မေတြ မိုသည္က သည္တိုင္းတြင္း သည္တိုင္း သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္
Color	on gni- ion.	####################################
	ln nud	8 8
TION	Foss on 1g, n. Sus- Diss on 1g, n. Sus- Diss on 1g, n. Sus-	8546288858888888888888888888888888888888
POR	Loss 1.10 T	
RESIDUE ON EVAPORATION	pended.	100 00 00 00 00 00 00 00 00 00 00 00 00
E ON	solved.	10 m 04
ESIDI	-std	25
2	Total.	2
	торО	
53	or.	##
APPEARANCE	Color	kakakakakakakakakakakakakakakakakakaka
PPEA	Sedi- ment.	Little Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Little Li
7.	Turb'y.	
	1	######################################
DATEOF	1900 Exami- nation.	23
DAC	1900 Collec- tion.	Jan. 15 12 12 12 13 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15
.01.	Serial	6600 [19] 6601 [19] 6601 [19] 6601 [19] 6601 [19] 6602 [19] 6603 [19] 6003 [
		XXXXXXXXXXXXXX

Color on Ighthon-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 46.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS—PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, MORRIS.

No. of Isac.	per Cubic Centi- meter.	286.000 1.550.000 2.550.000 2.550.000 3.650.000 3.650.000 3.650.000 3.650.000 4.5000
6 or ('olf.	Presence To set A	· + + : : · · · · · · · · ·
	Tempera Of Air, C	- 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
'J. 'J.	Tempera of Wate	0000 000000 H0000000000000000000000000
1.	HgioH ots7/	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
AS	Nitrates.	※主 主が始め発展器 が窓がる※主送干のお子登録ではおよばに対信主写を発表表表表
NITRO- GEN	Nitrites.	$r^{\prime\prime}$ $g_{1}^{\prime}g_{2}^{\prime}g_{2}^{\prime}g_{3}^{\prime}g_{3}^{\prime}g_{2}^{\prime}g_{2}^{\prime}g_{3}^{\prime}g_{3}^{\prime}g_{3}^{\prime}g_{4}^{\prime}g$
οż	pended.	<u> </u>
ORGANIC NITROGEN	Dis.	8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
NO TIN	Total.	28.8.2.8.2.2.2.2.8.8.8.8.8.8.8.8.8.8.8.
	i p.pud	2
SN AS		\$\frac{\pi}{2}\$\
NITROGEN AMMONIA	Albuminoid Dis- Div'de	86 4 4 4 4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Nr	mohia.	8 + 5 4 9 5 8 8 8 2 5 5 5 4 8 4 8 8 4 5 8 5 2 5 2 5 2 5 2 5 2 5 2 5 2 5 2 5 2
. o	By Suspd Matter.	$\frac{4\phi_{1}\psi_{1}+-\psi_{2}\psi_{2}}{\phi_{3}\psi_{2}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}} = \frac{4\phi_{1}\psi_{1}+-\psi_{2}\psi_{2}}{\phi_{3}\psi_{3}+\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3}+\psi_{3$
OXTGEN CONSUMED.	By Dis.	<u> </u>
Con	Total.	కృథానికింగాలో అంది. స్టార్జ్ కాయలు గాగా అలగా అంది అందు అంగా అంది అంది. అంది అంది అంది అంది అంది అంది అంది అంది
.əu	Chlorin	<u> </u>
Color	on Igni- tion.	
	p,pud	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
ATION	Piori Los on Los	8-6-48-48-68-68-68-68-68-88-88-88-88-88-88-88-88
APOR	Lorr Lorr	
N EV	pended.	※ 1 × 2 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3
RESIDUE ON EVAPORATION.	.bevios	4
ESID	-sid	0.
bring	Total.	
	TobO	Gasy M'sty M'sty Gas'y
5.0	lor.	
RANC	Color	sakasakaka kakakakakakakakeeeeee
APPEARANCE	Sedf.	Cons.d Little Cons.d Little
	Turb'y.	** + * * + * * * * + * + + + + + + + * + * * * * + + + * * * * * * + + * * * * * * * + + *
5.		12 13 14 14 15 15 15 15 15 15
DATEOF	3 3	4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Da	1900 folled tron.	Mar
o.o.N	Iniros	88 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
		2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

Color on Javeton-G. Gray. B. Brown. DB. Dark Brown. LB. Light Brown. RB. Reddlsh Brown. BG. Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy, VW, Very Muddy, T., Turbid, C., Cloudy, Terbinety * Decided. § Very Decided. + Distinct. + Very Slight. Slight.

TABLE 47.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

A
h. 1
-9
H
-
\circ
0.0
ER,
- 23
-
-
40
8
~
OX
-
200
2
2.5
120
-
4
. '
=
-
E.
1
0
-
31
-
RCE
20.00
-
\supset
-
0

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

		0	00		*	. .	÷	00	عا د		0	٦	0 1	- :		0	0 :	D	. c	0	0	0		0	-	٠ ت	-	00	5 0
No. of Bac.	per Cubic Centi- meter.	3.35	(€. 3 (€. 3 (€. 3 (€. 3 (€. 4 (€. 4 () () () () () () () () () () () () () (300.55 500.55 500.55	150,00	16.50	245.00	12.50	000000000000000000000000000000000000000	2.50 2.50 2.50 2.50	18,20	1.45	Ž.	20.0		3.95	Ø: : :	5)+'2	06.5	£.30	1,00	2,70		18.50 18.50	20.12	1.50 1.50 1.50 1.50 1.50 1.50 1.50 1.50		S	4.70
	Presenc	+	++	-	+	+	-				+	+		-	++	+	:	ı	+	-	۸.	1	:	a .		ī			1 1
Temperature of Air, Cent.		ાં	ei e		٠ ښو	i }-	16.	Ξ,	· · ·		6.	3	19.		20. 13. 13.	3	:	:		:	:	:	:	:		:	:	:	
Temperature of Water, C.		65	တ် က	, -	د ي					: .		-			ु हा		26.	 G	200	23.		 .:	: 1		5.5	65	0.5	2 i z	
Height of Water.		62	ස ව ව	- - -	÷ -	· -	+	- 11							000			0.2	2.6	30 24	<u>ം.</u>	3.6		9.7		3.6	- · ·	300	0.0
O- NA AS	Zitrates.	.56	हां ह <u>ै</u>	253	20	. 33 . 33 . 33	9.1	1.16	2 8	ું હાં	.36	ਰ.	23.8	250	7. 6.	.36	91.	9:	: :	80.	. 16	82.	83.	.; 	35	85	91.	: : :	988
NITEO-GEN	Zitrites, S			0.00									-	•												.003		.003	.005
2 c	·pəpuəd ·sns	.16	208.9	.208	25.5	.608	1.408	.416	57.5	48	.548	.332	388	212.	001.	124	141	25.		.176	272	880.	584	.736	. 10x	.028	899	:	128
ORGANIC NITROGEN.	Dis-	.52	56. S.	673	000 000 000 000 000 000 000 000 000 00	632	795	.664	0.00	708	.612	.568	£:	400	320	966.	308	55.	2 .	.388	.561	.46	212	<u></u> چو		.576	.673	:	513
ORIN	Total.	89.	200	- x	# ā	ंक	e	 80 80 80 80 80 80 80 80 80 80 80 80 80	368	88.	.16	<u>.</u> ص	86	8000	152	£.	1000	564	548	.564	.836	.548	2.5	.116		1 09.	: 5 .	- 188 - 188	 .6.1.9.
	i p.pud						<u></u> 5.				_										<u>. </u>			=			=		010
A .																									_				116
NITROGEN	Albuminoid 1.10T																												\$ 5
Nir	monia.						band																						142 .4
	Matter. Free Am-	0. 4	200	e 60 :	ان فرن بو فرن	, eq.	<u> </u>		5 C	0	<u>හ</u>	0.				0.	0.0	2 5	7	1 .0	5	-	~ ·	2		9	<u>ح</u>	5	2. 1.
OXTGEN	By Suspd	च्	ن تن	9.	<u> </u>	- 77	+ 1 + ·	,	000		ಣ				- 0 - ∞	0	.0	<u> </u>	. T.	-	9.	ا	7. 1.	90	χ χ	€	9.	χ; :	1
OXTGEN	Total. By Dis-	9.8	ن ت بر بر	· ₹~ :	φ φ γ	0.1.0	3.5	. 4	<u>و</u> و	200	<u></u>	6.	7.0	n. c	0 ∞ 0 ∞	5.5 6	₹~ °	ວ ຜ : ;.	9	9.3	2.1.7	6.7	8 .0 6 .	න න	 	000	2-	5.3	3.8
.ani	Сріогі	6:		re c	χ. :3	-	œ ∕	2) α —	0	 	.6	9.	ઓ . 	_	ەت تە	9.	7 6	3 æ	2 7	9:	<u>=</u> ः	rċ.		? ·	7	<u>∞</u>		7:0	0 9
H	Igni- tion.		D (-	- 71	ri en	- 		कश का 			4	9						9 6			9			 	<u>ज</u>		ر چيد ا		ы Б.:
_	1 1 1				r r				H 00	<u> </u>	න 		<u>क</u> ्र			음 국	<u>جر</u> د	0 ==	7	3 3	~ :	음 : 다	<u>क</u>	ψ i	¥:	es es	φ 	27 F	1 Bh
NOI.	I'10T Solution l'10T o service l'10T b'Vlos -suS -suS -suS -suS -suS -suS	40	9	00 t	00 0 00 77 77	·	?}	ە - -			30. 2.	구 ? }		40.0	o 0 01	-		95.0			:	3) (-)		∞		00	∞ 23	6.5	6 10
ORAT	sso -sid		5. 7. 5. 6. 7. 6.	38	5. 5. 5. 5.	36.		हरू हैं कर य	2 20	4	-	00			+ 9	00	000	H 60			•	100 H	9			3	.6 3	∞. «	6 8.
25		81.2	9. 6		+; ±		20	<u>ත ද්</u>	3 % +		33	33		0.14.0	7.5	36		0.22		45				.6 41					
NO 3	Sus. Sus.	10	= -	* ## ·	30	108.		25.5	2 9	52.8		15.6			3.6.6									3 173.6					∞.
SIDUI	Dis- solved,	150			7.09.1 0.00.1			199.2				_	986	282.0	25 25 26 25 30 50	293.2		388.		263.			_				367.6	280.4	288.8 291 6
RE	Total.	160.	318.8	202.4	202.8	281.6	626.8	7 986 386	2000	302.8	405.8	300.8	291.	2588.2	350.0 350.0 4.0 4.0 5.0 5.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7	301.2	321.2	304. 313.6	316.	305.	358.	8.4%	301.2	440.4	297.6	304.4	27.1.2	296.	304. 299.6
r,	opO		:		:	: :	:	:	:		:	:	:	:			:	:					SI.Gs.		:	:	:	:	
	01.	F.03	F. 69	3.15	500 200 200 200	888	F.30	F. 15	3.	F.06	F.04	1.1	F.05	F.02	300	F.01	王 35.85 8.85 8.85 8.85 8.85 8.85 8.85 8.8	F.20	五 30.F	F.02	F.02			E.02	F.05	ડ -	F.01	F.05	F.2
ANCE	Color	M. F		M. I	Z.	- I		N. H.	17	N			N.		7. 7.				. T.									-	
APPEARANCE	Sedi. ment.	Little	: :	Cons'd	Much	Much	9.9	Cons'd	Much	Cons'd	;		Cons'd	Fittle		Much	Little	77	9,9	Much	Little	33	99	= 1	Cons'd	Little	"	9, 3	,,
Turb'y.		+	==+ ==+		* *	*	*	* *	+ + 50	- * - *	* 01	<u>}-</u>	+ -	- +	500	*	* 1	e ÷	*	*	* 6	*	*	*	+	+	-	100	-++
E 0 F	1900 Exami- nation.	Jan.	3 3	7.9	Mar.		Apr.	: :	: :	May	:	:	:	June	: :	9.9	July	: :	9 4	*,	Aug.	4			Sept.	,,			
DATEOF	1900 Collection.	Jan. 4	:::	:	Mar. 12	ः	Apr. 3	::) T	May 1		15	3.0	c oung	: :	** **	July 2		. 53	30	g. 6	13	3000	3	Sept. 4	.: 10	17	FG	Oct. 1
Serial No.		6629	6678	6772		919		7288	075-1		7.489	7517	7599	7670	2017	7786	7820	1000	7096	8047		8171	% %	85396	%399 %399 %399 %399 %399 %399 %399 %399	%1333 %1333	£		8635 5635 5

Color on Ightion-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. VM., Very Muddy. T., Turbid. C., Cloudy. Colon-M., Muddy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS. PARTS PER MILLION.

T,
-
-
1
7.5
\vdash
-
_
- 5
pro-pr
1.0
p.liq
partie.
-
~~
-
5.
-
0
11
-
-1
_
_
40
ponq
122
<u>-</u>
-1
4
2
2
2
-
-
-
OF W.
OF W.
E OF W.
CE OF W.
CE OF W.
TICE OF W.
BCE OF W.
DUBGE OF W.
DUBGE OF W.
TICE OF W.
DUBGE OF W.
DUBGE OF W.
DUBGE OF W.
DUBGE OF W.

Report of Arrhub W. Palmer, T. J. Burrhull, University of Illinois.

No. of Bac.	2000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
resence or describe of the	
hio') , ri /. li '. hio') , ri /. l	
omberadine Chater, C	
Height of Water,	
GEN AS	X 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7
NITRO- GEN GEN	
Susans	
ORGANIC NITROGEN Dized	
o'Y .Isto'l	第4章第※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※※******
id Am.	#
	o.
Nitrogen Ammonia Albumino	T 8 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
ve Am-	
Suspd Bass	(a) and and a to employed the and all and a supplied the
ONYGEN ONYGEN ONYGEN ONYGEN ONYGEN ONYGEN ONYGEN ONYGEN ONYGEN	
Chlorine,	\$\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
Color on Igni- tion.	
No. 18 19 19 19 19 19 19 19 19 19 19 19 19 19	A
PORATI	
EVAP	n, %
RESIDUE ON EVAPORATION DISSEMBLE ON EVAPORATION DISSEMBLE ON EVAPORATION DISSEMBLE ON EVAPORATION	
Dived.	
S Jeto'	######################################
.tobO	N N N N N N N N N N N N N N N N N N N
or.	FF9889898989898
RANCE.	katatatata zakakakakatatatatatatatatatatatatatatata
APPEARANCE.	Cons. d Little
्रं,वृज्ञ	.1, + + + * + * + * + + + + + + + + + + +
OF 1900	
E	
oallo)	Nar
Jos Juines	इ. १ वर्ष दे हे है है है है इ.

TUBRIDITY *Decided. § Very Decided. * Distinct. ‡ Very Slight. Slight. (older M. Middy. VM., Very Muddy. T., Turbid. C., Clondy. (older On Leveridea, C., Cray, B., Brown, DE., Dark Brown, LE., Light Brown, RE., Reddish Brown, BG., Brownish Gray, Bh., Brownish, R., Red, Rh., Reddish, W., White.

STREAMS EXAMINATION -CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burriel, University of Illinois,

SOURCE OF WATER-BIG VERMILLION RIVER, LA SALLE.

No. of Bac.	per Cubic Centi: meter.	8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Presence or Abs. of Coli.		
Temperature of Air, Cent.		
Temperature of Water, C.		
Height of Water,		
NITRO- GENAS	Nitrates.	0.000
N.	Nitrites.	100 100
NIC	Sus- pended.	yand * yand *
ORGANIC NITROGEN	Dis-	8
	Total.	######################################
S A .	p,pud -sng	00000000000000000000000000000000000000
	Albuminoid Pistoria Divios	56 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
NITROGEN	Toor.	\$\frac{1}{2}\frac{1}\frac{1}{2}\f
Z	Free Am- monia.	5.3448155851588888888895588889595959589535344888754
ED.	By Suspd	-0 r
OXYGEN CONSUMED.	By Dis-	
Total.		$-i\psi ij \omega \omega ij \omega ir ir ir i\omega \omega + i\psi \omega ir ij ir iv iv \omega \omega \omega ir iv iv iv iv iv iv iv$
	Chlori	42.83 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 -
Color	on Igni tion	年代は単元はまなはははははははははははははははははははははばればばばらら
N.	i p,pud sng	$\frac{\omega-4}{\omega}-\frac{\omega}{\omega}\frac{1}{\omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial \omega}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}\frac{\partial}{\partial \omega}$
RATIC	Pivos Pivos bivos bivos -sid -sus -sus	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
VAPO	T.10T	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RESIDUE ON EVAPORATION	bended.	○ 元 元 元 元 元 元 元 元 元 元 元 元 元 元 元 元 元 元 元
DUE	Dis-	88
RES	Total.	2584 - 24 - 24 - 24 - 24 - 24 - 24 - 24 -
Odor.		
	or.	8 9 8 8 8 8 9 8 8 8 9 9 9 8 9 9 9 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9
APPEARANCE.	Color	a zazgazgazzak a kazazzetetetetezak aza
	Sedi- ment.	Little Cons.d Cons.d Much Little Little Cons.d Cons.d Little Little Little Little Little Cons.d Cons.d Cons.d Little Little Little Little Little Little Little
	Turb'y.	word20122-122-122-222-222-222-222-222-222-
EOF	1900 Exami- nation.	Jan. Jan. Nar. Nar. Nar. June June Sept.
DATEOF	1900 Collec- tion.	Jan. 18. 18. 19. 19. 19. 19. 19. 19. 19. 19. 19. 19
,o.Y.	Serial	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8

Color on Ignition-G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bb., Brownish. R., Reddish. W., White Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. # Slight.

TABLE 50.

STREAMS ENAMENATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, LA SALLE.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

		and the second of the second o
No. of	Per Cubic Centi- meter.	23.1.000 111.0
	Presence Abs. of 6	+++++
Temperature of Air, Cent.		
1.1 (918A 10	
Temperature Temperature		
A. S.	Zitrates.	$\mathbb{R}^{3} \times \mathbb{R}^{3} = \mathbb{R}^{3} \times \mathbb{R}^{3} $
NITRO-GEN	Nitrites.	g-2328725252222222222222222222222222222222
	-sus	868888855588848488845888458555555555555
ORGANIC NITROGEN	Dis-	20.8
O.Y	Total.	$\frac{888886}{8888} \times \frac{2}{8} \times \frac{2}{8$
	g p,pud	4 1 2 2 3 4 5 4 5 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5
NITROGEN AS AMMONIA.	M. M	54 5 4 8 8 6 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
AMM	I.10T	空存沒住的契 [∞] 表表的查表的答案的发表示背限的特殊的型式的重要的程度的型性的
4	Free Am- monia.	第5544894644888888888888888888888888888888
ED.	By Suspd	
OXYGEN CONSUMED.	By Dis- solved.	$ \begin{array}{c} \Box \varphi \alpha \varphi \varphi \varphi \varphi \alpha \alpha \alpha \gamma \varphi \alpha \gamma \varphi \alpha \alpha \alpha \alpha \alpha \gamma \varphi \varphi \varphi \varphi$
<u> </u>	Total.	— မြေရန်မှ ထုတ်သွေတွင်ရှိ မြန်မာ့မှ ရေရန်မှာ မြေရန်မှာ မြောင်းများ မြောင်းများများ မြောင်းများများ မြောင်းများများများများများများများများများမျာ
	Chlorit	会後4247774444200000100010000000000000000000
Color	on Igni- tion.	
7	p,pud sps	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RESIDUE ON EVAPORATION.	on 187 D'vios	\$652.857.89.288.87.128.7288.8288.8288.838.652.857.888 4. 4. 4. 4. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6.
APOL	Loss on Said	$\frac{1}{10000000000000000000000000000000000$
N Ev	popuod Sus-	$\frac{\omega}{\omega} \frac{\partial \omega}{\partial \omega} \frac{\partial \omega}{\partial$
UE O	postos	कर रह
ESID	-51(1	
	Total.	
	robO	
ដ្ឋ	Color.	######################################
RANG	၁	
A PPEARANCE.	Sedi- ment.	Cons.d Cons.d Little Little Little Little Cons.d Little Li
	L.J. and	·
) F	19au 1900 Collece Exami- tion, nation.	
DATEOF	, c. E.	- 10 m m m m m m m m m m m m m m m m m m
Q	19au Collection.	Jan. Field Mar.
No.	[6]102	666 Jan 6652 III Aug 8552 III Aug 8553 III A

Color on fantion G. Gray. B. Brown. DB. Dark Brown. LB. Light Brown. RB., Reddish Brown. BG. Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon - M., Muddy, V.M., Very Muddy, T., Turbid, C., Cloudy, TCRRDITY-* Decided. § Very Decided. † Distinct. † Very Slight. Slight.

STREAMS ENAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrul, University of Illinois.

123
_3
3
Э.
-74
S
_4
-4
\vdash
Ŷ
7
-14
Z
-1
53
-
N-pr
7
A P
0
H
5
\equiv
Z
-
_
AND
7.
-1
4
100
-
0
NOIS
Date:
-3
-1
1
04
TER
-
ME.
-
-
0.5
0
23
0
URC
1
2
10

No. of Bac.	cubic Centi- meter.	25,000 24,000 25,000 25,000 26,000 27,500 26,000 27,500
	Presenc	++++* ++ + ++++++++++++++++++++
Juo5	Tempers of Air, C	
	Tempera	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
,16	ngioll etall	
TRO- GEN AS	Zitrates.	$5 \times 8 \times 4 \times 8 = 4 \times 8 \times 4 + 6 \times 8 \times 6 \times 6 \times 6 \times 6 = 4 \times 4 \times 6 \times 6 \times 6 = 6 \times 6 \times 6 \times 6 \times 6 \times 6 \times 6$
NITRO-	Nitrites.	88
	Sus-	84± <u>Egg2</u> 28888888588856888658865666668
ORGANIC NITROGEN.	solved.	25.25.25.25.25.25.25.25.25.25.25.25.25.2
O.N.	Total.	**************************************
	p,pud sns	50000000000000000000000000000000000000
EN AS	Albuminoid Silvid	45,24,89,84,48,148,89,99,99,99,99,88,89,89,89,89,89,89,89
TROGEN	Tol'I	20000000000000000000000000000000000000
N	Free Am- monia.	8
SD.	By Suspi	0.000 0.000
OXYGEN JONSUMED	By Dis-	$-\infty - \omega + \omega - \omega + \omega - \omega + \omega + \omega + \omega + \omega + \omega$
CON	Total.	□ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □ □
не.	Chlori	\$\frac{1}{2}\frac{1}{2
Color	on Igni- Lion.	
	i p,pud	$\frac{-4\pi}{6} \times 9 \times 9 \times 9 + -\pi \times 6 \times $
TION	Solution 1997 1997	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
POR	Tot'l	
RESIDUE ON EVAPORATION	pended.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
CE O	solved.	
RESII	-sid	TH 0101999 000000000000000000000000000000
	lsto'T	
•,	TobO	Nistry Nistry Nistry Nistry Nistry
RANCE.	Color.	<u> </u>
	တိ	azanan janazanan kanan k
APPEARANCE	Sedi- ment.	Cons'd Nuch Cons'd Cons'd Little Cons'd Little Cons'd Slight Little Cons'd Little Cons'd Little Little Little Little Little Little Little Little Little
	Turb'y.	$\frac{1}{1+++++++***+****+***+*+++***********$
OF	1900 Exami- nation.	Jan. Jan. Jan. Mar. Mar. June
DATEOF	1900 1900 Collee- Exami- tion. nation.	Jam. Nav. Apr. : : : : : : : : : : : : : : : : : : :
	-	
.07	Seríal I	6606 6653 6653 6653 6653 6653 6653 6653

TURBIDITY-*Decided. § Very Decided. † Distinct. ‡ Very Slight. † Slight. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Colon on Ichtrion-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Reddish. W., White.

TABLE 52.

STREAMS EXAMINATION CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS PARTS PER MILLION.

SOUTH

and the
2.4
Z
-
-
_
0.0
-
EII.
-
1-1
00
-
10
_
-
0
0
0.00
180
LINO
180
LINO
LLINO
LLINO
-ltiino
-ltiino
R-LLINO
n-lunno
R-LLINO
TER-LLINO
ATER-ILLINO
JATER-ILLINO
JATER-ILLINO
ATER-LLINO
WATER-ILLINO
WATER-ILLING
WATER-ILLING
WATER-ILLINO
OF WATER-ILLING
WATER-ILLING
E OF WATER-ILLINO
E OF WATER-ILLINO

Report of Anthur W. Palmer, T. J. Burrhill. University of Illinois.

No. of Bac.	per Capie Centi- meter.	6.400 103.0
	Presence Ju-sdA	
	Tempera	하는데 이번 부번 때문 근무 때에는 이 전문로 왕 로 만든 경영장
	stoqmoff otaW lo	वर्गन् । वर्गन्त्रम्मन्द्रम्मन्द्रम्भन्द्रम्भन्द्रम्भन्द्रम्
	Heigh ofaW	$\frac{1-2\alpha r}{2} = \frac{1-2\alpha r}{2} = 1-$
N. AS	Nitrates	
NITHO-	Nitrites.	<u>enangangangangangangangangangangangangang</u>
	рориод	24 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
ORGANIC	solved.	\$555778 \$25 \$25 \$25 \$25 \$25 \$25 \$25 \$25 \$25 \$25
ORC	Total.	82°8844585864 <u>2888</u> 2°×888 <u>85566666</u>
	1 • 1	※第5號等號5套子裝訂至數第三至5套第三數單數 2 2 3 2 3 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 5 6 6 7 6 7 6 7 6 7 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 <tr< td=""></tr<>
AS.	K Sus	- 48
TROGEN	Fror Parent Property of Parent Property	
NITE	I'10'T	本党をおよりに対象を記述する *** *** *** *** *** *** *** *** *** *
	-шл ээтЧ лиони	
ED.	By Suspd	
OXYGEN	By Dis-	$\infty \oplus x \times F \oplus \oplus \oplus \oplus \oplus \oplus G \times \infty \times \oplus \times G \times \infty \times F \oplus F \oplus \oplus \oplus \oplus \oplus \oplus G \times G =$
o S	Total.	$\frac{\alpha \alpha \sqrt{3} r_1 x_1 r_2 - \sqrt{2} \sqrt{2} x_1 \sqrt{3} - \alpha \alpha \sqrt{2} - \sqrt{2} \alpha \alpha \sqrt{3} - x_1 x_2 r_2 - \alpha \alpha \alpha \alpha \alpha \sqrt{3} - \alpha $
.91	Chlorit	888
ofor	on gmi- ion.	
	= p,pud sus	
TION	a sid b'vios	8% 8 2 4 8 8 8 5 5 5 7 7 7 5 8 8 8 8 8 8 8 8 8 8
PORS	Tor.	
Eva	Dended	μοι φυρφορούν ο σχακοφοκού οχή πουσκ
NO.	-sus	91
RESIDUE ON EVAPORATION.	Dis-	8
RES	Total.	2
	1000	
	robO	
<u></u>	Color.	STATATATATATATATATA STATATATATATATATATATATATATATATATATATATA
RANG	3	प्रवेद्य प्रवेद्यद्यद्यप्रवेद्य प्रवेद्यद्यद्यम् ।
APPEARANCE	Scali- ment.	Little Little Nuch Cons.d Little
1	-	
	ि हे हैं	crassanspanskansr2-25880588=5727=220580
OF	Lien 1900 Collect Exami- tion, nation.	Tank and the second sec
DATEOF	C.C-11	교통원왕대급교왕대급교왕의교통원왕노편원왕극도국원의교통왕교교왕원교급왕
	Collection.	Jan. Jan. Nar. Nar. Nay. Nay. Nay. Sept.
No.	Listers	### ### ### ### ### ### ### ### ### ##
		X X X X X X X X P P P P P P P P P P P P

Tuberity-*Becided, §Very Becided, † Distinct, ‡ Very Slight. | Slight. | Slight. | Colon | M. Minddy. | V. Very Maddy. | T., Turbid. | C., Clondy. | Colon | C

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE O

.:	
E	
LL	
=	
1	
RYVI	
ERY	
\simeq	
田	
-	
_	
.4	
-	
FER,	
(41)	
5	
RI	
-	
m	
=	
0	
CINOIS	
H	
II	
heet	
1	
200	
rer	
LE	
T	
N	
-	
F	
1	

Report of Arthur W. Palmer, T. J. Burnell, University of Illinois.

No. of Bac.	per Cubic Centi- meter.	28.680 1.38.680 1.38.680 1.38.680 1.38.600
Presence or Abs. of Colf.		11++ +++11+++111
	Tempera O'Air, O	8 1 1 1 1 1 1 1 2 1 1 1 1 1 1 1 1 1 1 1
ture r, C,	Tempera of Wate	
10.1	Heigh ofa'W	24.825
NITRO- GENAS	Nitrates.	
Z.	Nitrites.	5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5
NIC GEN.	-sus	# + * * * * * * * * * * * * * * * * * *
ORGANIC NITROGEN	Dis-	### ### ##############################
-72	Total.	2
sv.	E p,pud	2. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.
	T'10T Solv'd	£15885588888888888888888888888888888888
NITROGEN AMMONI	I'to'T	86.85
	Prec Am- monia.	2
EN MED.	By Suspd	
OXYGEN	By Dis-	∞ with ∞ the ∞
	Total.	$\frac{w}{\varphi \otimes \varphi \otimes \overline{\varphi} \otimes \overline$
	Chloric	
Color	Igni tion	<u> </u>
oN.	S on 1'10'T S'S on 1'10'T S'N'108 S'N'10	0.00
RESIDUE ON EVAPORATION.	ao sid b'y los	+6884848857777777848888877848888888888888
EVAP		83358893666688666666666866866846866666666666
0.N]	-sus-	
SIDUE	Dis.	337.6 337.6
RE	Total.	385.6 35.6 35.6 35.6 35.6 35.6 35.6 35.6 3
	TobO	Wast V
6.1	or.	EXERCETERER EXERTERER EXECUTATION
RANCI	Color	zazak zazazazazazazazazete eletek zazazaz
APPEARANCE.	Sedi- ment.	Consideration of the considera
	S,qan,L	でして当っているようなよーでいっているのでのいまーでによるにこのです。 +++*+∞***+****************************
: 0F	1900 Exami- nation.	20
DATEOF	1900 Collec-1 tion.	- at - att
	757	6603 Jan. 6681 Jan. 6680 Jan. 6681 Jan. 6781 J

TURBIDITY—* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight. | Color On Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White.

STREAMS ENAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, WESLEY CITY.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

o. of	per Cubic Centi: meter.	66.000 113.
Abs. of Coll. Rac. Children Colling Co		
Presence or		- : • : · · · · · · · · · · · · · · · · ·
51111	Tempera of Air, C	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1 ure	BreqmeT'	
JO 1	idgioff ofic77	01-01-4-4-1-1-5-55 - 5 - 5 - 5 - 5 - 5 - 5 - 5 -
TRO-	Nitrates.	8+1355 8865 8-455 205 205 885 + +28 + 855 558 8+ +28 55
NITRO- GEN	Witrites.	年 2 2 3 2 3 2 2 2 2 2 2 2 2 2 2 2 2 2 2
IC EN.	Sus- pended.	88888888888888888888888888888888888888
ORGANIC NITROGEN	Dis- solved.	85.55
ON	Total.	86488652851428888888558848888848848848
on	P.pud esns	84 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
4 ;	Albumimoid 1'10'T sign	######################################
NITROGEN	I'10'T	8. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.
Z'	Free Am-	8 9 9 4 4 5 8 8 8 8 4 4 8 8 5 5 5 4 2 5 8 8 6 5 6 8 6 5 6 8 5 6 5 6 5 6 6 6 6
ZD.	By Suspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN	By Dis-	1-808881-081-01-181-181-181-181-181-181-
CON	Total.	
ne,	Chlori	##88##555000 ##5 # # # # # # # # # # # # # #
Color	on Igni- tion.	長氏性性ははははははははははははははははははははははははははははははははははは
	≓ p,pud	$\frac{14009146796714671467674}{9988499949499899999999999999999999999$
TION	ald sid	######################################
POR	Loss on 18	τ φρτισατι α ααιρταρ τοι στον αραν
RESIDUE ON EVAPORATION	bended.	
E ON	-sug	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
SIDU	·sid	357.12 367.12 367.12 361.8 382.8 382.12 381.12 381.12 381.12 382.12 382.12 382.13 382.13 383.
RE	T'otal.	367 12 370 8 6 16 16 2 38 1 2 370 8 6 16 16 2 38 2 1 2 370 8 1 2 3
٠.١	юрО	
ri.	or.	######################################
APPEARANCE	Color.	CNEWNHART HARMANN KNEWNNN KNEWNN KNEWNN KNEWNN KNEWNN KNEWNN KNEWNN KNEWN KNEWNN KNEWN
	Sedı- ment.	Little Cons'd Much Cons'd Cons'd Much Cons'd
	Turb'y.	+ * * * + * * + * * * * * + + + + + *
SOF	1900 Exami- nation.	Jan. 6 - 13a. 6
DATEOF	1900 Collec- tion.	に二寸形等。こことの工學程45下型は本元別程を記憶を記憶を記して記せることをは可可能を
No.	Inlros	6652 Jan. 6652 Jan. 6654 6756 6757 6757 6757 7721 7721 7721 7721 7721 7724 7724 7724 7724 7725 7727 7

TUBBIDITY-* Decided. § Very Decided. † Distinct. † Very Slight. Slight. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Colon on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish. R., Red. Rh., Reddish. W., White.

TABLE 55.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T.J. Burrill, University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, PEKIN.

No. of Bac.	per Cubic Centi- meter.	200.000 1.320.000 1.
Presence or Abs. of Colf.		0.+x.++
Temperature of Air, Cent.		-577-47-0 -644-8821-8824-88888888 -577-47-0 -644-8821-8824-88888888
ture r, C.	Tempera 91ste	<u> </u>
.7:	dgiəll ərs7/	ဖ
TRO- GEN AS	Nitrates.	88 88 88 88 88 88 88 88 88 88 88 88 88
NITRO	Nitrites.	<u> </u>
IC 3N.	Sus.	10
ORGANIC NITROGEN	Dis- solved.	25.50.00
ON	Total.	8.57.78.88.44.59.88.88.88.88.88.88.88.89.89.89.89.89.89
00	np.pud	25
SEN AS	Albuminoid Pistorial Divide	88.88.88.88.88.88.88.88.88.88.88.88.88.
NITROGEN	I'10'T	4.5.5.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.
	Free Am-	85 52988 444 88 4 5 5 4 5 5 5 5 5 5 5 5 5 5 5
N ED.	By Suspd	$\frac{\alpha \omega \omega - \alpha \alpha \omega \omega \omega + \omega + \omega + \omega}{\omega + \alpha - \omega + \omega} = \frac{\omega \omega \omega \omega \omega - \omega}{\omega + \alpha - \omega} = \frac{\omega \omega \omega \omega + \omega}{\omega + \omega} = \frac{\omega \omega \omega \omega + \omega}{\omega + \omega} = \omega \omega$
OXYGEN	By Dis- solved.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
CoS	Total.	$corr 30 cor 54 cor 37 corr 57 \mathsf{co$
.9n	Chlori	##83/-#85.000.000.000.000.000.000.000.000.000.0
Color	on Igni- tion,	は我は我のは我就我就就就就就是我就就就就就
	i p,pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ATIO)		38888888888888888888888888888888888888
APOR	no l'10T no l'10T h'vlos	
RESIDUE ON EVAPORATION.	pəpuəd sng-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
DUE 0	Dis- solved,	
RESI	Total.	389 9 9 3 3 9 9 9 9 9 9 9 9 9 9 9 9 9 9
Odor.		Cass.
APPEARANCE.	Coloi.	1
		MANANANANANANANANANANANANANANANANANANAN
	Sedi- ment.	Littile Cons'd Much Cons'd Littile VYMuch Cons'd Much Cons'd Much Cons'd Much Cons'd Cons'd Littile Li
	Turb'y.	+****
OF	1900 Exami- nation.	Jam. 6 1. 12 1. 12 1. 13 Nar. 8 Nar. 15 April 5 April 5 1. 18 Nay 4 1. 18 1. 18 June 7 1. 19 Ang. 9 Sept. 6 Sept. 6 Coc. 4
DATEOF		######################################
	1900 Coffee- tion.	6663 Jan. 6663 Jan. 6663 Jan. 6663 Jan. 66848 Feb. 6848 Feb. 66948 Fe
No.	Serial	668 688 688 688 688 688 688 688 688 688

Color on Ignition-G. Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Tunbibiry-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 56.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

₩
Z
3
. "
-
A
paties
-
2
国
-
α
32
10
6.
=
3-2
-
\vdash
20
pane.
田
H
4
_
54
0
0
E
RC
-
0
12

Paye 69 Appendix No. Appendix			
	No. of Isac.	per ('nbie ('enti- meter.	88 (100) 2 (100) 2 (100) 3 (100)
			++++ 0. ++00.0. 0. ++++++ 00.0.+:+0.0.+0.++0.
Collect Ramp Color Colo			- : ing cg : cg
180 180			
180 180	RO- EN AS	Nitrates.	0.00.00
	FIN	Zitrites.	
PATE OF 1500	0 ×		<u>8. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.</u>
PATE OF 1500	GANI		8
Horizon Hori	QN FIN	Total.	\mathbf{z}
Horizon Hori		p.pud esus-	1 - 6
	▼ ;		-
	TROGI	L'10T	8447444744444498888445888887198988884488 8454486486488488844888888448888844888884488888448888
	Z	minom.	_
Secondary Seco	Z.O.	Matter.	$\underline{\omega} \circ \underline{\omega} \circ \underline{\omega} + \underline{\omega} + \underline{\omega} \circ \underline{\omega} \cdot \underline{\omega} \circ \underline{\omega} + \underline{\omega} + \underline{\omega} - \underline{\omega} - \underline{\omega} - \underline{\omega} \circ $
Secondary Seco	SUM	solved.	
Color 1900	Cos		$\frac{\infty - \omega \omega \omega \overline{\omega} \omega$
Section 1900	'ou	Срјоци	2
PATE OF 1900	Color	on Igni- tion.	我我我我我我我我我我你你我我就我就就我我我我我就会你你你就你我我我
DATE OF ISOO ISOO		p.pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
DATE OF 1900	AT10.	= 1100	(0) ((0) (0) (0) (0) (0) (0) (0) (0) (
DATE OF 1900	APOR.	2	(S) (A) (A) (C) (A) (C) (C) (C) (C) (C) (C) (C) (C) (C) (C
DATE OF 1900	N Ev.	Depued:	8. 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
DATE OF 1900	re o	'pəatos	0 + 0 x 7 + + 0 x 0 + 0 x + x 0 0 = +0 0 + 0 0 + 0 x =
DATE OF 1900	ESID		$\begin{array}{cccccccccccccccccccccccccccccccccccc$
DATE OF 1900	H H	Total.	
DATE OF 1800 APPEARANCE 1900 1800		Tob()	
DATE OF 1800 APPEARANCE 1900 1800		or.	######################################
DATE OF 1900	ANCE	5	
DATE OF 1900	PEAF	ent.	de d
DATE OF 1900	N.		
			おやしゃでおいって思答しることとしては当立ることがある不思答してご答。
	EOF	E.Na nati	National May Natio
	DATI	λ(t) Hec- on.	
हिन्द्र हुन्द्र हुन्द्र हुन् हुन् हुन् हुन् हुन् हुन् हुन् हुन्	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
	'uN	Serial	इस्टर्स है इस है

Truminity *Decided. § Very Decided. + Distinct. ‡ Very Slight. | Slight. | Slight. | Color on Inversion—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownlsh Gray. Bh., Brownish. R., Reddish. W., White.

TABLE 57.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SOURCE OF WATER-SANGAMON RIVER, CHANDLERVILLE.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

		1101
ER,		III
PALM		JO
124	LL,	ersity.
=	REE	ver
THUE		Cniv
KTI		
7		
rt o		
\$C10		

No. of Bac. Per Cubic Centi- meter.	11.1.1.00 12.1.1.1.00 13.1.0
Of Air, Cent. Presence or Abs. of Coli.	
Temperature of Water, ('. 'Temperature 'Temperature	0.1.4.10.0.9.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.
Xitrites, ed and selection of the select	1 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -
bended.	10 10 10 10 10 10 10 10
Organial Nitrogram Polati Dis- Boylog -sil	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
nd Am.	000 000 000 000 000 000 000 000 000 00
AMMONIA. Albuminoid Tot'l	80 1128 80
Free Am- monia.	
ONY SEE ON TOTAL ON T	######################################
Chlorine.	$\begin{array}{c} \Delta \omega \nabla \nabla$
Color Color Lion.	$\frac{1-1}{4}\frac{1-0}{6}\frac$
Loss on 1g. 1 Sus. Sus. solv'd sus.	0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 + 0 +
Sus- pended.	1
Toral. Bassach Toral. Bassach Toral Control	324 + 4 334 8 327 + 6 282 8 337 + 6 282 8 338
Odor.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
ANCE.	N NN NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN
APPEARANCE Sedi- Col	Little Cons'd Little Cons'd Much Cons'd Much Cons'd Nuch Cons'd C
S B C J. I. I. J.	
DATE OF 1900 1900 Collec- Examition.	38 Scpt. 1 3 Scp
Serial No.	66612 Jan. 6662 6754 6883 Feb. 6894 6897 Mar. 7033 7048 71184 71184 7124 April 7248 7348 7348 7348 7348 7348 7348 7348 7350 7350 7360

Turbibry.—* Decided. \$ Very Decided. + Distinct. + Very Slight. | Slight. | Slight. | Color on Ish., Muddy. VM., Very Muddy. T., Turbid. C., Clondy. Color on Ish., Brownish Cray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bb., Brownish. R., Reddish. W., White.

TABLE 58.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER. BEARDSTOWN.

ER,		Illinois.
EN		of
7.	LL,	ity.
-	RRI	ers.
ARTHUR	T. J. Bu	Univ
101		
Keport		

No. of	per cubic Centi- meter.	21.500 23.5000 23.5
	Presence of Abs, of	
	Temperat of Air, U	4 + - q r - r - r - q r - r - q r - r - q r - r -
enre	Partequests of Marei	
	Height	
NITRO-	Nitrites.	全級第787228882222222828282823222222222222222
	pepuad-	g8628844554456848857888575588854858888888888
ORGÁNIC	Dis- solved.	8 2 1 2 1 1 1 2 2 2 2 1 2 2 2 2 2 2 2 2
ON	Total.	[8882288888888888888888888888888888888
N.S.	p.pud -sng	58855568888888885558555555555555555555
ITROGEN A	Por l'10T sid	<u> </u>
NITROGEN	E I'10T	本意義
	Free Am-	28 218 2 21 2 2 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3
OXYGEN CONSUMED.	solved. By Suspd	$\frac{\omega_1\omega_2\omega_1}{\omega_1} = \frac{\omega_2\omega_1\omega_2\omega_1\omega_1+\omega_2\omega_1\omega_1}{\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_2\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_2\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_2\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_2}{\omega_2\omega_1} + \frac{\omega_1\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_1} + \frac{\omega_1\omega_1}{\omega_2\omega_1} + \frac{\omega_1\omega_1}{\omega_1} + \omega$
Oxy	Total.	$\frac{-\infty \overline{\omega} + \overline{\omega} + \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} + \overline{\omega} \overline{\omega} - \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega} \overline{\omega}$
16, ,91	Chlorin	88588 : 500 = c + 10 + 4 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
Color	on Igni- tion.	我我我我你我我我我我我我我我我我我就就就就就就我我我我我我我我我说你你就我我我
	= p,pud = sug	$\frac{\alpha+5\pi\alpha+2}{4\pi}\frac{1}{8}$
RATIO	Loss on 1g	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Evare		84 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
RESIDUE ON EVAPORATION.	Sus-	
ESIDU	DIS- solved.	<u> </u>
"	Total.	
1	TODO	
CE.	Color.	ns'd N. F.09 ns
APPEARANCE		ANNAMANAMANAMANAMANAMANAMANAMANAMANAMAN
\APP.	Sedi- ment.	100 MM 100 MM M
	Furb'y.	にいる。
DATE OF	1900 Exami- nation.	Na N
DAT	1900 Collec- tion.	
No.	Serial	6827 Jan 6830 6837 6837 6837 6837 7785 7785 7785 7785 7785 7785 7785 7785 7787 778

TUBBIDITY * Decided. \$Very Decided. + Distinct. ‡ Very Slight. † Slight. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Colon on Isrition. G., Gray. B., Brown. DB., Dark Brown. LIS., Light Brown. RB., Reddish Brown. BC., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Anthur W. Palmer.
T. J. Burrell,
University of Illinois.

Sot

百
H
Ξ
20
AMPSVI
N
ΚA
~
VER
5
RIV
23
SIONI
Z
E
Ξ
T
24
ATER
L
1
OF
0
URCE
In
~

		. 00 20 5100100000000000000000000000000000000
No. of Bac.	per Cubic Centi- meter.	13.200 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 13.500 14.500 14.500 15.500 16.500 17.500
	Presenc	+ + + + + + + + + + + + + + + + + + +
ture Jent.	Tempers, (4 2 2 4 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
ature D., Te	Tempers	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	ilgiolf otali	$\pi \pi $
TRO- GEN AS	Nitrates.	88. 28.88.88.88.88.88.88.88.88.88.88.88.88.8
NITRO- GEN	Nitrites.	86.99.99.99.99.99.99.99.99.99.99.99.99.99
IC EN.	-snS	28 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
ORGANIC	Dis- solved.	66.6 6.6 6.6 6.6 6.6 6.6 6.6 6.6
ON	Total.	8.8. 1.13. 1
	i p,pud	851.15.1.15.8.8.8.8.8.8.8.8.8.8.8.8.8.8.
EN AS	Albuminoid Dis- Div'd Div'd	200
NITROGEN	L'10T	24
Z	Free Am- monia.	186
ZD.	By Suspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN	By Dis- solved.	ထားလူလွှင့် မေးမြောင်းတွေတွေလည်း မေးမြေးမှု မြောင်းများများ ကြောက်လွှာလွှာလည်း မေးမြောင်းများ မေးမြောင်းများ မေးမြောင်းများ မေးများ မေးများမှာ မေးမြ
S	Total.	$\alpha + \varpi \alpha \sin \alpha + \pi + \alpha + \alpha$
.əu	Chlori	######################################
Color	on Igni- tion.	表现基础的最高数据设置的数据设置设置的
	i p,pud	60000000000000000000000000000000000000
ATIO	no sid b'vlos	######################################
APOF	Tot'l	$\frac{45878819819824824828883}{85567788318748388283327378}$
RESIDUE ON EVAPORATION.	bended.	10. 10. 10. 10. 10. 10. 10. 10. 10. 10.
DUE (Dis- solved,	25.5
RESD	Total.	ನ್ಯ-ಎಯ್ಯಯಯಲ್ಲಿಲ್ಲ್ ಚಿನ್ಯಾಯನ್ನು ನಿ. ಚಿನ್ನಾ ಈಚ ಲೀನ್ನಾಯ ನಿನ್ನಾಗಿಕ್ಕೆ ಮ
		288.888 14.48
.1	topO	
JE.	Color.	8000000000000000000000000000000000000
RANG		REAR HARINGE REARINGER HERE
APPEARANCE	Sedi- ment.	Little Cons'd Wuch Cons'd Cons'd Wuch W.SFht Little Little Little Little Little Cons'd Wuch W.SFht Little W.SFht
	Turb'y.	$4 \% \% - \infty 76 \% \circ 5 \% \% \circ 5 \% \circ 5$
OF	1900 Exami- nation.	Jan. 1 Aug. 1 Au
DATEOF	e's	######################################
	1900 Collection.	Jan. Mar. Apr. Junk Junk Sept.
No.	Serial	6669 6774 6774 6774 6774 6774 6774 6774

LUBBIDITY—* Decided. * Very Decided. + Distinct. ‡ Very Slight. | Slight. | Colon On Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White.

TABLE 60.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arrhur W. Palmer, T. J. Burrill, University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON. (Regulars and Extras.)

No. of Bac. per Cubic Centi-meter.	25, 280 25, 200 25, 20
Presence or Allo, 100, 2011.	1 1944 14
Temperature of Air, Cent.	
Temperature	
To IdgioII TotaW	STANGE OF STANGE
Solvitics.	
N. Sitritiz	######################################
Pended.	\$\frac{2}{8}\frac{2}\frac{2}{8}\f
Ondanic National Ondanic Ondanic Silved.	
Total.	8.66.5.2.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8
Sus- Sus- Sus- Sus-	
NITROGEN AS AMMONIA. Mibuminoid Am. Programmer and programmer and	क्ष्म ते हैं के क्षेत्र के
ANN ANN I'10T	1
Ртее Азп- топіа.	455 × 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
By Suspd	$\frac{\omega}{m-1} = \frac{-\omega \omega +\omega \omega\omega \omega\omega \omega}{m-1} = \frac{\omega}{m} = \frac{\omega}$
Total, Construction of the Prince of the Pri	$\alpha F - \alpha \infty \alpha \alpha \alpha \alpha \alpha F - F - F - F - F - F - G - G - \alpha \alpha F - \alpha - G + G - \alpha G - \alpha G - G - G - G - G - G - G -$
Total.	TP-F-877-878-878-878-988888888888888888888
Срјогјис	8 4 5 5 6 9 6 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
Color on Igni- tion.	ක්රස්ත්ත්ර්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්ස්
id p.pad	$-\frac{46}{3} + \frac{1}{3} + $
on le sici	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RESIDUE ON EVAPORATION. Tolvi Loss on Evaporation. Tolvi Loss on Evaporation. Tolvi Loss on Evaporation.	表現在台灣路路路路台灣路路表表記單面隔离路場以路路上看在80回路高度在35点 五年6
bended Sus-	25 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
a portos	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
Dis.	12) ==0,00,00,00 = 0,00,00 = 0,00,00
Total.	88 88 88 88 8 8 8 8 8 8 8 8 8 8 8 8 8
Odor.	
or.	
APPEABANCE. Sedi-	
APPEAN Sedi- ment.	
C is is	
DATE OF 1900 1900 Collec- Examition.	H
DATE OF 1900 19 20Hec. Exation. nat	+5444488-07-00-00-00-00-00-00-00-00-00-00-00-00-
Serial No.	\$\$ 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6

TURRIDITY *Decided. § Very Decided. + Distinct. † Very Sight. Slight. Color Nunddy. VM., Very Muddy. T., Turbld C., Cloudy. Color on Sention G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Bi. W., White.

TABLE 60-Continued.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burnell, University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON. (Regulars and Extras.)

160,000 173,00	23.500 13.000 23.500
++++++++++++++++++++++++++++++++++++++	-1+:
Temperature of Air, Cent.	
Temperature of coccepce of the monococcus of Maler, C.	
$\text{Neight of the problem of the p$	± 55.55 ±
4 - 33 MILL	≈8 2
No. Assistant	013
34 5 6 2 1 5 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	
O 2 No 10	000
SE -2(d	
[] [[ftoT] ===================================	
## D'bud 85488567886525554 845784 8588 8888 8888 8888 8888 888	₹±±8
AMNONIA AND OLIVE AND ALL THE CASE AND A	571 160 160 160 160 160 160 160 160 160 16
A N N O O O O O O O O O O O O O O O O O	8 8 8 8 8 8
is in one in months in the first of the contract of the contra	36.2.2.2
1 1	0.00
2 - bavios 4-20-40-40-40-40-40-40-40-40-40-40-40-40-40	0000
$\begin{bmatrix} \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \bigcirc \\ \bigcirc \\ \bigcirc \bigcirc$	
Chlorine.	
20 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	i zi zi zi
	8.8 12.4 6.0 10.4 7.6 24.8
ON EVAPO Super- Supe	031.2 036.4 242.4
ON EVAL. 1.12 2.12	228.0 216.0 213.3
- 101 - 102 - 103	900
RESIDE TOTAL	
Ref. 1 1975 1 19	375.6 369.2 371.2
TobO.	
8 0 EREFERERERERERERERERERERERERERERERERERE	F.04 F.03
ANCE. COO.	NAN.
Sedi- Cons'd M. F Cons'd M. F W.W. V. W. F W. W. V. W. F W. W. F W. W. F W. W. F W.	: : :
	20 30 40
1 wedanth	-10001-
DATE OF 1900 100. 100. 100. 100. 100. 100. 100.	:::
	:::

TUBBIDITY—* Decided. § Very Decided. + Distinct. + Very Slight. | Slight. | Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish, Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy.

TABLE 60—Continued.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON. (Regulars and Entras.)

No. of Bac.	per Cubic Centi- meter.	11.000	9,000	2.000 2.000 2.000 2.000	16,000	16,500	7.500	3,900	9,750	8.300	33.300	6.700	99.400	(S).7	4.100	19,550	905.4	0001	0.000			560	
	Presenc	+-1	1	+1		1 ,	1	1	+		:	+		5.	+	-+-	+	1	+	:			1 . ;
	Tempera of Air, C												:				:			:			
	Tempers	∞ σ:		di di d	si di	ص ص ص	2:	==		: 22		2:	် ကို ရ		8 K		- 100 to		15	:		6: 6:	8 <u>8</u> 5
10.1	Heigh Dig/A		でででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででででで			0 11.9	الحط	6.0 10.0	::	12.4				3 oc	00 to	6.	01.0	?? ?:	10	:	: .	411.6	c
TRO- GEN AS	Zitrates.	9.1	라 :	\$ 16 3 	ਤ ਹੈ। ਤ ਹੈ।	3.2	1.7	ž (-	_	ંદા	\$\$ -	-	 8.8			-	1.3	- 3	- X	ズ さ -	1.1	ন্ <u>ন্</u>	- m m
NITRO- GEN	Nitrites.		50 S				-	9.8 9.8	-				•	• • •	038			200. 5		. OII			
IC EN.	Sus- pended.		196							(3.27. 	200 F		£ 5		Z 2			_		•	- 11	<u></u>	5 7
ORGANIC NITROGEN	Dis.	3.4.	568	100°	. 556 . 536	.504	0440	.536		504	.600	.50	287	15.5		.468	00 i	580	916	016.		375	18.8
0.7	Total.	76	38.5	28.00	8.80		1.016	. X. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.	000		786. 186.	.952	21 g	3.	- oc	Œ.	8:3	5966.	X:9:	192	.51¢.	619	0 min
vo.	p,pud -sus	44.0	986	287	182	9.9	.128	3.5	821.	× × × × × × × × × × × × × × × × × × ×	38	.08	176	11:0	8 = =	82.	960	5 8	=======================================	253	.000	500.	1 = 2
EN AS	Albuminoid Pis- Divios	176	208	803. 803. 803. 803. 803.	- T. T.	224	176	18	176	38	56.	303	4.6	208	38.6	.166	.176	2 7	99	156	===	176	32.3
NITROGEN AMMONIA	I'10T	.320	86.85 10.85	25.55 25.55	3000	61.88 61.88 61.88	304	x xx x xx x xx	304	200	355	27.5	284		8; 8; 0	3000	272	00101	27.2	9.10	012	80%	380
Z	Free Am- monia.	272 256	4000	208	121	.207 176	144	92.5	192	.176	.176	871.	900	07.2	86	3330	×10.	200	100	× 0	020	070.	8 8 8
N ED.	Matter:	10 m	٠٠٠٠ ن ت ن	0 7 - 0 6 1	ন কা । কা	ر د د د د د د د د د د د د د د د د د د د	2) C	0 10 20 0	9.4.6	} {~	ात - को द	∞ • • • • • • • • • • • • • • • • • • •		. 77	0 81 0 81	6.5	∞ 1	î -:	5.1	- x	9.	∞. c	==
OXYGEN	By Dis- solved.	6.4	4.00	0 .00 .00 .00 .00 .00 .00 .00 .00 .00 .		6.0 7.0	6.7	6.6	9.6	- 1-	æ ∞ ∞ −	6.2	0 %	10.	رن م م	9.9	10 = 00 11	0 0	91	9 5	5.6	6.9	. e. e.
0.00	Total.	3 12.0	12.0			20.2	4 10.9	8 II 9 0 II 5	20 10 10 10 10 10 10 10 10 10 10 10 10 10	· ∞ • ∞	5.1.5	2 10.5	1.0		7.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00	9.5	\$- 0 ₹- 0	c	= 1		101	1-1-	6 - 1 - 0
ne.	Chlori	4.0	ना चा।								10 4					≈	7 :	<u>: ::</u>	223	2 =	- 10	10 E	<u> </u>
Color	on Igni- tion.	≃ ≃	~~	~ ~ ~ ~	i zi :	~ ~	≃:	<u>~</u> ~	<u>=</u> =		~ ~	===	~~	: <u>~</u> :	≃ ≃	~	≃ <u>`</u> =	: ::::	≃:	ja	<u>: ::</u>	≃ ≃	<u>:~</u>
×	r p,pud	-	es es e	× π -	, ,	0.9 1.0 1.0	T. 0	20 20 20 20	47	: =	9 ?		0.5		0 0 0	7.	7-	- 3¢	∞ :	x w o e:		4.4	
SATIO	Loss on last of last o	36		n,∞, c	200	20 20 20 20	30.	53 63 53 63	6.61		17.6		23.00 20.00 20.00		37.32 26.32		37.2			0.00 0.00 0.00 0.00		31.2	18.88 0 × 0
VAPOI	1'10'I'	86.88 86.88	9.15 13.0	19.2 19.6	9,000	8. 6. 2. 6. 3. 6. 5. 6.	F: H: 9	2 7. 20 00 20 7. 20 00	0.45	5	\$ 18.5 \$ 00 \$ 00	8.36.0	38 4 91 8	75.636.0	∞ ∞ © ©	36.	0.27.6	2 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	380.23	x x = 2	33 +	0.35.6	28.0 26.4
RESIDUE ON EVAPORATION.	bended.	202.0 32.8 204.0 26.8	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	180.4		25 72 26 72 26 23		162.8	130.4	100.00	136.4	100	138.8	15.6	98.6	108.1	0.201	93.5		2 2 2		0.08	.0.03 € £ £6
DUE	Dis- solved.	159.6	0010	S. 51.5	181.0	200.0		0.038.0			4 186.0	16.0	56.4	53.6	S S		∞ o	8 267.6	6 2:36.4	7 19		253.2	2 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1
RESI	Total.	900	<u>∞</u> c ·	19.5 20.5 20.5 20.5 20.5 20.5 20.5 20.5 20	0 00	200.00 300.00 300.00	ic i	250.00 25	50 2	311.42	200 H	316.8 216.0	313.2 156.4	329.2 253.	371.× 265.	352 0 213.	360.8 258.		371.62	341.637.3	364.1.3	343.2.253	
			en en e	** *** **		-	_	70 00		-					- : - :								
	lobO	31.75				:::	•			•			-	-	G 15		-				7.	-	
CE.	Color.	F.07	F. F. 5	353 1113			breed to		M. F.03	-	F.06		E. E	-	66		E.0.3			5.5 5.5 5.5			
ARAN		N.N.	77.	7. 7. 7	7. 7. S	スズ	7.7	22	7.7	R. 72.	z, z	. Z	7.7		d V.N.	-		iz.		٦.		÷÷	-
APPEARANCE	Sedi- ment	p.suo.)	: :	: : :	:	: :	: :	: :	: :	:	: :	:	: :	:	Cons.d	:	:: 1	Much	VLit'le	Consid		11 11 1	;
	्राव्याम्,	* *	* * *	# # 4 31.00 =	+ + + + + + + + + + + + + + + + + + +	* * 0c =1	* :	* *	* *	*	* *	*	* *	*	10 -	*	* *	+ +	- 1	* *	* *	* *	* *
OF	1900 Exami- nation.	Apr.	: :	: : :	:	: :	: :	តិ ត : :	÷ : :	:	: :		May	:	: : : : : : :	une	:::	: 26	& :	: 1		::	<u>221</u> ±=
DATEOF	o ec. E	1-16	0.0	=212	<u> </u>	ए क	1-	<u>x 5:</u>	÷ ÷	3 23	53 50 0 50 0 50 0 50		7. 21 C	91	88	-	<u> </u>	S 151	52	-	-	೮೦	n = ::
j	1900 Collec- tion.	Apr.	: :	: : :	:	: :	:	. :	: :	:	: :	:	May.	:	: :	June	• :		7 .	: 1	£ 111 (*	4 :	: :
No.	Istros	1361	222	131	1316		7334	200	一次に	1320	1.385.1	133	1500	1561	1655	7674	371	7769	5355	をえい	7835	X7X-	1200 1200 1700 1700 1700

Color on Ignition -G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon M., Muddy, V.M., Very Muddy, T., Turbid, C., Cloudy, Trubunty -* Decided. § Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 60—Concluded.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON. (Regulars and Extras.)

MER,		Illinois
PAL.	RRILL,	versity of
ARTHUR W.	T. J. Burr	Univ
Report of		

o.	ir oic ti- er,	114 8 8 8 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
No. Isa	per Cubic Centi- meter,	
	Presence to the Abs. of the Abs.	
ture tant.	Тетрета от Аіт, С	
erure	Tempera of Wate	8. 46 50 50 50 50 50 50 50 50 50 50 50 50 50
10 1	Heigh	ware a a f f f a a rerea a a f f f a a a a a a a a rere re a a i i i
NITRO- GEN AS	Nitrates.	24
N. I.	Nitrites.	8 4 4 5 5 5 5 6 5 6 5 6 5 6 5 6 5 6 5 6 5
IIC EN.	-suS	84844455555555555555555555555555555555
ORGANIC NITROGEN	Dis- solved.	### ### ### ### ### ### ### ### ### ##
OZ	Total.	24
œ	Pud'd Sus-	98.00.00.00.00.00.00.00.00.00.00.00.00.00
TEN AS	T'ot'l Dis- pio- silo di	######################################
NITROGEN AMMONIA	Too'T	24
Z	Free Am- monia.	0.0988888888888888888888888888888888888
ED.	By Suspd	ususung innennasahanasapangang in simansah ing inakanasapangan ing inasipana
OXYGEN CONSUMED.	By Dis-	ου ο 4/-/-/- νινο 4 4 4 4 α 4 4 4 4 4 4 4 4 4 4 4 1 1 1 1
000	Total.	0001-01-000000001-00000000000000000000
	Chlori	848411184718888888888888888888888888888
Color	on Igni- tion.	ಪ್ರವಸ್ಥನವನ್ನು ಪ್ರವಸ್ಥನ ಪ್ರವಸ್ಥ ಪ್ರವ ಪ್ರವಸ್ಥ ಪ್ರವಸ್ಥ ಪ್ರವಸ್ಥ ಪ್
X.	Sus- brand	
RATIO	Toss on Tori	000001478-50888888888888 000000000000000000000000000000000000
VAPO	T.10.T	86
RESIDUE ON EVAPORATION	-sng -sng	
DUE	Dis- solved.	2. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
RESI	Total.	289.6 2233 2 24.4 218.0 2 289.6 228.6 2 289.6 218.6 2 289.6 2 28
.robO		
)r.	50.000.000.000.000.000.000.000.000.000.
APPEARANCE	Color	ROBERT REFERENCE SERVICE STREET RESERVICES
PEAR	Sedi- ment.	Little Little Little Little Nuch Little Cons'd Little Little Little Little Lons'd
		E GE G NGN EN EN
	S B C	F-0-1-000 19-1-0-1-00-1000 0-1000 -10000 000 100 10-00-10000 1000
EOF	1900 Exami- nation.	Sept.
DATEOF	1900 Collec- tion.	52888888-2000000000000000000000000000000
		7956 July 77656 July 7
.0%	Serial	200

TURBIDITY—* Decided. § Very Decided. + Distinct. ‡ Very Slight. Slight. Color Oloron—M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ignition—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Reddish. W., White.

TABLE 61.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Anthur W. Palmer. T. J. Burrill., University of Illinois.

SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON. (Regulars and Extras.)

No. of Bac.	per ('ubic ('enti- meter.	88.600 8.500 8
	Presenc	[+++++++++++++++++++++++++++++++++++++
	Tempera of Air, U	
oan re	Tempera	000000000000000000000000000000000000000
·L.	Helgh Helgh	$\frac{1+2n^2+4}{2}-\frac{1}{2}\frac{2n^2+1}{2}2n^2$
ITBO- GEN AS	Zitraties.	5884448441688162°5880888288282824860
NITEO- GEN	Ziteites.	S
	bended.	· · · · · · · · · · · · · · · · · · ·
ORGANIC	Dis.	2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 +
OLIN	Total.	6658°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°°
	i p,pud -sus	+ + + + + + + + + + + + + + + + + + +
EN AS NIA.	Albuminoid Pot'l Pot'los	<u> </u>
NITROGEN	Tol.	######################################
7	Free Am- monia.	852222411168
ED.	Matter.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN	By Dis-	$\frac{\pi}{\pi}\frac{\pi}\pi\frac{\pi}{\pi}$
Cos	Total.	$\frac{8767789678787878788888888787988797887}{86699999999999999999999999999999999999$
.9п	Chlori	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Jolor	on Igni- tion.	
	p.pud	$\frac{1}{10000000000000000000000000000000000$
RESIDUE ON EVAPORATION.	100 I	25898812856988888888756517615525755 084 -8 +568888 8+58808+58=+88085 085050000000000000000000000000000
APOL	Loss on Piston Dis-	<u>8988888888888888888888888888888888888</u>
N Ev	bended.	25
)1.E 0	Dis- solved,	$\frac{\infty+\infty\pm\infty\infty}{-}\frac{-\infty+\infty}{-}\frac{-\infty}{$
RESII	Total	23.2 2.5
	tobO	
	1	
CE.	Color.	######################################
A RAN		
\PPEARANCE	Sedi- ment	Cons'd Nuch V.M'ch Wuch V.M'ch Wuch V.M'ch Wuch Cons'd Cons'd Cons'd
	.v.q.m.J.	** + * * * * * * * * * * * * * * * * *
OF	1900 Exami- nation.	$\frac{\mathbb{R}}{\mathbb{R}^n} : : : \cdot : \frac{\mathbb{R}}{\mathbb{R}^n} : : : : : : : : : : : : : : : : : : :$
DATEOF	1960 Tollee-F	するにながらいのとってまるにもがおしいいいのおすると思います。
	Col.	Nar.
.0%	Intros	184 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

Color on Ignition - G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. RG., Brownish Gray. Rh., Brownish. R., Red. Rh., Reddish. W., White. Colon -M., Muddy, VM., Very Muddy, T., Turbid, C., Cloudy, Trumpury * Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER, T. J. BURRILL. University of Illinois.

SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON. (Regulars and Extras.)

No. of Isae.	per Cubic Centi- meter.	200,000 300,000 111,000 111,000 111,000 111,000 12,
e or Colf.	Presence Abs. of	+11+ ++++++ 1+1+1
ture Jent.	Tempera of Air, C	
ture r, C.	Tempera	re 4 g g g g g g g g g g g g g g g g g g
	ngioH otaW	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
TRO- GEN AS	Nitrates.	$\frac{4\cdot8\cdot2\cdot8^{\circ}\circ8\cdot2^{\circ}\circ2^{\circ}\circ5\cdot5\cdot5\cdot5^{\circ}\circ5\cdot5^{\circ}\circ5^{\circ$
NITRO- GEN	Nitrites.	99999999999999999999999999999999999999
ic SN:	-sng-	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
ORGANIC NITROGEN.	Dis- solved,	<u>ただちまる。+で、************************************</u>
N.O.	Total.	86 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
70	p,pud -sng	28. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
EN AS	Albuminoid Pis- Bis- Divios	1
NITROGEN	I'10T	### ### ### ### ### ### ### ### ### ##
Z	Free Am-	######################################
ED.	By Suspd Matter.	$\begin{array}{c} \phi \stackrel{.}{U} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \phi \stackrel{.}{U} \overline{\alpha} \overline{\alpha} \overline{\alpha} \phi \\ \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha} \overline{\alpha}$
OXYGEN	By Dis-	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
Cos	Total.	######################################
.9n	Chlori	88 9 - 63 80 80 80 80 80 80 80 80 80 80 9 4 5 4 4 70 4 4 4 4 4 8 65 65 65 65 65 65 65 65 65 65 65 65 65
Color	on Igni- tion.	热热热度玻璃度转换铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁
	i p,pud	$\frac{\omega_{\alpha}\omega_{\alpha}+\omega_{1}\omega_{\alpha}\omega_{2}\omega_{1}\omega_{1}\omega_{1}+\omega_{\alpha}\omega_{\alpha}}{\iota_{\alpha}+\omega_{1}\omega_{1}\omega_{2}\omega_{2}\omega_{2}\omega_{3}\omega_{3}\omega_{3}\omega_{3}\omega_{3}\omega_{3}\omega_{3}\omega_{3$
ATION	p,Alos	000000000000000000000000000000000000000
APOR	T'ot'l	40.48.48.49.42.88.88.88.88.48.48.48.88.88.88.88.88.88.
RESIDUE ON EVAPORATION.	pepued,	1650 1650
UE 0	solved.	198 66 8 1 8 1 8 1 8 1 8 1 8 1 8 1 8 1 8 1
ESII	-siQ	306.8 5024. 116.8 5024. 116.8 5024. 116.8 5024. 116.8 5024. 116.8 5024. 116.8 5024. 117.6 5024. 117. 117. 117. 117. 117. 117. 117. 11
	Total.	
Odor.		AO Z
	or.	######################################
APPEARANCE	Color	MANANAMANANAMANANAMANANAMANANAMANANAMANANAMANANAMANANAMANA
PEAR	nt.	Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Much Little Much Cons'd Cons'd Cons'd Cons'd Cons'd
AP.	Sedi- ment	Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Much Little Much Little Much Little Much Little Much Little Much Little Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Cons'd Cons'd
	Turb'y.	c - 5312f 248 c 5f 22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
OF	1900 Exami- nation.	Apr.
DATEOF		400==50x2440052500244201-00==50x3222
	1900 Collection.	Tata a second a secon
No.	Serial	2000

COLOR ON IGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BC., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. V.M. Very Muddy. T., Turbid. C., Cloudy. Colon-M., Muddy. Turbidity-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 61—Concluded.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICE.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON. (Regulars and Extras.)

Report of Authur W. Palmer, T. J. Burrill, University of Illinois.

4		
No. o Isac.	tubic Centi- meter.	ਲਾਗ ਜੋ ਨੀ ਜਾਂ ਵਲੀ ਜਾਂ ਜੋ ਲੋ ਲੇ ਜੇ ਲਾਜ਼ੇ ਗੇ ਹੈ ਜੋ ਗੇ ਗੇ ਗੇ ਜੋ ਹੋ ਗੇ ਗੇ ਗੇ ਜੋ <u>ਹੈ</u> ਜੋ ਗੇ ਜੋ ਹੈ ਜ
	Presence to selvence to selven	
	Tempera of Air, to	
arna oana	Temperation To	ក្រុងស្ត្រីក្នុងស្ត្រីស្ត្
10 11	11e1gh 1840	
NITRO- GENAS	Nitrates	8888 64 8888 98 4 × 8 2 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
N	Nitrites.	88
NIC GEN.	-sng	88.86 4.46 4.46 4.46 4.46 4.46 4.46 4.46
ORGANIC	Dis-	
- 7.	Total.	28. 29. 29. 29. 29. 29. 29. 29. 29. 29. 29
AS.	m P.pud	**************************************
	T'10T Piston Manipoid Piston P	892688888888888888888888888888888888888
NITROGEN	I'10T	6.74.23.48.88.48.48.48.48.48.48.48.48.48.48.48.
	Pree Am- monia.	0.000 0.000
EN MED.	By Suspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN	By Dis-	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
	Total.	$a \circ a \circ a \circ r \circ a \circ a \circ a \circ a \circ a \circ a \circ $
		81 91 91 91 10 mm m — m + m m m m m m 11 21 61 m 91 11 11 10 10 10 10 10 10 10 + — —
	on Igni- tion.	
10N.	= p,pud -sus p,ajos	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ORAT	o sid	11 11 1 + + + + 0 11 + x 2 0 0 x + 2 2 0 11 0 11 1
RESIDUE ON EVAPORATION	Pended.	25
E ON	-sus	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
SIDE	Dis- solved,	25
<u> </u>	Total.	2
Cdor		
	or.	######################################
SANCE	Color.	ZAZZEZZEZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ
APPEARANCE.	Sedi- ment.	Much ('ons'd Much ('ons'd Much ('ons'd Much ('ons'd Much Much ('ons'd
· v.gan.l,		
£4.	1900 1900 Tollee-Enami- tion, nation.	28 - 22 x 5 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
DATE OF	1900 Tollve- I	
	100	2 x 2 x x x x x x x x x x x x x x x x x
35%	Lifter	흊춪퐟칮쳁찞찞귳춪뜢뜢뜢뜢귳쯗퍞퍞퍞퍞똣똣똣똣똣

Color on lention-C. Gray. B. Brown. DB. Dark Brown. LB. Light Brown, RB. Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White, Colon-M., Muddy, V.M., Very Muddy, T., Turbid. C., Cloudy, Transpiry *Decided, & Very Decided, + DistInct, + Very Slight, i Slight.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON. (Regulars.)

No. of Isac.	per Cubic Centi- meter.	2.000 2.0000 2.0
	Presenc	+ + + + + + + + + + + +
ture ent.	Tempera of Air, C	ಹಿಎಟ್ ಆಪರ್ಕ್ ಚಿತ್ರಗಳ ಚಿತ್ರಗಳ ಪ್ರಾಪ್ತಿಸ್ತೆ ಪ್ರಾಪ್ತಿಸಿ ಪ್ರಾಪ್ತಿ ಪ್ರತ್ಯ ಪ್ರತಿ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರತ್ಯ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರತ್ಯ ಪ್ರತ್ಯ ಪ್ರತ್ಯ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರತ್ಯ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರಕ್ಷ ಪ್ರಕ್ಷ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರತ್ಯ ಪ್ರಕ್ಷ ಪ್ರಕ್ ಪ್ರಕ್ಷ ಪ್ರಕ್ತ
erare.	Tempers	
	Heigh Wate	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
A S	Nitrates.	5 34 5 5 4 8 4 4 8 8 6 8 0 1 2 8 6 9 8 4 6 8 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
NITEO.	Nitrites.	0.000000000000000000000000000000000000
	bended.	86 1 1 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
ORGANIC NITROGEN.	-sns	3)
)RGA ITRO	-sid solved.	La para para para para para para para pa
Z	Total.	86.6.5.5.8.8.5.8.5.8.5.8.8.8.8.8.8.8.8.8
ss	p.pud -sus	88515448886844 - 884476981311811888888888888888888888888888888
EN AS	Albuminoid Am. Sus. Sid. Sid. Sid. Sid. Sid.	84 98 88 86 88 86 88 88 88 88 88 88 88 88 88
NITROGEN AMMONIA	Tor.	24 8 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Z	Free Am- monia.	4.5.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.
SD.	By Suspd	$\frac{\omega}{10^{-17}} + \frac{1}{10^{-27}} + \frac{\omega}{10^{-17}} + \frac{\omega}$
OXYGEN CONSUMED.	By Dis-	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
CON	Total.	
.en	Сһ1огі	8:43024-9-8-1-8-1-8-1-8-1-8-1-8-1-8-1-8-1-8-1-8
Color	on gni- ion.	
	In nud	
10N.	Sus.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
RAT	Torilloss on 1g, n. Sulvid Sulvid	884488544888888888888888888888888888888
VAPO	1.10L	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ON E	pəpuəd sns-	83738833333338888888888888888888888888
RESIDUE ON EVAPORATION.	Dis- solved,	# (4 # 0: 6 mm m m m c: 6 6 6 c m m c: 0 6 c c m c: 0 6 c c c c c c c c c
RESI	Total.	385 8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	торО	<u> </u>
	1	
	Color.	######################################
RANG	ŭ	
APPEARANCE.	Scdi- ment.	Con'sd Much Con'sd Nuch Cons'd Nuch Con'sd
	Turb'y.	r 1 2 2 2 3 2 2 2 3 2 2 2 2 2 2 2 2 2 2 2
0.5	1900 Exami- nation.	Nan. Nan. Nan. Sept. 21. 1. 22. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
DATEOF		457222, 12-1128-15800000000000000000000000000000000000
D	1900 Collec- tion.	626 Jan. 655 .
.o.V	Serial	\$5.50

Turbidity—*Decided. §Very Decided. † Distinct. † Very Slight. | Slight. | Slight. | Color on Is intron—G., Gray. B., Brown. BB., Brown. BB., Brown. BB., Brownish. R., Reddish. W., White.

TABLE 63.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of ARTHUR W. PALMER, T. J. BURRILL. University of Illinois.

SOURCE OF WATER MISSISSIPPI RIVER, GRAFTON. (Regulars.)

No. of Bac.	per Cubic Centi- meter.	24.700 24.700 24.700 24.700 24.700 25.000 26.000 27.000
	Presence Abs. of	1++++1+++++++++++++++++++++++++++++++++
	Pempera of Air, C	호 · · · · · · · · · · · · · · · · · · ·
ture r, C,	rempera	00%+0000000+66==62%54%8% : 58%8%8%8%8%8
1.1	Неіgh Ияте	$\frac{31+4446+3440}{62646+346+346+346+346+346+346+346+346+346+$
0- IN AS	Zitrates.	$= \frac{1}{8} + $
NITRO- GEN	Nitrites.	00.00
S. C.	sus- pended.	656 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
ORGANIC NITROGEN.	solved.	\$\$\frac{1}{2}\$\fra
ON	Total.	た 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	p.pud	+ 20 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
N AS		857774488788888888888888888888888888888
NITROGEN	Tor'l	85 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Nr	monia.	85988588888888888888888888888888888888
	By Suspd Matter.	$ + \omega \circ - + \psi \circ (\psi + \varphi \circ f + \varphi \circ i) \cdot (\psi + \psi \circ \psi \psi \circ \circ i) \cdot \varphi + i \varepsilon \cdot (\psi \circ \varphi \circ \psi \circ \psi - \psi \circ \circ i) \cdot \varphi + i \varepsilon \cdot (\psi \circ \varphi \circ \psi \circ $
OXYGEN CONSUMED.	solved.	$ \cos \alpha \cos \alpha + \cos \alpha + \sin \alpha \cos \alpha + \cos \alpha \cos \alpha + \cos \alpha \cos \alpha \cos \alpha + \cos \alpha \cos \alpha$
Cons	Total.	
ne.	Съдоти	လွှဲ့နှံ့လွန်္ကလေး လလလုပ်ပေး လုပ္ပလုပ္ပလုပ္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့
Color	on Igni- tion.	虽良钱我感就就我说我我就就就就就就就就就就就就就就就就就就就
· i	i p,pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
SAT10	Loss on 18. Poly of 19. Poly o	4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
APOI	Tot.1	$\frac{1}{100} \frac{1}{100} \frac{1}$
RESIDUE ON EVAPORATION.	bended.	25
DUE (DIS.	88686888888888888888888888888888888888
RESL	Total.	200 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	robO	
jei jei	Color.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RANG	5	RANKEN PROPERTY OF THE PROPERT
. APPEARANCE.	Sedl- ment.	Much Cons'd Much Much Much Cons'd Much Much Cons'd Muc
-	Jurb'y	にこるでののでできょうと然につたがいったがとしては多いってない。 **+**********************************
OF	1900 1900 Collect Exami- tion, nation.	
DATEOF	lee-le	
	Collection.	Mar
Zo.	Serial	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$

Color on lengtion -G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Reddish. W., White. Collon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Treenerr * Decided. § Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 64.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arrhur W. Palmer, T. J. Burner. University of Illinois.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM ILLINOIS SHORE, ALTON.

No. of Bac.	per Cubic Centi- meter.	16, 400 16, 400 16, 800 16, 800 16, 800 17, 800 18,
e or Coli.	Presenc	+ + + + + + + + + +
	Tempers of Air, (4.0 r. Q.W.W. D.W. W. D.W. W.
er. ature	fielgh of a variable of the stranger of the st	: 01-00 - 01-01-01-01-01-01-01-01-01-01-01-01-01-0
A 38	Zitrates.	$\underbrace{\text{sign}_{S}\text{sign}$
NITRO-GEN	Nitrites.	86.90.90.90.90.90.90.90.90.90.90.90.90.90.
z c	·pəpuəd	2000 2000
ORGANIC NITROGEN	Dis- solved.	0.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4.4
QN IN	Total.	
	į p.pud -sns	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
EN AS	Torr	41 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
NITROGEN	I'10T	4.84
Ž	Free Am-	1.00.00.00.00.00.00.00.00.00.00.00.00.00
ED.	By Suspd	
OXYGEN	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ిప్ప	Total.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
	Chlori	$\frac{\cdot}{\exp - \exp (3)} \frac{1}{2} \frac{1}{4} \frac{1}$
Color	Jgni- tion.	
ow.	i p.pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ORATI	Discon Loss on	
RESIDUE ON EVAPORATION.		<u> </u>
ON	Sus-	### ##################################
SIDUE	Dis-	283.2 2 2 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3
RE	Total.	283. 2.261 283. 2.261 283. 2.261 283. 2.261 283. 2.261 283. 2.261 283. 2.261 284. 4.175 287. 4.113 287. 5.28 287. 5.28 287. 6.123 288. 8.213 288. 8.2
.1	орО	
	01.	######################################
RANCE	Color	ZERZEZZEZZEZEZEZEZEZEZEZEZEZEZEZEZEZEZE
APPEARANCE.	Sedi- ment.	Cons'd " Much Cons'd Nuch Con
	Turb'y.	で こので もの で ご も し こ に ご む し こ こ と い こ こ こ こ こ こ こ こ こ こ こ こ こ こ こ こ
OF	1900 Examination.	Jan. "" "" April "" June June "" Aug. Sept.
DATEOF	1900 Collection.	Jan. Mar. Mar. 1104. 110
	Serial	\$88.00 \text{Constraints}

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. I.B., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Color-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Turbidtry-* Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER - MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM ILLINOIS SHORE, ALTON.

ARTHUR W. PALMER,	T. J. BURRILL,	University of Illinols.
Jo		
teport		

o. of Bac.	per Cubic Centi- meter.	11.60 12.64 13.64 14.60 15.35 16.00 17	050 050 000
Colf.	Presence Abs. of		11
	Tempers of Air, (
arnar	Heigh Wate Temperation of Wate	0 0 1 - 20 0 10 10 10 10 10 10 10 10 10 10 10 10	ාල ක දුරු දුරු දුරු
O- IN AS	Nitrates.	£ \$ £ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$	85.55
NITRO- GEN	Nitrites.	8199999999988899998889999999	200
	Sus.	1	155
ORGANIC NITROGEN.	solved.	811-15: 85: 45: 45: 45: 45: 45: 45: 45: 45: 45: 4	· ×
OHU	Total.	<u> </u>	
	g p,pud	55 2 5 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	<u> </u>
SN AS		- 4888	169
ITROGEN	Torl Dis-	88555458854555885455988588558555	
Nr	Free Am- monia.	45.7.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	
D.	By Suspd	αρού τρορμού - ε ορμοτικού συρο Τυσον	ن -
OXYGEN CONSUMED.	By Dis- solved.		-11-
CON	Total.	$ \begin{array}{c} $	∞ ∞ Ω Ω
.əu	Chlori	$\overbrace{000} \\ 000 \\ 001 \\ 0$	- oc
Color	on Igni- tion.	据报准证证 改良我就就就就就就就就就就就就就就就就就就就就就就就就就就就就就	0.13.
ż	i p,pud	్లు ఒక్కా ఈ క్రైడాగా లోని అక్కు కార్యాల్లు అక్కా చెప్పాని ఏ — ప్ర ఈ ` బ్రీ ` ` బ్యా ఈ ఉంపు ఈ ` బ్యా ఈ ఈ బ్యా ఈ ఆ బ్యా ఈ బ్యా ఈ ఆ బ్యా ఈ ఆ బ్యా ఈ బ్యా ఈ ఆ బ్యా ఈ బ్యా ఈ ఆ బ్యా ఈ బ్యా ఈ బ్యా ఈ ఆ బ్యా ఈ బ్యా ఈ ఆ బ్యా ఈ ఆ బ్యా ఈ బ	∞
RATIC	p _i Alos		∞. ∞. ∞. ∞
7AP01	Tot'l	※8887-4817-888888223233333822-822388222228※8887-4877-828233333333※8887-4877-82823333333※8887-4872333※8887-487233※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733※8887-48733<li< td=""><td>SI</td></li<>	SI
ON EVAPORATION	bended. Sus.	888.888.888.888.888.888.888.888.888.88	197.6 5.0 8.0
	Dis- solved.	्ळळळ चक राग्यक च ककक न राग्यक च न करा चळळ च राष्ट्र	00 00 12 00 12 00
RESIDUE	Total.	1 6 4 4 00 0100 01 400 400 60 1 401 1 101 101 100 40 1 400 00 60	च्या ध्य
	10.00		320
.1	tobO		
ធំ	Color.		전 전 연 연 연
RANC	S	संस्थानस्त्रेत्रस्त्रेत्रस्त्रेत्रस्त्रेत्रस्त्रस्	7.7
APPEARANCE	Sedi- ment.	Consider Con	Much
	:X.q.m.J.	10日本の15mの15mの15mの15mの15mの15mの15mの15mの15mの15m	* *
2 O E	1900 Exami- nation.	Nar carried Nar ca	3
DATEOF	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	→ ままにおいた まどいで まどが - 50 変形 いなる 82 変 と 50 を 82 24 + 1 ∞ - x で 31	25 10
DA	0	5. 10	+
DA	1900 Collec- tion.	6621 Jan. 6632 Jan. 6632 Jan. 6633 J	8324 **

TURBIDITY -* Becided. § Very Decided. + Distinct. ‡ Very Slight. | Slight. | Color | Color | Color | William | Color | William | William

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

No. of Bac.	per Cubic Centi- meter.	8. 4. 8. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6.	9
	Presenc		
	Tempera of Air, C	4.0	
arman	Tempera	000001000040000001080440448888888888888	
10 11	dgi9H otaW	0.1-8 0.000 0.00000000000000000000000000000	
TRO- GEN AS	Nitrates.	$\frac{1}{\circ \text{tr} \text{tr} \text{sec}} = \frac{1}{4 \cdot \text{sec} \text{tr} \text$	
NITRO- GEN	Witrites.	888 888 888 888 888 888 888 888 888 88	200
IC SN.	Sus- pended.	4.55.4.4.5.5.5.4.5.5.5.5.5.5.5.5.5.5.5.	
ORGANIC NITROGEN.	Dis- solved,	865 865 865 865 865 865 865 865	100.
ON	Total.	2.5. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	
-	E papud -sns	86.50 8.4440 14.46 8.50 8.50 8.50 8.50 8.50 8.50 8.50 8.50	200
EN AS	Pionimino pionim	2	-
NITROGEN AMMONIA	I'10T	8. 4. 4. 4. 8. 8. 8. 4. 4. 4. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8.	-000.
N	Free Am-	<u> </u>	
9.6	By Suspd	$\frac{0.4-100}{100} = \frac{1.40}{100} = \frac$	7.00
YGEN	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	-
OXYGEN CONSUMED.	Total.	$\frac{15654998}{9999999999999999999999999999999999$	5
.əu	Срјогі	က်တန္-လုပ္သံုန္ ကယ္ေၿးယည္းမယ္မယ္မရာနယ္လုပ္သံုနယ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုတ္လုပ္သံုလ္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုလုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုလုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုတြက္လုပ္သံုလု	2
olor	on Igni- tion.		
	s p.pud	9-07-1-88 : 8888 6 6 6 5 6 6 6 6 6 6 6 6 6 6 6 6 6	
ATION	Loss on sid sid o sid o sid o sid	$ \begin{array}{c} 888989 - 489 & 121 - 512 - 5$	3
RESIDUE ON EVAPORATION.	Tot'l Loss		-
N EV.	.suS	0.000000000000000000000000000000000000	*
UE O	solved.		2
ESID	Dis-	ထြတ္လုံး မေတြသည္။ မေတြကို သည္ မေတြကို	-
<u> </u>	Total.	8.82 8.82 8.83	
٠,	торО		
	or.	PERFERENCE REPERENCE REPERENCE REPERENCE PERFERENCE REPERENCE REPE	ρ,
ANCE	Color.	MANANA IN TANANA MANANA	
APPEARANCE	Sedi- ment.	Cons'd Nuch Cons'd Much Whuch Whuch Cons'd Whech Whech Cons'd Whech Cons'd Whech	
	Turb'y.	* * + * * * co. · * co.co.* * * * * * * * * * * * * * * * * * *	
OF	1900 Exami- nation.	Jan. Jan. Mar. 38 Sept. 6 Sept	Oct.
DATEOF	1900 Collec-	40-44-4-1200-1284-558400088000884-18-80688 m 558	0
.oV	Serial	6622 Jan. 6652 Jan. 6652 Jan. 6652 Jan. 6653 Mar. 7002 Mar. 7005 Mar. 7173 Mar. 7173 Mar. 7173 Mar. 71745 Mar. 7175	

Turbidity—* Decided. § Very Decided. † Distinct. ‡ Very Slight. ∥ Slight. Colon.—M., Muddy. V.M., Very Muddy. T., Turbid. C., Cloudy. Colon on Ignition—G., Gray. B., Brown. DB., Bark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Reddish. W., White.

TABLE 67.

STREAMS ENAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Z
0
5
7
- 4
53
四四
0
provide particular par
.73
-
×
00
33
35
=
-1
7
0
500
I
回
0
7.
P
S.
=
-
E
~
D
,0
-
亩
1%
0
_
В,
ER,
1
1
RIVER,
1
1
1
1
1
1
1
1
-MISSISSIPPI RIV
-MISSISSIPPI RIV
ER-MISSISSIPPI RIV
-MISSISSIPPI RIV
ATER-MISSISSIPPI RIV
ER-MISSISSIPPI RIV
WATER-MISSISSIPPI RIV
ATER-MISSISSIPPI RIV
OF WATER-MISSISSIPPI RIV
OF WATER-MISSISSIPPI RIV
OF WATER-MISSISSIPPI RIV
RCE OF WATER-MISSISSIPPI RIV
F WATER-MISSISSIPPI RIV

Report of Anthur W. Palmer, T. J. Burrill, University of Illinois,

No. of Bac.	Cubic Centi- meter.	10,000	16,800	20,600 20,600	36.000	118.000	34.(80)	SE (SE)	162,500	66.99	72,500	16.000	15.000	16.08 20.08	29.500	1.000	1.000	3,500	1.750	3.600	2,300	8 8 8 8 8 8 8	6,000	(X)F-P	2.900	6,400
Presence or Abs. of Coll.		11	+	+1	1 +	++	.] -	++	-] -	+	+	1+	-	1 1	+-	+-+	-+	:	- :	٠.	S	-	-	: 1	1	٥.
Juot	Tempers of Air, C	4.0	00		9.0		0.0	တံ့ မှ				2.8	£5		55	S &	.96	£ 33	. o.	667	.96	25.00	0; 0 0; 0	300.	30.	30.
ature er, C.	remper of Wate	0.0	00		= -		0.0) -	0	ယ် ဝ	i oc	<u> </u>	30.	S. 57	19.		- 			∞ 000 000 000 000 000 000 000 000 000 0	36.	688	· ·			
	fgielf faist	61	= 0	න : න :	97.0	. vc	±;	2 <u>1</u>	=	1.5	7	2 E	10.1	± ∞.	× 1	9:5	7C 5.5	10 1 10 1	- 70 5 to	4.2	÷: نۍ	6.5	م 5.	. 6	10	9
TRO-GEN AS	Zitrates.	70	80.0	1.08	89.		5.6	ž 5	30	7.7	1.08	2 27	æ	£	÷. 9	<u>.</u> T	80	80.1	2 89	.36	.16	%.	÷ §	1.00	=	.36
NITRO- GEN	Zitrites.	.013											_													-
S.S.	Sus- pended.	.313	.418	888	4.	1.168	e	1.24x	1.336	E %	1.41	1.76	.608		1.633	.616	± ±	.336	159	513	168	98.	元 文 で	0.1	681	
ORGANIC NITROGEN.	Dis- solved.	8448	355	5.00 5.00 5.00 5.00 5.00 5.00 5.00 5.00		672	3.	515	861	9. 10.0	9.	+ - -	3	X 22 X 22 X 23 X 24 X 24 X 24 X 24 X 24 X 24 X 24 X 24	308	151	45%	##60	286	23	2000	101	5	312	53.3	× = = = = = = = = = = = = = = = = = = =
ON	Total.	.76	φ ξ	g; ∞	€ 2	Z.	00 f	50 61	03	86.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1.8 1	10:	Z 21	.038 830.	8,15	6.1	- oc	G:	35.0		133	97	- 8	5 5	9000	8	.76
	E p,pud	### ### ##############################	208	25 28 138 138	48.8	576	91.		3	.355 .355 .355	541	913	352		384	107.6	<u>%</u>	200	38		.035	33.	3, 3	300	333	200
EN AS	Albuminoid Piscolvid	.24	.16	으 은 참	.144	30.5	288	808	886	10.1	256	8 15 8 15 8 15 8 15	176	96.1	128	3 2	19.5	92.	<u> </u>	176	33	x :	X 20	26	199	301
NITROGEN AMMONIA	[1'10'F]	384	368	368	384	5 %	2448	2 22	876	576.	- oc	200 200 200 200 200 200 200 200 200 200	30.00	576	515	ζ σ. () ~ ()	2.5	000 S	301	33	27.53	523	9 3	2 00	<u>x</u>	133
Z	Free Am-	54.63			-			_	•			_	•													
	By Suspd	6.1		J			1- (20 01	723	{- {-		x -	10	- 1-	. 1	- 00	\$3	بان <u>-</u>	· · · ·	9.	9.		- 0	 	≈ ∞ ∞	5 5
OXYGEN	By Dis- solved.	6.0	. 60	23 :	2 - 0 2 - 0 2 - 0	0.00	0X 0	0 00 0 10 0 10	ee :	910	- ∞ - ∞	12.00	-	x -	-:	, ka		10.1		-:	{- i) - ·	_ x	2 :	7.1	S.
Cos	Total.	10.7	10.9	- x	ලා බ ස ග	0 00	13.5	- 15 - 15	19.5	2 7	9:	: F		5. 3. 2. 3. 3. 3. 3.	7.12	2 2 2		= 15	= =	10.7	= 3	× :	2 =	1 = -	5.3	16.5
.эп	Chlori	12.	تن. ئن	9.6	ac c.	9.4	٠: ٥ ٠: ٥	35 25 25 26) in	-	 	.01	n 00.	- i	- 0	7.	33.0	- 00	± ∞	~ :	· ·	x, ±	- 73	~	···
)r																										
010.	on Igni- rion.	≃ ≃	≃ :	± ±	~ ≃	R.B.	~		-	≃ ≃	: ≃:	<u>~</u> ~	≃:	<u>~</u> ~	<u></u>		-	<u> </u>	· ~	~	~:	<u>.</u>	· ·	· ~	~	1018.
.	= p,pud			3.6 G.	. 0	 د ن	oc /	21 C	_	x ≒	- 50	<u>ت</u> د	oc .	ж <u>Ф</u>	o	C 23	2)	+ 5	-		\$. 0	2 =	- 01	??	10.4 DB.
.	= p,pud	. î. 9		3.6 3.6	. o	1 53.6 E	oc /		_	x ≒	19.6	<u>ت</u> د	oc.	2 S S	1 30 1	C 00	15 2	÷ 0	\ \(\infty \)		5.6		15 5 15 15	- 01	3.5	2 10.4
.	i p, pud -sng p, v los	8. 24. 8 24. 8 18. 8 6. 4	31.2	6 30. H.c		30.153.6	8. 8.05 9.		15.6 22.1	x :0: x = -	6 18. 19.6	16.25 20.35 30.56	(5) (5)	25 50 25 50	8 38.4.20 1	0. 4. 0. 4. 0. 6. 0. 6.	1 27.9 15.3	- 50 - 50 - 50 - 50 - 50 - 50 - 50 - 50	(o)	∞	16.1 7.6	25.	26 113 61	18. 39.2	21 21 21 21 21	6 21.2 10.1 1
.	Torri Posson 18:0 Solv'd Sus- pended.	626.8 24.8 2. 25.2 18.8 6.4	34.2 31.2	631.6 30. H.C		30.153.6	× 51.6 20.8	6.5 10 10 10 10 10 10 10 10 10 10 10 10 10	838. 15.6 22.1	6.07.6 30.8 6.8 16.1 13 4 14	37.6 18, 19.6	16.25 20.35 30.56	(5) (5)	136. 131.2 21.8 4 21.6 9.6	858.8 38.4 20 1	2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1 27.9 15.3	.6.40.1 38. 35.1	(o)	∞	16.1 7.6	25.	26 113 61	18. 39.2	21 21 21 21 21	6 21.2 10.1 1
.	- hovios	13.626.8 24.8 2.	20. 34.2 31.2	61.2 29.6 26. 3.6.	13 42 8 20. 20. 30. 30. 50. 50. 50. 50. 50. 50. 50. 50. 50. 5	× 401.681 30.153.6 H	6 (9) 821.6 20.8	20% FO. 16.% 19.10. 1.11	150.838. 15.6 25.1	250 C.27.6 20.8 C.8	901 81 97.6 10.6	28. 4.20. x 17.2 x 6 26. 10. 16. x 20. 6	150 6'22.8 22. 8	38. 136. 13. 14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15	502.858.8 38.4 20 1	6 182 8 11.4 21.9 23.0	.8 82.8 12.4 er. 2 15 2	2000 200 300 300 400 400 400 400 400 400 400 4	130. 150. 150. 150. 150. 150. 150. 150. 15	121, 16.8 38.1 8.4	111.624. 16.1 7.6	.6 46 856. SE.	121.458. 42.115.64 128. 16.1.38. 10.11	770.857.2 18, 39.2	6 240.139.2 39. 17.3	176.431.6 21.2 10.1 1
.	Piped Sus- Prod. P	211.6 13.626.8 24.8 2.	189.2 20. 34.2 31.2	27.6 61.2 29.6 26. 3.6	13 42 8 20. 20. 30. 30. 50. 50. 50. 50. 50. 50. 50. 50. 50. 5	156 × 101.2,81 30.153.6 I	\$161.6 (8) \$21.6 \(20.\times\)	[0.1 183 6 5 6 5 6 5 6 6 6 6 6 6 6 6 6 6 6 6 6	116.1 160.838. 15.6 22.1	135.6 History 6.25.6 10.8 6.2	2116. 407 237.6 18. 19.6	6 110 4 863 9 16 1 16 8 29 6	. 4 168.8 159 622.8 22.	25 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2 116.4 502.858.8 38.4 20 1	1 1 1 6 182 8 1 4 2 1 2 23 3 3	6 180.8 82.8 12.4 27.9 15.2	6 192. 83.6 40.1 38. 2.4	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 195.6 121. 16.8 38.1 8.4	6 168. 111.624. 16.1 7.6	1 27.6 116 x 36. 32. 11.	110 K 78 16 1 36 10 1	126. 770.857.2 18. 39.2	111.6 240.139.2 22. 17.3	8 60 t 176.431.6 21.2 10.1 I
	- hovios	13.626.8 24.8 2.	189.2 20. 34.2 31.2	. 1 [64.8 157.6 31.6 20. 41.6 18.8 27.6 61.2 29.6 26. 3.6	13 42 8 20. 20. 30. 30. 50. 50. 50. 50. 50. 50. 50. 50. 50. 5	558. 156 × 401.281 30.153.6 I	252 4 161.6 (8) 8 21.6 20.8 .8	550, 113, 558 40, 16,853,81 1584, 100,1 1183 6 15 5 10, 35, 51	577.2 [116.4 460.838. 15.6 22.4	259 2 125 6 253 6 27 6 20 8 6 8	613.2 116. 497.237.6 18. 19.6	200 1 132 138 130 18 17 2 3 6 1 913 6 110 1 803 9 16 1 16 8 29 6	. 4 168.8 159 622.8 22.	38. 136. 13. 14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15	2 116.4 502.858.8 38.4 20 1	11.6 182.8 11.4 21.2 23.3	6 180.8 82.8 12.4 27.9 15.2	192. 83.640.1 38. 3.4	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	121, 16.8 38.1 8.4	6 168. 111.624. 16.1 7.6	. 127.6 446 856. 32. 34.	121.458. 42.115.64 128. 16.1.38. 10.11	896.8 126. 770.857.2 18. 39.2	380 111.6 040.139.0 30.17.0	336.8 160 4 176.431.6 21.2 10.1 1
RESIDUE ON EVAPORATION.	Piped Sus- Prod. P	211.6 13.626.8 24.8 2.	189.2 20. 34.2 31.2	27.6 61.2 29.6 26. 3.6	13 42 8 20. 20. 30. 30. 50. 50. 50. 50. 50. 50. 50. 50. 50. 5	156 × 101.2,81 30.153.6 I	252 4 161.6 (8) 8 21.6 20.8 .8	[0.1 183 6 5 6 5 6 5 6 6 6 6 6 6 6 6 6 6 6 6 6	577.2 [116.4 460.838. 15.6 22.4	259 2 125 6 253 6 27 6 20 8 6 8	2116. 407 237.6 18. 19.6	6 110 4 863 9 16 1 16 8 29 6	. 4 168.8 159 622.8 22.	25 6 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	2 116.4 502.858.8 38.4 20 1	1 1 1 6 182 8 1 4 2 1 2 23 3 3	6 180.8 82.8 12.4 27.9 15.2	6 192. 83.6 40.1 38. 2.4	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	6 195.6 121. 16.8 38.1 8.4	6 168. 111.624. 16.1 7.6	1 27.6 116 x 36. 32. 11.	507 6 140 6 141 408 15 10 60 507 6 140 8 158 15 1 35 110 1	126. 770.857.2 18. 39.2	380 111.6 040.139.0 30.17.0	336.8 160 4 176.431.6 21.2 10.1 1
RESIDUE ON EVAPORATION.	Total. Polyos-sid Sus-bended. Disconding Sus-bonded. Prof. From Sus-bonded. Sus-bonded. Sus-bonded.	211.6 13.626.8 24.8 2.	209 2 189.2 20. 34.2 34.2	F.1 (288.8.8.27.6 61.229.6 26. 3.6)	7.1 214. 201.2 42.5.20. 20.	2.4 558. 156 × 101.2,81 30.153.6 I	17.4 220 4 161.6 GD 8-21.6 -20.8	570 112 50X 10 10 10 10 10 10 10 10 10 10 10 10 10	577.2 [16.4 190.838. 15.6 22.1	259 2 125.6 233 5.27.6 30.8 6.8	F.10 613.2 116. 407 237.6 18. 19.6	F.(5) 17.2 3.6 17.2 3.6 17.2 3.6 17.3 3	F.(5) 328.4 [168.8 159 6 22.8 228	F.40 531. [(8, 6, 421 26, 21.2 + 8 F.2 152, 8124.1 33× 4.31.2 21.6 9.6	10.1 619.2 146.4 502.858.8 38.4 20 1	F.30 321 111 6 182 8 1 4 2 1 2 13 3 3	F.01 : 263.6 180.8 82.8 12.4 27.9 15 3	F.05 83.6 f02. 83.6 f0.1 38. 3.4	311.2 [81.2 [30] 27.3 24.8 25.1	319.6 195.6 121. 16.8 38.1 8.1	279 6 168. 111.624. 16.1 7.6	57. 57.6 46 506. 35. 51.	507 6 140 6 141 408 15 10 60 507 6 140 8 158 15 1 35 110 1	896.8 126. 770.8 37.2 18. 39.2	380 111.6 040.139.0 30.17.0	336.8 160 4 176.431.6 21.2 10.1 1
RESIDUE ON EVAPORATION.	Potal. Dis- Sus- Ponded. Sus- Ponded. Pisses Pisse	255 2 241.6 13.6 26.8 24.8 2.	209 2 [89.2] 20. 34.2 31.2	F.1 (288.8.8.27.6 61.229.6 26. 3.6)	7.1 214. 201.2 42.5.20. 20.	55. 156 × 101.3,81 30.153.6 I	17.4 220 4 161.6 GD 8-21.6 -20.8	570 112 50X 10 10 10 10 10 10 10 10 10 10 10 10 10	577.2 116.1 160.838. 15.6 22.1	259 2 125.6 233 5.27.6 30.8 6.8	F.10 613.2 116. 407 237.6 18. 19.6	913 6 110.4 803.2 16.1 16.8 29.6	F.(5) 328.4 [168.8 159 6 22.8 228	152.8 124.1 328.4 31.2 21.6 9.6	16.1 502.858.8 38.4 20 1	F. 30 34.1 2 130.4 14.20 11.4 21.2 23.3	1.01	275 6 102 83.6 40.1 38 2.4	311.2 [81.2 [30] 27.3 24.8 25.1	319.6 195.6 121. 16.8 38.1 8.1	279 6 168. 111.624. 16.1 7.6	57. 57.6 46 506. 35. 51.	270. 15.6 121.45% 12.1 15.61 27.6 6 110 6 128 16 1 36 10 11	896.8 126. 770.8 37.2 18. 39.2	380 111.6 040.139.0 30.17.0	336.8 160 4 176.431.6 21.2 10.1 1
RESIDUE ON EVAPORATION.	Total. Polyos-sid Sus-bended. Disconding Sus-bonded. Prof. From Sus-bonded. Sus-bonded. Sus-bonded.	255 2 241.6 13.6 26.8 24.8 2.	M. F.05 209 2 189.2 20. 34.2 31.2	F.1 (288.8.8.27.6 61.229.6 26. 3.6)	M. F.1 214, 201.2 42 8.20, 20,	N. P.4 30.1558. 156 × 101.2,81 30.153.6 I	N. F.4	570 112 50X 10 10 10 10 10 10 10 10 10 10 10 10 10	N. F.5 577.2 [16.1 19.838. 15.6 22.1	N. F. S.	M. F.10 613.2 116. 497 237.6 18. 19.6	N. F.(5) 280 132 128 17.2 3 6 1 1 1 2 8 2 9 6 1 1 1 8 8 2 9 6	N. F.(5 328.4 [168.8 159 622.8 22. 8	F.20 531 169 6 421 136 21.2 4 8 F.2 152 8124 1 328 431.2 21.6 9.6	M. P.1	F. 30 34.1 2 130.4 14.20 11.4 21.2 23.3	M. P.01 263.6 180.8 82.8 12.4 27.2 15 2,	F.05 83.6 f02. 83.6 f0.1 38. 3.4	16 1. F.03 311.9 181.2 130. 27.3 24.8 2 1	T. F.05 319.6 195.6 121. 16.8 38.1 8.4	T F.02 279 6 168. 111.624. 16.1 7.6	7. F.O 574. 127.6 146 856. 52. 141.	270. 15.6 121.45% 12.1 15.61 27.6 6 110 6 128 16 1 36 10 11	N. F. 1	N. F.5 382 111.6 540.139.2 35 17.5	336.8 160 1 176.4 31.6 21.2 10.1 1
RESIDUE ON EVAPORATION.	Turb'y. Total. Odo: Piss- Solv'd. Piss- Solv'd. Piss- Portyl. Piss- Portyl. Odo: O	* (Cons.d M. F.04 255.231.6 13.636.8 24.8 2. ** N. F.07	4 N. F.05 200 2 189.2 20. 34.2 31.2	M. F.06	* M. F.1 214. 201.2 42 520. 20	* Much M. F.4 555 7 156 7 101.031 30.153.6 1	* ('ons'd M. F.4 325 + 161.6 (9) 8 21.6 30.8	670 113 55 40 16 68 68 113 55 40 16 68 68 68 68 68 68 68 68 68 68 68 68 68	* Much M. F.5 577.2 [116.1 460.838. 15.6 22.1	* Conv. N. E. S. 241 132 245 16 14 24 25	* Much M. F.10 613.2 116. 407.237.6 18, 19.6	8 CONSTITUTE OF THE STATE OF TH	* ('ons'd N. F.65 328.4 168.8 159 6'22.8 22	* Much M. F.40 531. 109-6-121 126. 21.2 1 8 * Cons.d. M. F.2 152.8124.1 328.431.2 21.6 9.6	\$ V.M'ch M. E.1	* ('ons'd M. F.07 321 111 6 182.841.4 21.2.23.0	* N. P.01 263.6 180.8 82.8 12.4 27.2 15.2	* N. 12.05 275.6 192. 83.640.1 38. 2.4	* Little T. F.02 311.2 181.2 180. 27.3 24.8 21.1	* 17. F.05 319.6 195.6 121. 16.8 38.11 8.4	* T F.02 279 6 168. 111.624. 16.1 7.6	# ('ons'd N. F.02 574.4 127.6 446 856. 53. 24.	* 134 10 11 10 10 10 10 10 10 10 10 10 10 10	* Consid M. F. 1 896.8 126 770.857.3 18 39.2	* N. F.3	§ Much M. 336.8 [60 4 176.431.6 21.2 [6.4]
APPEARANCE. RESIDUE ON EVAPORATION.	Turb'y. Total. Odo: Piss- Solv'd. Piss- Solv'd. Piss- Portyl. Piss- Portyl. Odo: O	5 * Cons'd M F.04	M. F.05 209 2 189.2 20. 34.2 31.2	Consed M. F.06 288.8 27.6 61.2 29.6 26. 3.6	V M. P. V. M. P. S.	* Much M. F.4 555. 156 × 101.0×1 30.153.6 I	* ('ons'd M. F.4 325 + 161.6 (9) 8 21.6 30.8	VN 15.15	Much M. F.5 577.2 [116.1 40.838. 15.6 22.4	* Conv. N. E. S. 241 132 245 16 14 24 25	13 * Much M. F.10 613.2 116. 407 237.6 18, 19.6	8 (ons.d M. F.65	3 * ('ons'd N. F.65 328.4 168.8 159 622.8 228	10 * Much M. P.40 531. 105.6 421 126. 21.2 4 8 17 * Cons.d M. P.2 152.8 121.1 328, 431.2 21.6 9.6	S V.M'ch M. E.1	* ('ons'd M. F.07 321 111 6 182.841.4 21.2.23.0	14 * M. F.01 263.6 180.8 82.8 12.4 27.2 15.2	M. P.05 275.6 192, 83.640.1 38, 2.4	* Little T. F.02 311.2 181.2 180. 27.3 24.8 21.1	* 17. F.05 319.6 195.6 121. 16.8 38.11 8.4	* T F. (2) 279 6 168. 111.624. 16.1 7.6	# ('ons'd N. F.02 574.4 127.6 446 856. 53. 24.	1. N 1.03	* Consid M. F. 1 896.8 126 770.857.3 18 39.2	* N. F.3	6 § Much M. 336.8 ltn + 176.431.6 21.2 10.1 I
RESIDUE ON EVAPORATION.	1900 Nedi. Color. Dis-pended. Portil Dis-pended. Portil Dis-pended. Sus-pended. Sus-pended	4 Jan. 5 * Cons.d M. F.04 255.231.6 13.636.8 24.8 2.	4 N. F.05 200 2 189.2 20. 34.2 31.2	25 * Much M. F.D. 322. 164.8 Jan 531.6 20, 11.6 1, 3 * Constd M. F.1 288.8 27.6 61.2 29.6 26, 3.6	* M. F.1 214. 201.2 42 520. 20	* Much M. F.4 555 7 156 7 101.031 30.153.6 1	3 Mar. 3 * ('ons'd M. F.4 323 4 161.6 (9) 8 21.6 20,8	670 113 55 40 16 68 68 113 55 40 16 68 68 68 68 68 68 68 68 68 68 68 68 68	22 2 Much M. F.5 577.2 116.1 (6).838. 15.6 22.1	29 *	13 * Much M. F.10 613.2 116. 407 237.6 18, 19.6	19 * (ons d M. F. 65	2 May 3 * ('ons'd M. F.65 328.4 168.8 159 6'22.8 228	10 * Nuch N. P.40 531. 108 6 121 136. 21.2 + 8	24 \$ V.M'ch M. P.1	51 * NHCH M. F.07 321 111 6 182.8 11.4 21.2.2.3.3.	14 * M. F.01 263.6 180.8 82.8 12.4 27.2 15.2	* N. 12.05 275.6 192. 83.640.1 38. 2.4	* Little T. F.02 311.2 181.2 180. 27.3 24.8 21.1	* 17. F.05 319.6 195.6 121. 16.8 38.11 8.4	1× * 19 *		* 134 10 11 10 10 10 10 10 10 10 10 10 10 10	1 * Const M. F. 1	* % * N. F.3	. 5 Sept. 6 \$ Much M. 336.8 http://doi.org/10.1111
APPEARANCE. RESIDUE ON EVAPORATION.	Turb'y. Total. Odo: Piss- Solv'd. Piss- Solv'd. Piss- Portyl. Piss- Portyl. Odo: O	5 * Cons'd M F.04	17 1 1X + 1	25 * Much M. F.D. 322. 164.8 Jan 531.6 20, 11.6 1, 3 * Constd M. F.1 288.8 27.6 61.2 29.6 26, 3.6	* M. F.1 214. 201.2 42 520. 20		Mar. 2 Mar. 3 * ('ons'd M. F.4 222 4 161.6 60 8 21.6 20.8	670 113 55 40 16 68 68 113 55 40 16 68 68 68 68 68 68 68 68 68 68 68 68 68	21 ' 22 * Much M. P.5 577 2116.1 190.838. 15.6 22.1	25 4 2 2 3 4 2 2 3 3 4 2 2 3 3 4 3 4 3	12 " 13 * Much M. P.10 613.2 116. 497.237.6 18. 19.6	19 * (ons d M. F. 65	May 2 May 3 * ('ons'd M. F.65 328.4 [68.8 159 622.8 228	10 * Nuch N. P.40 531. 108 6 121 136. 21.2 + 8	23 . 24 \$ V.M.ch M. E.1 619.2 116.4 502.858.8 38.4 20 1	51 * NHCH M. F.07 321 111 6 182.8 11.4 21.2.33.3	. 13 . 11 *	20 · 31 * 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3 × 3	* Little T. F.02 311.2 181.2 180. 27.3 24.8 21.1	T. P.05 319.6 195.6 121. 16.8 38.1 8.4	: 1× · 19 * · · T F.02 279 6 168. 111.624. 16.1 7.6	Ang. 1 Ang. 2 * Cons a N. F.03 574-1 121.6 116 256. 32.	. 9 * 13 * 13 * 12 * 13 * 13 * 15 * 15 * 15 * 15 * 15 * 15	29 · 21 * Constd N. F.1	. 29 : 30 * N. F.C 382, 111.6 240.139.2 42, 17.0	Sept. 5 Sept. 6 \$ Much M. 336.8 ich 1 176.431.6 21.2 10.1 1

TURRIDITY *Decided. *Very Decided. *Distinct. *Very Slight. Slight. Color. M., Muddy. V.M., Very Muddy. T., Turbid. C., Cloudy. Color on Interior Color. Brown. Dr., Durk Brown. LB., Light Brown. RB., Reddish Brown. BC., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White.

TABLE 68

STREAMS EXAMINATION--CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM MISSOURI SHORE, ALTON.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

No. of Bac. per Cubic Centi- meter.	15.20 16.00 18
Presence or Abs. of Coli.	111+11+11++1111++++++++++++++++++++++++
Temperature of Air, Cent.	400000000000000000000000000000000000000
Water. Temperature of Water, C.	1, - 16 16 16 17 17 17 17 18 18 18 18
Nitrates.	81351883888888888888885151524484885155485888
Signature S.	282825225252525252525252525252525252525
pended,	4 4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Onganic Dis- solved. Sus-	24
One Sid	8 % 4 % 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
ig b'breq T'otal.	1
A Sus-	88 2 2 3 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Anabuminoid Albuminoid Tot'l Fist	
AMD AMD TOTAL	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Free Am-	4.50 4.50
By Suspd	-31-49
Total. OON WAS TOTAL OON WAS TO WAS TOTAL OON WAS TOTAL O	
Total. C	0.000
Chlorine,	<u>ાં તે મેં જે છે છે જે એ મેં મેં મેં કો કો જે મેં જે જે જો જો જો મેં મેં મેં મેં જે મેં મેં મેં જો જો જો જો જો જો કો કો કો કો </u>
Color on Igni- tion.	
Loss on Light Constitution of Loss on Light Constitution of Light	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Tor.1 Loss o	राम क्रम रक्षिम रक्षिम कराखरामरा म रू रा म न
RESIDUE ON EVAPORATION. Disconding Disconding Colors on 1gr Colors on 1	23.6 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Dis- B	#\$387.536.83888888888888888888888888888888888
T'otal.	200. 8 244. 4 220. 8 246. 8 220. 4 200. 4 220. 6 1797. 2 220. 2 17
Odor.	
ANCE.	MANANA MA
APPEADANCE. Sedi- nient.	Cons'd Wheeh Cons'd Wheeh Weeh Wheeh Cons'd Wheeh Wheeh Cons'd Wheeh Wheeh Cons'd Wheeh Wheeh Wheeh Cons'd Wheeh
G ig ig	で 二 元 代 以 の
1900 Exami- nation.	Jan. Feb. Mar. April May Lune Lune Lung
DATE OF 1900 19 Collec- Exa tion. nati	Jan. 18. 18. 18. 18. 18. 18. 18. 18. 18. 18
Serial No.	6624 1 1 1 1 1 1 1 1 1

TURBIDETY-*Decided. §Very Decided. † Distinct. † Very Slight. | Slight. | Color | Colo

TABLE 69.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.
SOURCE OF WATER-MISSISSIPPI RIVER, 400 PEET FROM ILLINOIS SHORE, AT CHAIN OF ROCKS, PUNPING STATION NT. LOUIS WATER WORKS.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

No. of 18ac.	per Cubic Centi- meter.	8.8.8.9.4.6.8.5.4.4.4.8.4.4.4.8.8.4.4.4.8.8.4.4.4.8.8.4.4.4.8.8.4.4.4.8.4
:ent. :e or	Tempers of Air, (Presence Abs. of	+ + · · · · · · · · · · · · · · · · · ·
er. nture er, C.	Helgle Water Tempera of Water	$3 \times 3 \times 4 \times $
TRO- GEN AS	Nitrates.	515 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
NITRO- GEN	Zitrites.	00000000000000000000000000000000000000
IC EN.	Sus-	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
ORGANIC NITROGEN.	Dis- solved.	25.50
ON	Total.	\$25.55.55.55.55.55.55.55.55.55.55.55.55.5
200	Sus- Sus- prd'd	100
EN A	Albuminoid 170T sid b'vlos	25-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1
NITROGEN	I'10'T	844884 10044488 1004488 100448 100448 1004488 100488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 100448 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 1004488 100448 100448 1004488 1004488 1004488 1004488 1004
Z	Free Am- monia.	44255455555555555555555555555555555555
SN IED.	By Suspd	$\frac{\omega\omega\sigma_0\overline{\omega}_1\overline{\omega}_1+\omega_{\ell_1}+\omega_{\ell_2}+\omega_{\ell_2}\varepsilon_1\overline{\omega}_2\overline{\omega}_2\overline{\omega}_1\overline{\omega}_1\overline{\omega}_2\overline{\omega}_2\overline{\omega}_1\overline{\omega}_1\overline{\omega}_2\overline{\omega}_1\overline{\omega}_1\overline{\omega}_2\overline{\omega}_1$
OXYGEN CONSUMED.	By Dis-	Φ — 17 ∞ 13 ∞ ω 13 ∞ ω 12 ω ω 17 + 14 — 21 f ω 13 ω 12 Φ + 10 π + 10 π + 10 Φ + 10 π + 10 Φ + 10
	Total.	<u> </u>
	Сијог	<u> </u>
Colo	on Igni- tion.	数数数数数数数数数数数数数数数数数数数数数数数数数数数数数
.X.C	s sns p,pad	8.39 19.30 1
RATI	Loss on 1.10 T	$\frac{99}{8} \frac{1}{8} 1$
VAPO	3 1.10,1,	8.91-86-66-88-88-88-88-88-88-88-88-88-88-88-
RESIDUE ON EVAPORATIO	Sus- sus-	298. 273.2 24.8 38.3 374. 191.9 17.2 38.3 374. 191.8 170.2 38.3 174. 191.8 170.2 38.3 174. 191.8 170.4 4.9 170.2 38.3 170.2 4.9 170.2 39.3 170.2 4.9 170.2 4
IDUE	Dis.	298 273.2 24 179 179 179 179 179 179 179 179 179 179
RES	Total.	288 241 250 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
• .	юрО	
ANCE.	Color.	MANAMAN MANAZAMANANAZAMANAN MANAMAN MA
APPEARANCE	Sedi- ment.	Much N.W.ch N.W.
	.X.qana.	* * * * * * * * * * * * * * * * * * * *
OF	1900 Exami- nation.	Jan. 123 124 125 125 125 125 125 125 125 125 125 125
DATE OF	ec- E	BEST STEER S
	1900 Collection.	May. Nprij
No.	Serial	

Color on Ignition G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rb., Reddish. W., White Colon-M., Muddy, VM., Very Muddy, T., Turbid, C., Cloudy. Trubmity * Decided. § Very Decided. + Distinct. + Very Slight. Slight.

TABLE 69—Concluded.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISHPH RIVER, 400 FEET FROM ILLINOIS SHORE, AT CHAIN OF ROCKS, PUNITING STATION ST. LOUIS WATER WORKS.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois,

7,500 17,000 16,000 10,000 1,800 3.000 Bac. per Cubic Centi-meter. No. of Presence or Abs. of Colf. Temperature of Air, Cent. Temperature of Water, C. Height of Water, GEN AS Nitrates. 00024538488°8955 NITRO-800 9 900 800 000 Nitrites. рәриәд ORGANIC NITROGEN. solved. Dis-899 92 856 696 4.32.82.82 66£ 33xx88 92.0 22 Total. p,pud -sng Albuminoid Am. NITROGEN AS 808 AMMONIA. p, rjos .384 352 ['30T 136 136 136 136 136 136 88=88 Free Am-monia. By Suspd Matter. CONSUMED OXYGEN 13.9 9.3 16.9 13.0 16.9 13.1 22.1 17.3 24.8 17.3 By Dis-solved Total. 7-888.07.0 4. 80.00.0 Chlorine. Igni-tion. Color on p,pud -sng Loss on 1g'n. RESIDUE ON EVAPORATION. 139.640.8 136.828.4 596.854.4 213.257.2 154.4 32.4 185.2 36. 134.4 31.6 161.6 60. 65 H हा हा हा ∞ 1,10,1, 350.4 191.2 159.2 202.8 163.6 390.2 8 163.6 391.2 1492.2 6 542.7 552.2 1452.2 376.8 542.4 162.8 292.4 162.8 292.4 162.8 164.7 pəpuəd -sng 1020.8 155.6 E 576.2 155.6 E 592.4 148.8 5778.3 171.6 4 332.2 156. 330.8 176.8 330.8 176.8 136.9 165.8 1 156.4 solved Dis-Total. Odor. FF.05 F.02 F.02 F.03 F.01 Color APPEARANCE. NANANANHHH HERERERERERE V.M'cb Sedi-ment. Cons'd Cons'd Much Cons'd Much Cons'd Little Much Much Turb'y Exami-nation. 1900 DATEOF 1 Oct. : : : : : Collec-384488453 8885555568 1900 tion. 8581 Oct. 8618 ··· 8117 8161 8184 8198 22.22 22.22 25.22 Serial No.

Bh., Brownish. R., Red. Rh., Reddish. W., White. C., Cloudy. T., Turbid. VM., Very Muddy. COLOR ON IGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Coron-M., Muddy, | Slight. TURBIDITY-* Decided. § Very Decided. † Distinct. † Very Slight.

TABLE, 76.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS—PARTS PER MILLION.
Source of Water-Mississippi River, Midstream, at Chain of Rocks
Pumping Station St. Louis Water Works.

Report of Arrhur W. Palmer, T. J. Burrill, University of Illinois.

No. of	per Culbio	Centi- meter.	13.000	13.500	192,000	3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3	123.95	103.000	129,000	59.000	97,0(4)	000.00	000.65	3000000	24.500	33,000	53,000	0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.	000.63 20.030	(100), 16	00200	13.500	20.000	25.000	24,000	38,000	12.000	61.08.0	000 22	23.500
		Threse		+	+	+	-	+	1	1	•	+-					,	+	+	 †-	: +		:	:	•	1	1	+		
	Temperature of Air, Cent.		ic. #	-		- 1	, T	10.		; - ;	}- '	e i g	3 -	7	$\frac{\infty}{2}$	33.			± 5 5 5	7 6			:	:			:	:		
97111		TempT W To	.0 .0	0	 	- i-	0	-	oi.	<u>.</u>	72 .	: : : : :	3 12	1 x	~	_	30.	A) .	-tr 2	- ^	97.6		× 22.	6.27.	3 26.	1.27.			. 5 ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° °	8 26.
10	ußi	He				o o	-	_		-											13.9		<u> </u>	2	=	_		5. 0		
TRO-	.89.	Nitrat				6 G		_					_	4								56	19.	9.	E.	-		- 3	3 x	33.
NITRO-GEN	·sə·	HHIN				88													3 3					-		100.	-		9.0	9.
IC EN.	.bo	sus Sus	.33	£		2.16	1.568	3.04		5/	130	35.5	3.6		1.35 	1.216	1.76	200 · ·		1.076	2.676	2. IGH	2.212	300 3	1.48	39.	E03:	757		2.001
ORGANIC NITROGEN.		Solve	.53	٠ ١	416	$\frac{\pi}{2}$.512	.64	9.	ာ	100	. X		38	333	324	22	9i 9	202.	7	1 2	.516	33.	37.5	27.2	or.	987	21.00	101	. 136
No.N	J. It	NoT'	₹.	3.0	xç ğ	5.55	80.3	3.68	10.0	1.304	1 3	2.12	9 6	33	1.5.1	1.54	20 9	33.63	2 d	388	26:00	25. CS	14.8	= ::	.73	<u> </u>	-:	3 3	1.61	5.4
20	<	p,pud -sng	80.	11	.192	000	.576	1.264	. 432	8	3330	010.	544	× ×	.56	.G	.576	576	0.46	200	1.024	XX.	1.056	.061	1961	<u>x</u>	146.	2 10	2	1321
4 ;	Albuminoid	Dis-	.176	200	250-	. 16	1224	.176	77.	272	176	2003	200	38	.192	.16	.176	91.	220	200	.176	.113	11.	. 15	300	=	X :	0.1.0	113	25
NITROGEN	Albur	Tot'l	.256	51.0	XX 6	3 %	တင္	#	67.5	207	. 5 I G.	50.00	736	.61	.752	×.	.752	989	200	96.	63	96:	3.	176	.576	505	100	A SE	. 26.	.021
7		Free A	.304	890.	× ×	355	22.2	2561	£ 5	3000	200	200	9.5	800	840.	:0:	.052 .052	X :	010.	030	?}	90.	<u>.</u>	-	689	970.	260.	8 5	2 2 2	.056
SD.		By Sus	00 61	<u>∞</u> ,	0.0	- i c	3	00	9.0 63.0	x 0		C 4	o	5.5	8.9	9.1	٠ ١ ١	9.	2.0	=	10	∞. 1G	6.3	_	ात : करे	-:	x :	3) X	??	-
OXYGEN	.ba	solv solv		63.6 (C)		20:	- j-	∞ ;~	 च्याः ऽ १०० ३	x .	- 0	T 10	5 50	1	5.5	£	∞ :	0.0		0.0	5 9 1	6.6			33	0.5	?? : ;	n ==	6.6	≎! ∞
Cox		RioT	9.3	∞ ;	200	19.6	18.6	26.3	্ন ন	27.0	20.5	21.2	. 2	- C	19.3	<u>8</u> .8	<u>₹ 0</u>	5:2	910			20.4			16.7		1 2	2 - 2		19.6
.91	1110	СРЈ	33.	₹~ ₹ 50 \$	- 000		9.8				÷:	0 0 0 10	-	5.6	. č.	6.	တ္ ၁	x o	C 1-	- } -	6.1	6. 4	oc i	7.	x :	- :	- c	C 1U		7.5
Color	on Ieni-	tion.				<u>i</u> ==i				2.0	200	<u>:</u> =	-	~	7.		~ :		<u>-</u> -	-	13.	7:	≃;	· .	2			<u> </u>	~	13.
ż		p _i pud -sng	6.8	च्याः व्यः	0.0	39.6		16.4			0.75		· · ·	000	- C	?? ???	51 -	9.6	1. R. S.	9.11							G 1		3	× ×
NATIO	101	sid b'vios	G	± 0 ∞ ≀		9	30	2.	300	0.4	mps.	-	. 23	19.6	29.6	30.00	26.	× 05.	90.10	8.01	30.	00	16.4	13.6 58	89. 63.	02 × 40	21.6 60	3 61	2)	
APOL	Loss	1,10,1,		x:02:	0.10	£	9 11	64.8	31	20° F	3.5	2.5	2 21	159	÷5		33.3	3 5	9 (1 4401	30	oc.	35 (3	7	7.5	200	11.6	6 101.	80.
RESIDUE ON EVAPORATION		puod sng	26.4	47. 5 47. 5 47. 5	2 α 2 α 3 α 3 α 3 α 3 α	111.4	100	301		mbre	555	a.	201 P. 202	673.6	619.2	1016 156	0.95.661	1010.4 0101	3119 1		•					ardit.	1514. 07	16:29:2 11	2365.6	-
DUE O		ovios -	303.6	. 3	, ; -	- 53	21	??	:5	- ·	#H. 6	41 000	n .	.5	_	x	0			or.	198.801	1965.8	196 1 2580.	7.6	100.			200	8	50
RESU	-	rioT'		410.4.408	030 C-010	663.6 219	3		1 3 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		70.01		01	??	1.3 17	9.2 515	1.6 206.	X65 22 22	2 ~	4 4-04	-	∞. ≥= ∞.		9	: ود	٠ و	- 0	7 23	23	©3
			530	4:				1210	× :		-	1001	1017		7.91	1559	1001	1000	933	2001	. 2100.	3.00		2011	100	50.	5 2 5	25.	2000	. 1537.
	1019	0	:	:	:			:	:	:		:				:	•	:				:	:	:	•	:	:			:
		or.	F.03	30 31 31 31 31 31 31 31 31 31 31 31 31 31	30.7	F.03	F.0.5	S: :	ا ا ا	E 8	2012	200	88	F.(165	F.50		0) S	93	000	39	₹.10	E. C.	36	0.0	10.7	20%	1 3 3 M	30.2	10.	·.05
RANCI		Color.	N.	7.7	. 7		7.		7. 7				-	7.	N.	7.	7. 7	1.		1.1	N.					7. 7	17			N. I
APPEARANCE	2000	ment.	p.suo.	i. dent	HCII ;;	,	*	. M.cb	Much	9.		:		9 .	**	.M.ch	40		9 9	.M.ch	Cons'd	:		0		: :		:	*	9.0
	1.7.	1,111,	* *	* *	+ 4+	.//.	.11.	·DC :	lb 41	10	*	*	*	*	*	.00	0.4	t 31	- 00	. 10.	*	*	* 1	H -	₩ 4	+ 4	6 4	*	*	*
<u>G</u> .	1900	Exami- nation.		223	1. C. 1.	Feb. 9		ar, o				06 9	100			£ !	65 - C	7	1	1000	\$? ;	uly 6	<u> </u>			77	0 -0	3 51		
DATEOF		c. E.	5 Jan.		96		3	7.		-	7.00	101	300	3 May	. 01	-		-			· ·	-	on y	- :		ر د م	× 500	3 22	155	:
Us	1900	Collec. Exami- tion. nation.	Jan.	: :	*	reb.		Mar.						May			: :	3		9	0.5	200						1 21	:	
1000	[B]:		3633 J	6670	61.67	6567 F			1150		Carry Com	2523	1126			(26)	100 B	10101	7	1911			700	1812	1922	N. Comp	202	100%	8050	6036

TUBBLUTY-*Decided. \$ Very Decided. † Distinct. † Very Slight. Slight. (OLOR-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. Color on Ishinon-G., Gray. B., Brownish. R., Red. Rh., Reddish. W., White.

TABLE 70—Concluded.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Peport of ARTHUR W. PALMER,

APPEARANCE.

DATEOF

Color.

Sedi-ment.

Lan,L

Collee-Examination.

Serial No.

Odor.

of Illinois.	No. of Bac. per Cubic Centi-meler.			6,000	10,00	4,000	16,06	6,000		1.8()	- 1986 -	2.150	1.900	£	7,30)	19.400	0.600		3,100	œ æ	3 3 3 5 5 5	29.100	500 ti	3.400
LL, o		Presence or Abs. of Coli.				+-		-;	pr. 0				, -	1		7							- 1	1
T. J. BURRILL, University of I	Temperature of Air, Cent.					:		:		:										:				
1.5	Temperature of Water, C.			96.7		000	. 68		જું જું જું	29.5	g; ∞ (§)	.936.	9.26.		27	- 1			30.	∞ 9	. 03	3	္က ဂ	
	Height of Water,			63.7	10.929	⊕ ⊕ ∞ -	T T	6.1	5	ر د د د	σ. χ. σ.	∞		20.0	10.	T :	Ω (1·	- 1	6.1	ۍ ن و	10	5.6	ž c	:
	Vitrates. CEN A S			88 88	. e.	==	==	=:			e 15		9:	ج =			٠ د د		= = = = = = = = = = = = = = = = = = = =			56.	5. 5. 5. 5.	<u> </u>
	NITRO-	Nitrites.			.00						88	•	**	38	9.		3.5	• •		88	• •		38	8
	οż		-sng	1.992	966:	250	.676	1.00	367	98. 8.	$\frac{1.02}{1.116}$	1.292	768	21.5. 20.5.	.76		5000	.632	312	425	1.128	968	7364	1.536
	ORGANIC NITROGEN	.b	-si(I 97Ios	.348	324	5. 5. 5. 5.	324	جن جن چن چن	3I6.	ين _د	2 8 3	.188	395	0000	.416	: 3	9 5	. 355	384	.336	133	. 132	20.00	809.
OF ROCKS.	ON	J.	BioT	1.64	1.32	9.5		26.7							_		بشر				عبال		1.18	2.144
	NITROGEN AS AMMONIA.	Am.	p,pud -sng	.56	.432	1273 178 178 178	308	368	.368	.16	541	.35	528	967.	.464	515.	238	. 112	37.5	45.	513	809	188	<u> </u>
<i>;</i>		Albuminoid	b'vlos	176	.176	4.5	75	1.76	176	86.3	193	.16	11.300	11:3	.138	356	000	336	308	.16	88	.16	77.55	304
AT C KS.		Albur	1.10T	736	809	310.4	544	##¢:	tte:	£18	.736	.48	15.	809	.592	292	0140	148	×4×	7. 0	9.35	7.68	5 6 6	I.024
MIDSTREAM, AT WATER WORKS.	Z		A 9914 sinom	.044	07.0	000	10.	.960 -	133	.116	152	80.	.14	135	.113	961.	116	38	890.	36.		280.	2 21	<u>₹</u>
DSF	ED.	By Suspd		15.2 13.3	11.2	4.7	∞ }~	တ် (C	-	6.4 5.4	14.4	<u>ن</u>	e : = :	1 T	7:	10.0	ء ج ≥ ∞	, <u>i</u> -	7. C	(- c	- x	0.0	57 to	13.1
i, M	ONYGEN	By Dis- solved.		6.9	7- 70:		5.5	6. 2 I	6.9	တ္း) : : :	5.7	က် (၁	5 <u>1~</u>		ر ا ا	+ ± - ∞		9.4	ء و د	0.0		- 8. - 8.	12.6
RIVER, LOUIS	000	J·	Tota	20.2	18.7	15 16.	15.7	55 57 45 7	= :	20 20 20 20	20.0	18.4	2. 3 2. 3 3. 3	20.8	21.5	17.4 10.0	10.0 70.0	20.6	· · ·	 	.¥. 28. 18.	21.6	C 60	24.7
PPT RIVER ST. LOUIS	Chlorine,			5.4	30	0 0 0 0		က် က် က်		တ <u>~</u>		8.6	?? O	11.	10.3	0	0 0 0	5.00	5.6	4.0) ~	= "	o 4. `∞	5.
WATER-MISSISSIPPI RIVER, MIDSTREAN, AT CHAI PUNPING STATION ST. LOUIS WATER WORKS.	Color on Igni- tion.			æ. æ.	m ;	ವ ಜ	<u> </u>	ವ ವ	i pi	≃ ≃	i	ii i	2 2 2 2	DE:	D E E	200		D13.	DE:			<u> </u>		
S ST	ż	: n.	p,pud sng	30.4	68.4	33.54	18.8 2.8		21.6	.0	51.5	27.52	27.6 7.7.6	10.8	39.6	0	ر ان ان	<u>∞</u>	हा !	17.6	36.4	93.6	0 00	55
VATEI	(ATIO)	Loss on 1g	Dis.	33.2.28.30.8	26.	97.0	26.4	36.6	24. 21.	17.6	300	 	\$ \frac{1}{2} \frac{1}{2} \f	19:93 19:01 19:01	16.4	T 0	2 2		9	3.55	38 i 32 i 30 i	26.8.33.6	310	8.83
OF V	APOI	Los	1,10,1	. 23 19	74.4	# E	45.2	54.4	45.6	41.6	10.4	40.	25.5	36.4	46.	40.4	3000	36.	00 s	43.0 97.0	59.5	50.1	54.00 54.00	56.
SOURCE OF WATER PUNPING	RESIDUE ON EVAPORATION		Sus	914.4	546.	377.6	331.2	303.6	210.	201.841	673.2	627.6	628.4	579.6	581.4	197.4	418.8.37.6	337.2	291.252	254.8443.6 670 5 55 6	076.8	630.	452.	974.8
So	DUE		Solve Solve	2.1 [48. 6.8 [49.6	51.6	161.6 16.16		20 00 20 00		101	130 8	158.	146. 173. e.	138.8	80.5	2.1.2		7	25.50	. 6	229.61	195.2	150.1	170.8
	RESI		rio'T	1062.1 836.8	697.6 151.6	506 538 538		502.8		819.2				718.4		0.087	جبيات		्र	212.00	1306.4	835.3	602.4 150.1	
			: :	:		:	: :	: :	:	: :	:	:	: .	:	:	: :		:	:	: :	:	: :		

Cons'd M. F.02

M. F.02

M. F.02

M. F.02

M. F.02

M. F.03

M. F.03

M. F.04

M. F.05

M. F.

Sept.

: % 55

8116 8149 8159 8153 8196 8224

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. ‡ Very Slight. | Slight. Turbidity-* Decided. § Very Decided. † Distinct.

Ξ.8

8571 " 8582 Oct. 8617 "

8481 8498 8516 8555 8550

TABLE 71.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burkell. University of Illinois. SOURCE OF WATER-MISSISSIPPI RIVER, AT CHAIN OF ROCKS, INLET TOWER, ST. LOUIS WATER WORKS.

o. of	per Cubie Centi: meter.	188 188 188 188 188 188 188 188 188 188
A bs. of Colf.		
10.0	Ттеменс	
	Tempera Of Alr, C	+ +- 0 = 1- 1- 0 8 5 2 2 8 5 8 5 8 5 8 5 8 5 8 5 8 5 8 5
9mn	Tempera	
	dgioH otrW	9) 20 9) 10 9) 20 10 20 10 20 10 10 10 10 10 10 10 10 10 10 10 10 10
TRO- GEN AS	Sitrates.	
NITROGEN	Sitrites.	88 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
o. N.	-sns Deuged	8. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3.
ORGANIC NITROGEN.	Dis-	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
QN EN	Total.	<u>~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~</u>
	i p,pud -sns	28 2 3 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
N AS		五 6 2 6 3 6 4 6 4 6 4 6 5 6 6 6 7 6 7 6 8 7 8 7 9 7 9 7 10 7<
NITROGEN	Albuminoid froft for	888488848848484848484848484888484888888
NITA	monia.	4 + 1 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2
	Pree Am-	
OXYGEN	By Suspd	「 「 「 「 「 「 「 「 「 「
OXYGEN	By Dis-	$\frac{(d-1)^2}{(d-1)^2}\frac{d}{(d-1)$
Total, C		$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Chlorine,		ကြည်း တို့ ရှိသို့ လိုက်ကို ကိုမေတာ်မှ မေတာ့ တည်း တည်း ထို မြိတ်မြိတ်မြိတ် မြိတ်မြိတ်မြိတ် ကိုမေတာ့ မြိတ်မြိတ်
Colo	on Igni- tion.	
N.	i p.pud	8527 + 18
RATIC	Loss on 1g	
APO	1.10.L	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RESIDUE ON EVAPORATION	popuod sus-	+ + + + + + + + + + + + + + + + + + +
OUE (Dis- solved.	$\alpha + \omega + \infty$ and $\omega + \omega$ and $\omega + \omega$ and $\omega + \omega$
RESII		367 2 342 8 619 6 362 8 619 6 362 8 1850 210 8 1015 220 8 957 6 123 9 957 6 123 9 1012 8 173 6 1171 8 171 8 1371 6 20 1 1371 6 10 0 1371 9 10 0 1371 1 10 0 1371
	Total.	18.00 18.00 19
.robO		
APPEARANCE.	or.	######################################
	Color.	AND
PEAR	di- int.	
AP	Sed1.	Nuch W.W.ch W.W.ch W.W.ch Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd Cons'd
	Turb'y.	- 5 5 6 5 6 5 6 6 6 7 7 7 8 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9
OF	1900 Exami- nation.	July Nar.
DATEOF	0 ec. F	16-148 x 8 488 16 55 58 65 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
-	for0 Coffee- tion.	25
		66866999999999999999999999999999999999

TUBBIDITY * Decided. § Very Decided. † Distinct. † Very Slight. Slight. Color M., Muddy. V.M., Very Muddy. T., Turbid. C., Cloudy. Color on Interion—C., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. TURBIDITY * Decided. § Very Decided. + Distinct. + Very Slight. Slight.

TABLE 71—Concluded.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISSIPPI RIVER, AT CHAIN OF ROCKS, INLET TOWER, ST. LOUIS WATER WORKS.

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois.

·					
No. of Bac. Per Cubic Centimeter.			9,100		
	Presenc Abs. of	[++++++++++]			
	Tempera of Air, C				
	Tempera of Wate	ន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន្ទន	:		
	dgi9H eta7/	$\begin{array}{c} \overline{3}\overline{3}\overline{1}=\overline{0}\overline{0}\overline{0}\overline{0}\overline{0}\overline{0}\overline{0}\overline{0}$:		
TRO- GEN AS	Nitrates.	888888484448485000000000000000000000000	5		
NITRO	Nitrites.	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	100.		
oż	bended.	25.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.	. Clefin		
ORGANIC	Dis- solved,	1	10m		
0.7	Total.	8,			
	i p.pud -sng	0.00	ماممدا		
DEN AS	Tot'l Dis- bio Dis- bio Dis-	0 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1		
NITROGEN AMMONIA	Tot'l	6004 6004	1		
	Free Am- monia.	6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8			
	By Suspd	$\frac{2}{6}$	-		
OXYGEN	By Dis-	(1) (1) (1) (1) (1) (1) (1) (1) (1) (1)	?		
CON	Total.	84888955744188889888884989449988888 664469644696686666666666	7		
Chlorine.		$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	2		
olor	on Igni- tion.				
	. In nud	# 1888 0 1 1 1 1 2 1 2 1 2 2 2 2 2 2 2 2 2 2 2	-		
T10.	e b'vlos	838 88 88 88 88 88 88 88 88 88 88 88 88			
RESIDUE ON EVAPORATION.	Loss on l'10T l'10T l'10T l'10T l'10S l'10S	<u> 4 3 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0</u>	?		
Eva	bended.	223.4 8 1815.1 1818.8 8 8 8 8 111.3 1818.8 8 8 8 8 8 8 11.3 1818.8 8 8 8 8 11.3 1818.8 8 8 8 8 11.3 1818.8 8 8 8 11.3 1818.8 8 8 8 11.3 1818.8 8 8 8 11.3 1818.8 8 8 8 11.3 1818.8 8 8 8 11.3 1818.8 8 8 11.3 1818.8 1818.8 8 11.3 1818.8 8 11.3 1818.8 8 11.3 1818.8 8 11.3 1818.8 1818.8 8 11.3 1818.8 8 11.3 1818.8 8 11.3 1818.8 8 11.3 1818.8 1818.8 8 11.3 1818.8	2		
NO 3	-sns	8 1 1 1 1 1 1 2 2 2 2 2 2 2 2 3 3 4 4 8 1 1 1 2 3 2 3 3 4 5 2 3 3 4 5 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	3		
IDOIS	Dis- solved,	25	200		
RES	Total.	247. 6 172. 8 2231. 8 84 1981. 6 165. 6 1490. 1 80 155. 6 1490. 1 80 1257. 2 178. 4 1338. 8 82 1257. 2 178. 4 1338. 8 82 1257. 2 178. 4 1338. 8 82 125. 6 1055. 2 125. 6 1055. 2 125. 6 1055. 2 125. 6 1055. 2 125. 6 1055. 2 125. 6 1055. 2 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 1161. 4 125. 6 12			
Odor.		M'sty			
APPEARANCE.	ı;	<u> </u>	-		
	Color.	MINNE T PERFERENCE FERE ER			
	Sedi- ment.	Cons'd "" "" "" "" "" "" "" "" ""			
	·Y'din'l' & Ħ	***************	0		
DATE OF	1900 Exami- nation.	\$\frac{2}{80} \frac{2}{80} \fra			
	- Exe	288 284 25 25 25 25 25 25 25 25 25 25 25 25 25			
	1900 1900 Collec- Examition.	Aug. 141y 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3. 3.			
Serial No.		86888888888888888888888888888888888888	-		

TURBIDITY-* Decided. § Very Decided. + Distinct. ‡ Very Slight. ¶ Slight. © Slight. © Slight. © Slight. © Slight. © Slight. © Color on Jention-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rb., Reddish. W., White.

TABLE 72.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS PARTS PER MILLION.

Report of Arthur W. Palmer.
T. J. Burnell.
University of Himors.

SOURCE OF WATER-MISSISSIPPI RIVER, 400 FEET FROM MISSOURI SHORE, AT CHAIN OF ROCKS. PUMPING STATION ST. LOUIS WATER WORKS.

	*, *	66.180 66.180
No. o Bac.	tubie Centi- meter.	66, 160 66,
Golf.	Presenc lo ,sd A	
	Pempers of Air, to	
omn	rioqinol otisM lo	888888888888888888888888888888888888888
10 1	ranni daləti əts7/	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ITRO- GEN AS	Sitrites.	######################################
7.	sus-	+ (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2 (2
ORGANIC NITROGEN.	Dis- solved,	88999888889989999999999999999999999999
IO TIN	Total.	
s v	p,pud	900 900 900 900 900 900 900 900
	Albuminoid biological bivid	
NITROGEN AMMONIA	1'10'I	.,
	тее Ат. попів.	
GEN IMED.	solved. Sy Suspd Matter.	4rc 4 2 4
OXYGEN	Total.	10の1-の12以外の19日から19日かり19日かり19日から19日から19日から19日から19日から19日から19日から19日から
ne,	Chlori	8982895000000000000000000000000000000000
Color	on Igni- tion.	摄取线铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁铁
	i p,pud	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
RATIC	For Plass on 19, 10 Sus-	\$44774372005528828852448853855858
EVAPO	J	28. 28. 28. 28. 28. 28. 28. 28. 28. 28.
RESIDUE ON EVAPORATION.	Sus-	2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
ESIDU	-sid ,boylos	286. 8 248. 8 246. 8 24
2	Total.	25
r.	tobO	
CE.	Color.	######################################
APPEARANCE		REMEMBERS AND
APP	Sedi- ment	Cons'd Much Winch Wi
	i i i i	# * * * * * * * * * * * * * * * * * * *
DATEOF	Exami- nation.	May May
Dat	1900 Collection.	Jan. NP: : : : : : : : : : : : : : : : : : :
.o.X	Serial	6638 6930 6930 6930 6930 6930 6930 6930 6930

COLOR ON IGNITION-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. 8G., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy, Turbidity-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

Table 72—Concluded.

 N_0

Serial

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISSIPPI RIVER, 400 PEET FROM MISSOURI SHORE, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

University of Illinois.

Report of Arthur W. Palmer, T. J. Berrille.

36,000 2.900 10,100 47,950 27,100 26,750 15,000 6,000 6.000 5.000 4.000 1.000 per Cubic Centi-No. of Bac. meter, Presence or Abs. of Coll Temperature of Air, Cent. Temperature of Water, C. 8888889999877799967888 8-688899555877799967888 Height of Water. GEN AS £8282 NITRO-900 300 888 8 8003 003 900 9 89 8 9 11.528 .24 2.68 .244 3.076 404 pepued -sns NITROGEN. ORGANIC 809 288 188 188 solved -sid 375 [1336 372 [131 304 [92 224 [16] 1736 2 91 64 2 924 64 2 924 11.728 2.92 1.168 3.32 1.88 2.92 .896 2.6 .608 1.96 682242866 3 3 Total. 336 416 782 784 784 865 <u>∞</u> Albuminoid Am. NITROGEN AS .128 822.85.7 208 193 960 860 AMMONIA. 89999 84.948 056 056 168 37.6 88858 809 £218 1,10,1 98.75 88.88 051.064 Free Am monia. 3.7.19.6 4.916.3 4.919.1 4.20.2 5.20.1 11.8 19.2 6.4 12.8 11.6 15.4 5.1 10.3 13.2 16.8 8.4 8.6 12.4 20.7 7.4 13.3 27.6 24.9 7.5 17.5 15.4 26.4 6.8 19.6 16.2 23.8 6.3 17.5 3.9 16.3 4.3 17.3 4.5 15.6 4.9 92.5 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 4.1 21. Matter. 9.616.2 9.616.2 10.611. 15.211. 15.211. 17.211. 18.41. OXYGEN CONSUMED. By Dis-10. 27.7 6.621.8 9.424.9 8. 25.1 9.820.9 Total. Chlorine. Color gnition. on 37.6 5.5.4. 5.2.4. 5.2.4. ∞ p,pud sng Loss on ig n. RESIDTE ON EVAPORATION. 23.45 S S S 1039.6 139.6 810. 66. 21 976.4 125. 2 751.2 63.6 33 918.4 125. 3 618.8 16. 844.4 142. 3 618.8 16. 844.4 125. 3 618.8 16. 845.4 125. 3 618.8 120. .9. 234.8 168.8 2066. 110. 2040.8 170.8 1870. 109.2 1740.8 193. 1547.8 54. 1178.8 51.6 1591. 290.8 1303.2 34. 2148.4 285.6 1862.8 65. 1880. 282.4 1597.6 90. 1.10,1 384.4205.6 -sid bartos Total. Odor. E 0.03 E Color. APPEARANCE. zzzzzzzzzz ZZZZZ Sedi-ment. Much Much Much Larply * \$\\\ \= Collec-Examination. 1900 DATEOF Sea . Oct 36.2 2011255588 tion. 1900 8038 July 8052 ... 8076 Aug. 8583 Oct 8000 88888 850 850 850 850 8100 8518 8277 8386 S180 8499

Color on fention-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White C., Clondy. T. Turbid. VM., Very Muddy. Color M., Muddy. Trundity * Decided. & Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 73.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Arthur W. Palmer, T. J. Burrill. University of Illinois.

SOURCE OF WATER-MISSOURI RIVER, FORT BELLEFONTAINE, WEST ALTON, MO.

No. of Bac. Per Cubic Centi- meter.		21,000 399,000 43,600 43,600 37,200 7,500 1135,000 206,000 206,000 118,000 111
	Presenc	
ennie Jent.	Tempers of Air, C	では、ないで、可で、ないである。これで、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、では、
	renner otsW lo	
Helght of Water.		$-\omega + v \otimes v \otimes \omega \otimes \overline{\omega} \otimes \overline{\omega} + v \otimes \overline{\omega} \otimes \overline{\omega} = \overline{\omega} + \overline{\omega} \otimes \overline{\omega} \otimes \overline{\omega} + \overline{\omega} \otimes \overline{\omega} \otimes \overline{\omega} = \overline{\omega} \otimes \overline{\omega} \otimes \overline{\omega} \otimes \overline{\omega} = \overline{\omega} \otimes $
TRO- GEN AS	Nitrates.	8888288888888888888888888888888888888
NITRO- GEN	Zitrites.	8 5 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
S. S.	Sus- pended.	21 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
ORGANIC NITROGEN.	Dis- solved.	$ggggggggggggggggggggggggggggggggggg$
OF	Total.	8 8 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	g p.pud	\$\frac{1}{2}\$\frac
SN AS NIA.	bio bivios	
NITROGEN	Albuminoid Dis-sid b'vlos	$\frac{8}{8}$ $\frac{1}{6}$ $\frac{1}$
Nir.	sinom.	98 98 98 98 98 98 98 98 98 98 98 98 98 9
	By Suspd Matter. Free Am-	
OXYGEN	By Disselved.	
OX	Total.	89999977777777777777777777777777777777
ne.	Chlori	$\frac{\partial \mathcal{B}}{\partial \mathcal{B}} \frac{\partial \mathcal{B}}{$
Color	on Igni- tion.	
	i p,pud	uullouuka
RESIDUE ON EVAPORATION.	o sid n b'vios	89.0.884-484-4454-5668-4888-8982-288-856-664-665-7-1-6-6-4-4-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8
PORA	Loss on 1g	# CH # CO CO CO # # CO
Eva	pepuad	4) - 2 10 2 - 10 10 10 10
E ON	solved.	# 386 6 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9
ESIDI	Dis.	45.4 + 3.6
H H	Total.	
лорО.		
APPEARANCE.	01.	FTFTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT
	Coloi	
	Sedi- ment.	ch c
	Turb'y.	W.Y. W.Y. W.Y. W.Y. W.Y. W.Y. W.Y. W.Y.
DATEOF		
	1900 Exami- nation.	Mar. Nepri
	1900 Collec- tion.	
		28 28 28 28 28 28 28 28 28 28 28 28 28 2
No.	Serial	88888888888888888888888888888888888888

Color on Ignition - G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Colon-M., Muddy, VM., Very Muddy, T., Turbid. C., Cloudy, TURBIDITY * Decided. & Very Decided. + Distinct. + Very Slight. Slight.

TABLE 74.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS—PARTS PER MILLION

Source of Water-Mississippi River, 100 Yards from Illinois Shore, Jefferson Barracks, Mo

University of Illinois.

Report of ARTHUR W. PALMER, T. J. BURRILL,

000:12 per Cubic Centi-No. of Bac. meter. Presence or Abs. of Colf. of Air, Cent. Temperature femperature of Water, C. - 83 83 10 10 30 00 Height of Vater, GENAS Nitrates. NITRO-584 336 02 616 341 008 5512 768 011 544 32 013 384 32 013 64 4 02 536 1.504 016 656 776 016 536 1.504 016 656 776 016 656 776 016 656 776 016 656 776 016 Sus-sug ORGANIC NITROGEN. Dis-solved, 1.02 | 1.44 1.02 | 1.04 1.03 | 1.04 1.04 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 | 1.04 1.05 Total. Albuminoid Am. NITROGEN AS AMMONIA. Pis-.386 .36 .861 Totil Free Am-monia. By Suspd Matter. OXYGEN CONSUMED. By Dis-solved. Total. Chlorine. Color Jgni-tion. on 255.88 255.88 255.88 255.90 25 p,pud -sng Loss on 1g'n. RESIDUE ON EVAPORATION. Dis--sns solved. -sig Total. Odor. F OF Color APPEARANCE. Sedi-ment. Rurbyy * * Collec-Examination. 1900 DATEOF tion. 986 6614 Jan. 66712 " 67712 " 6781 Feb. 6874 " 7010 Mar. 7010 T191 " 7191 " 72243 Apr. Serial No.

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White, C., Cloudy. T.: Turbid. VM., Very Muddy. Colon-M., Muddy. Tunnintx-*Decided. §Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 75.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISHPH RIVER, EAST OF MIDSTREAM, JEFFERSON BARRACKS. M.

Report of ARTHUR W. PALMER, T. J. BURRILL. University of Illinois.

Illinois.	No. of	per Cubic Centi- meter.	1.800 24,100 26,800 38,400 38,400 140,000 133,000 187,500 187,500 38,000 38,000 38,000
13.01		Presence Abs. of 6	+++++++++++
unversity		Tempera Of Air, C	
C	ture r, C.	gradmall of Mate	
		Heigh Mate	- 55 51 10 10 10 10 10 10 10 10 10 10 10 10 10
	NITRO- GEN A	Nitrates.	
	7.	pended.	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
0.	ORGANIC NITROGEN.	solved.	25
s, MO	ORG	-siQ	5-0-8-6-4-4-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8-8
KKRACI		Fibrad F	87.50.00 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
N DA	AS	pio p.Alos	985 58 58 58 58 58 58 58 58 58 58 58 58 5
ERSO	NITROGEN AS AMMONIA.	Dis-	85574 55 55 55 55 55 55 55 55 55 55 55 55 55
JEFF	AX	Binom F 1'10'T	86 20 20 1 1 20 20 20 20 20 20 20 20 20 20 20 20 20
EAM,	OXYGEN CONSUMED.	Free Am-	0000 00400001-x
IDSL		solved. By Suspd	000001-1000000000000000000000000000000
30		Total.	0 1 1 4 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
LAST	Color fron. Chlorine.		24. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0.
11 E16,			最级的数据的数据数据数据数据
A LES	ż	i p,pud	#6####################################
SOLICE OF MAIER—MISSISSIFF	RESIDUE ON EVAPORATION.	FOLL DISCOULTS	# 10 # 22 # 24 # 2 # 2 # 1 # 1 # 1 # 1 # 1 # 1 # 1 # 1
	VAPO	1.10T	24 - 24 - 24 - 24 - 24 - 24 - 24 - 24 -
ALE	ON E	Sus- pended.	56.66.68.88.88.88.85.66.66.68.88.88.88.88.89.89.89.89.89.89.89.89.89
	HDUE	.sid .bovlos	8.68.28.28.28.28.28.28.28.28.28.28.28.28.28
	RE	Total.	339. 318.8 308. 239.6 309.2221.8 387.2221.8
	·,ı	орО	
		or.	25252525252525 2525252525252525 25
	RANCE	Color.	NAMES OF STREET
	APPEARANCE	Sedi- ment.	Much Cons'd Much Cons'd Wuch Cons'd Wuch Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Cons'd Much Much Much Much Much Much Much Much
	,	Lurb'y.	*****
	0 P	1900 1900 Collee-Exami- tion. nation.	Jan. 5 19 19 19 19 19 19 19 19 19 19
	DATEOF	66-E	- Jan
	I	Collection.	Jan. Feb.
11	.0%	Serlal	8615 6678 6678 6678 1113 6678 1113 1113 1113 1113 1113 1113 1113 11

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. Color M., Muddy. VM., Very Muddy. T., Turbid. C., Cloudy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 76.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of ARTHUR W. PALMER, T. J. BURRILL, University of Illinois.

الما

	No. of Bac.	per Cubic Centi- meter.	
20 0		Presenc Abs. of	+++++
Carro	ture tart.	Tempera of Air, C	
	r, C.	Tempera	00,04,000,000,000,000,000
		Heigh otaW	<u> </u>
	TRO- GEN AS	Nitrates.	48488855855555555555555555555555555555
	NITRO	Nitrites.	2000 2000 2000 2000 2000 2000 2000 200
	IC EN.	-sus	312 32. 36. 496. 56. 88. 1.44. 68. 88. 1.312 1.3
	ORGANIC NITROGEN	Dis- solved.	844. 98. 98. 98. 98. 98. 98. 98. 98
	ON	Total.	68888888888888888888888888888888888888
	92	e p,pud	1.55 - 1.00 - 1.
	EN A	Pioni Pioni	208 208 208 208 208 208 208 208 208 208
	NITROGEN AS AMMONIA.	L'10T	8.448888884484444444444444444444444444
	Z	Free Am- monia.	34 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
	N. ED.	By Suspd	463800004865477777 665777777788654777777
	OXYGEN	By Dis- solved.	awaaaaawaararr -ar amaa rear wer-
	000	Total.	9 5 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	'əu	Срјоц	######################################
	Color	on Igni- tion.	医抗抗抗抗抗抗抗抗抗抗抗抗抗抗
	RESIDUE ON EVAPORATION.	= p,pud -sns	4 6 6 9 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
		o sid b'vlos	0.60 % 10 % 10 % 10 % 10 % 10 % 10 % 10 %
		1.10L	4 6 8 8 8 6 6 6 4 4 4 4 4 6 8 4 8 6 8 6
		bended.	85.6 8 85.6 8 8 85.6 8 8 85.6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
		solved,	332.4 238.8 300.4 300.4 300.4 143.2 130.4 143.2 144.8 144.8 144.8
	RESI	T'otal.	388.8 381.6 386.8 308.8 308.8 308.8 313.6 313.6 314.8
		1	<u></u>
		robO	
	æ.	Color.	M. F.09 M. F.09 M. F.03 M. F.03 M. F.55 M. F.55 M. F.06 M. F.0
1	RANC	ర	
	APPEARANCE	Sedi.	Cons'd Wuch Cons'd V.M'ch Wuch Cons'd V.M'ch Much Cons'd
		Lucy,	105024008100F48000 *********
	OF	1900 1900 Collec-Examition.	
	DATE OF	ec- F	4-1-2 12 22 23 23 24 25 25 25 25 25 25 25 25 25 25 25 25 25
1		1900 Collection.	Jan. Feb. "" Mar. Apr.
1	No.	Serial	6617 6678 6688 6888 6888 6888 710 710 7118 718 718 718 718 718 718 718 718 71

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. C., Cloudy. T., Turbid. VM., Very Muddy. COLOR-M., Muddy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 77.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

Report of Arthur W. Palmer. T. J. Burrill. University of Illinois. SOURCE OF WATER-MISSISSIPPI RIVER, WEST OF MIDSTREAM, JEFFERSON BARRACKS. MO

57.200 19.900 97.200 35,500 125,000 216,000 135,000 000.101 46,000 61.000 per Cubic Centi-meter. No. of Bac. Presence or Abs. of Coll. Тетрекатиге оf Аіт, Септ. Nitrites.

| Nitrates. | Nitra 0114108 0114108 0114108 011 ∞855558,∞ NITRO-310 32 64 48 256 48 3 256 44 8 3 392 44 8 3 552 556 1.84 408 1.632 ORGANIC NITROGEN. Dis-64 2. 2 336 1.336 224 1.33 624 2.01 448 1.88 .568 3.84 Total. 048 p,pud -sus Albuminoid Am. NITROGEN AS AMMONIA. Totrl By Suspd Matter. OXYGEN CONSUMED. -sid ya solved. Total. Chlorine. Color Igni-tion. 372 1 316. 56.4 29.6 29.6 ... 382 1 297.2 115.2 31.2 28. 3.2 1 207.2 115.2 31.2 28. 3.2 1 20.8 310.2 30.2 31 p,pud -sug Loss on lg'n. RESIDUE ON EVAPORATION. Pages, d 1,10,7, Pis-Total. Odor. Much M. F.03
Much M. F.03
Much M. F.03
Consid M. F.03
Much VM.F.
Consid M. F.03
Much VM.F. 26848248 Color. APPEARANCE. p.suo.) * Much Sedi-ment. Cana 路路江西部 Collec-Examition. nation. x 表数数 to 50 数 : : : : : 3 Mar. DATE OF 1900 6826 Feb. 6962 " 7011 Mar. 7044 .. 1938 Apr. 1956 : Serial No.

COLOR ON IGNITION—G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. C., Cloudy. VM., Very Muddy. T., Turbid. Colon-M. Muddy. TURBIDITY-* Decided. § Very Decided. + Distinct. + Very Slight. | Slight.

TABLE 78

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

Report of Arthur W. Palmer, T. J. Burrill, University of Illinois. SOURCE OF WATER-MISSISSIPPI RIVER, 100 YARDS FROM MISSOURI SHORE, JEFFERSON BARRACKS, MO.

No. of Bac.	per	Cubic Centi- meter.	16,200	00f'6	51.600	19.300	21,000	69,000	84.000	33,000	134,000	320,000	106,000	59,000	55,000	84,000	38,000					
		Prese Abs.	+	+	+	+	+		+	+	+	۵.	+	+	+	+	+					
		qməT if A lo				:	:					:			:		:					
er.	Height of Water. Temperature of Water, C.				2.6																	
AS	Nitrates.		1		1				8f.													
NITRO- GEN	Nitrites.		810.																			
ic in.	· pa	sng -sng	.288								GTD.	₽ 1	$\overline{}$		-	_	-	:				
ORGANIC NITROGEN.		-siq	.552																			
ON	·I.	Tota	18.	22	.73	so.	.616	1.2	1.13	88.	·	e3.€	2.36	1.336	1.4	2.04	1.96	:				
50	Am.	p,pud -sns	.16	. 192	176	.208	960.	.33	.304	.064	1.568	<u>&</u>	.816	.352	304	.608	F9F.	:				
Nitrogen as Ammonia.	Albuminoid	b'vlos	176	.112	91.	. 128	14	. 192	. 224	808.	. 192	-24	808	£22.	.16	.16	144					
TROG	Albun	Tot'l	.336	304	.336	.336	₹;	.512	558	.272	.76	. 12	.034	.576	.461	7.68	.608					
Z	·u	Free A sinont	- T-	. 133	- .	104	44.	575.	: 575; 575;	80:	352	- 7	.33 E.	. 224	.144	.16	1 90.					
ED.	pq r.	By Sus	3.1	ව වෝ	4.3	 	1.5	6:	6.9	3. 8.	22.4	31.6	.6.	, o	9.7	11.5	11.9					
OXYGEN		By Dis			Ţ-	4		īĞ.	ပ်	v.	ŗ-	<u>.</u>	6.		6.	5	٠.					
000	I.	Total	1.118.	ा । ।	211.4	6 11.9	7.9	4 13.8	6 13.5	4 9.5	29.4	4 30°	e: 25 	.91		2 17.1						
	Chlorin		23.	8	: ::	17.	19.	<u></u>		-	6.	{~	<u>-</u>	<u>.</u>	=	<u>.</u>	<u>-</u>	:				
Colo	on	tion.	Ö.	<u>:</u>	?	?	~	~	ä	≃:	≃:	<u> </u>	<u>:</u>	~	≃:	~	===					
).X.		p,pud sng	_ :	v.		16.	0	; - 	33.	4.88 8.89	.8 39.6	46.8	33	19.2	33	18.	18.4					
RATIC	Loss on 1g	p'vloz	21.2	3.0° 	15.6	17.6	17.5	14.		180	22	+	133	12.4	9.6	20.1	ડ્ડ સ					
RESIDUE ON EVAPORATION	To	1.10.L	8 21.2	2 26.	5 20.	÷ 34.	4 28.	831.2	50.	2 41.2	6.852.4	6 60.8	34.	8.8 3I.6	8 31.6	4 38 4	6.38.4	-				
ON E		bende -sus		131.	252.5 20.			356.8 31.	166	19	ರಾ	6	7.6	86	8	8	1387.					
IDUE		-si(I solve	338.8	292.	324.	.2272.8	.2312.8	2246.4	216.	321.2	149.6	141.6	162.8	178.	278.8	240.8	9. T. 2					
RES	1	Total	377.6	423.2	576.5324	619.2	411.2	603.2	683	514.4321	2746.4 149.6	105.2	338.8	. 764.8 178. 5	079.6	337.2	614.8					
	10t	00			:	:		:	:		· · · · · · · · · · · · · · · · · · ·		• • • • • • • • • • • • • • • • • • • •				:					
61		or.	F.03	F.03	F.03	F.04	F.03	F.02	- -	F.03	F.3	F.I	F.25	五.05	F.30	F.05	F.03					
RANCI		Color.	M.	M.	M.	M.	N.	M.	1. N.	M.	M.	VM.		M.	M.	M.	M.					
APPEARANCE		Sedi- ment.	Cons'd	7,7	Much	,.	Cons'd	Much	29	Cons'd	V.M'ch	.,	Much	77	**	,,,	;					
	Ī	Turb'	*	12 *	19 *	* 98	÷.0	* 01	33 *	ر *	co co	1-	**	30 *	* 9	*	% 	:				
0 P	1000	Exami nation	Jan.	9,9	7.9	9.9	Feb.	,,,	.,	Mar.	7.9	7,9	.,	"	Apr.	:4	"					
DATEOF	1000	ė,	Jan. 4	11 ,,	18	ફુક	Feb. 3	6 ,,	:	Mar. 3	00	,, 16	::	66 .,	Apr. 5	2	06 .,					
.oV	al	Seri	6619 J	6671	6710				6963		70H5	7102	7142	7188			7376	:				

Color on Ignition-G., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. C., Cloudy. T., Turbid. VM., Very Muddy. COLOR-M., Muddy. TURBIDITY-* Decided. § Very Decided. † Distinct. ‡ Very Slight. | Slight.

TABLE 79 and 80.

STREAMS EXAMINATION CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS PARTS PER MILLION.

SOURCE OF WATER-ST. LOUIS, MISSOURI, TAP WATER. (Regulars and Extras.)

Jo	
Report	

Comme	6)	1.150 1.
No. of Bac.	Cubic Centi- meter	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1
	resend to sqv	
	Temper of Air,	
eathre	roquoT isW lo	
	lgioH - iaW	$x_{0}x_{0}x_{0}x_{0}x_{0}x_{0}x_{0}x_{0}$
NITRO- GENAS	Sitratics	200 200 200 200 200 200 200 200 200 200
7.	Sitrites	
NIC GEN.	-sns	
ORGANIC NITROGEN.	-si(I ,b9vfos	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
	Total.	
AS . 3	p.pud -sus	
TTROGEN	Torr Dis- Dis- Dis-	5.55.55. 5. 4.55.4. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.
Nrreg AMS	Torr	255252355245454545454545455555555555555
	Free Am	80.000
EN MED.	By Suspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ONYGEN CONSUMED.	By Dis.	<u>ထိုကို ကို မိုကို မိုကို လို လို လို လို လို လို လို လို လို လ</u>
	Total.	$ \delta u u u u \omega \omega \omega \omega \omega \omega $
-	TO[(1')	
Color	Igni- tioni.	$0 = \frac{1}{2} + $
	E p.pud	က် ကြိမ်းက ေ မြိတိန္တိုက္ပြည်သိတ္ကမိုးရိန်မိုက်သည် မြတ္လည္က ၁၈ မြန္မာကိုလိုသည်။ နေမာက်တွေ ၈ တွေ တရာကက မွာ နေမာက္သမွာ မွာလနေကာက္သည်။ မြန္မာကိုလိုသည်။
ORAT	Figure 1 10 T 1	\$64686666666666666666666666666666666666
Evar		90000000000000000000000000000000000000
RESIDUE ON EVAPORATION	sus Sus	88122222256688882775286895576667262766776
Sibt's	-sid Joylos	337 6 28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
23	Тота	28.88.88.88.88.88.88.88.88.88.88.88.88.8
1.*	obO	
	or.	තුන්තතත්වන් වන්නේ සම්බන්ධ වන්නේ වන්නේ සිසු අතු අතු සිසු සිසු සිසු සිසු සිසු සිසු සිසු සි
ANCE	Color.	ZEZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZZ
APPEARANCE.	Sedi: ment.	Little Cons'd Much Cons'd Cons'd Much Cons'd Much Cons'd Much Cons'd Con
1.	· v,qang,	* * + * + + * * * + + * * * * * * * * *
Gu	1900 Exami- nat on.	H
DATEOF	c. EN	
DA	1900 Tollection.	Mar
So.	Intros	を変数を表現を表現を表現を表現を表現を表現を表現を表現を表現を表現を表現を表現を表現を

Color on lengton -C., Gray. B., Brown. DB., Dark Brown. LB., Light Brown. RB., Reddish Brown. BG., Brownish Gray. Bh., Brownish. R., Red. Rh., Reddish. W., White. T., Turbid. C., Cloudy. Coloi: M., Muddy. VM., Very Muddy. Trubbuty -* Decided, & Very Decided, + Distinct. + Very Slight. Slight.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Edwin O. Jordan,

f Chicago.	No. of Bac.	per	Centi- Centi- meter.		• • •	1 910 000	995 000	145 000	1 415 000	1 020 000	1,010,000,1	015 (20)		950.000	000 070	1.190,000	1.370.000	600 000			1 360 000	1,700,000	1 850 000	9 650 000	1 680 000	1,000,000	630,000	970,000	3,190.000
University of			Prese Abs. o			•																		:	•				
Jniver	ature, C.	:19 :11	qm9T A 10			12	0.00	200	5	57	 	5	01	19.	5.	10	10	<u>∞</u>	9	00	10	=	-		000	7	+	60	ο
	ature D'. T	919 91£	qmaT 377 lo			: 2	. <u>-</u>	9	· {~	23	10.	19.	19	2	6.	13.	13.	10.5	00	2	0	6	10	10	10.0	000	6.	.6	6.5
	10 1	epr pyr tyr	giəH SW			15	100	0.7	1	- 1C	4.7	50.00	4.5		ın	5.9	-	5.6	5.6	20	70.	5.6	2.0	10	310	20.	i.c	٠ <u>٠</u>	÷
	NITRO- GEN AS	80	Nitrati			•		_							_						-						0.		
		!	Nitrit		:	: 0	2		000	0.	0.	0.	0:	0.	0.	0.	0.	0.	0.	0.	8	0.	0	0	0	0	0	0.	0.
	ORGANIC NITROGEN.	_	-sus		:	: .							:	:	:	:	:	<u>, :</u>	:	-								:	
ILL.)RGA VITRO	-	sid solve		:	:				:		:	:	:	:	:	•	:	:									:	:
		"T	b'bnq B10'T		:	:			80		9			-		ः ः	-	-		3	-	9			9			-	9
BRIDGEPORT,	25 4	oid Am	sus	1.6	3.6		ু কয়		^)		96:	_	_				_		-	_	_	-	_	্	_		1.9	_	<u></u>
	ITROGEN AS	Albuminoid	-sid b'vios	<u> </u>	o; o				.59.	-	£8.	:0	_		_	-		-			_	-		_	_			_	
CANAL,	NITR AM	Alk	Tot'l	ಣ	. o	3 ec	(0) (0)		1.4	1.8	e;	- T	o)	3.5	2.6	2.6	1.9	1.5	₽. 51 1. 4	30	1.9	e5		65	20.00	13.	(0) (00)	: · · ·	5.
GAN C		Freekm- monia.		35	16.																								
Міснівам	EN IED.	pd L	BySus	13.7	36	0 - 1	15.8	13.2	6.4	6	19.4	14.3	20.3	3.5	10.5	13.3	17.5	10.3	15.	16.3	16.6	13.4	20.4	25.4	19	3.5	9.	14.4	16.8
AND A	OXYGEN		By Dis	28.3	20.00	0 0 0 0 0 0 0	=	15.	12.6	8.61	15.	13.5	15.4	16.2	19.3	35.	8.6	6	- 1-1	_	12.8	9.4	12.8	5.	10.2	23	7.	12.4	17.
	့	1.	E10'I'		501.3																								
LLING	·ət:	ıi1	Срдо		130																	+	126.	156.	106	103	135.	118.	166.
ren—]	ON.	pe	puod Sus		113.6	351	104.8	38.4	182.4	46.	74.	-13	33	7.4.	81	100	54.	55	(3	40.	61.	34.	90.	107	98	18	.69	38.	.89
SOURCE OF WATER-ILLINOIS	RESIDUE ON EVAPORATION		Solve		00																	374.	472.	576.	433	454.	463.	166.	534.
TRCE O	RES EVAI	.1	rioT'	582.	7.04.4	165.	492.	544.	650.4	536.	442.	416.	468.	512.	502.	710.	504.	377.	567.	420.	481.	408.	562.	683.	530.	459.	534.	504.	602.
Sot		Odor		Gassy		:	:	4.	,,	,,	9.	*	;	99		,,	,,	,,,	,,	1.0 1.0	9 .	9.7	9 9	9 9	**	9.7	9.1	9,9	9.7
		_	.:	<u> </u>	:			_	_				>					· V		_	_		[A]		_		-		
			Color		:	• •	9.	9.	ıc	ıc	***	ಬ	Muddy	9.9	7 9	*	7.	Mudd	ಣ	-41	77.	₹.	Muddy	9.9	*	9.9	9.9	9.9	4.
	APPEARANCE		Sedi- ment.	V. Little	V. Much	Little	Cons'd	Little	Cons'd	:		p,suo,)	94	Much	:	d d	Little	Cons'd	Little	Much	9.9	Little	9 9	,	Much	Little	Cons'd	4.4	:
	AP		Turb'y.	Much	: :	Decided	Much	Decided	9.7	Much		**************************************	• • •	2 .	9 9	:	6.9	9 4	-9 -0	9.9	99	"	9 9	- 18	9.	9 1	79	*	9 9
)F	1899	Exami- nation.	ay 27	ine 11	65		01 ,	- 5	· 32					6:1				, 17					21	88	3c. 6		33	37
	DATE OF	_	c- E;	27 May	11 June 1	٠ 2 %	3 July	, 01		16	31	6 Aug.	. 58	1/2	18		2 Oct			23	30 08	6 Nov	13 "		25	4 Dec		300	32
	DA	1899	Collec- tion.	May 2		91	Inly	:	**	19		Aug.	• `	:	*		Oct.	,,,	*	2 @5		Nov.	;	99	33	Dec.			3.
	.o.X	[8]	Seri	6	76.7	88	23.1	569	308	343		416		8 909	633	_	_	759	791	830	_		216	086	6001	_		1107	1141

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LOCKPORT, ILL.

7.
4
-
2
السا
-
\circ
VI.V
1.
print
part.
.5
(E)
_
٠
0
-
- Secret
3-4
0
Š
5
05
~
2

University of Chicago.

No. of Bac.	per Cubie Centi- meter.	1.115.000 1.055.000 1.330.000 1.330.000 1.330.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.85.000 1.310.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000 1.340.000	1,000,000
	Presence lo selv		
time C.	Tempers,	45588888888888888888888888888888888888	
r, C.	sroqmoT otsW lo	왕 말 왕 말 왕 말 왕 말 왕 말 왕 등 영 등 영 등 장 등 작 등 명 등 등 중 구 등 등 중 구 등 등 등 등 등 등 등 등 등 등 등 등	9
	Height State		
RO- N AS	Nitrates	- 19 + 0 + 0 + 0 + 0 + 19 0 + 18 0 = 0 + 19 0 + 18 0 = 0 + 18 0 +	. O., A
NITRO-GEN A	Nitrites		
GEN.	pepuad -sns		
ORGANIC NITROGEN.	Dis-		
ON	Total.		-
	a sus	65.2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
TROGEN AS	Pionimoid Pionim		
NITROGEN	Torr	60 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1
	FreeAm- monia.	8 1 <td></td>	
EN.	BySuspd Matter.	కాంట్రాశానా చాయబుగునుగు : చాగా బ్యబుబుగా : నుబుబుబుడా కాకే కాంట్రా శారుగు కారాలు అందారు : చారు అందారాలు ప్రశాస్త్రము చారు చారు చారు చారు చారు చారు చారు చార	
OXYGEN	By Dis-	<u> </u>	•
33	Total.	1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0	
16,	Chlorin	<u></u>	
ON TON.	pəpuəd -suş	S = 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
RESIDUE ON EVAPORATION.	Dis-	8. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	
RE	Total.	28.50	
	Odor	Gassy	
	Color.	Muddy	-
APPEARANCE.	Sedi- ment.	Much Cons'd Cons'd Cons'd Cons'd Much Cons'd Much Cons'd Little Little Much Cons'd Little	-
AP	Turb'y.	Much Decided Much Decided Much Decided Much Much	
40	1899 Exami- nation.	Nay	
DATE OF	9 ec. E		5
D D	1899 Collec- tion.	May. July Sept. Se	
.02	Serial	845888888888777788888888888888888888888	727

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, LOCKPORT, ILL.

Z	
73	
4	
0	
ORD	
\simeq	
0	
~	
EDWIN	
2	
~	
-	
-7	
-	
0	
\cap	
~	
2.	
por	
0	
2	
43	
rel	
-	

EDWIN O. JORDAN,	University of Chicago.
t 0 f	
Report	

No. of Isac.	per Cubic Centi- meter.	6.900 6.9000 6.900 6.900 6.900 6.900 6.900 6.900 6.900 6.900 6.9000 6.9
Seor Coll.	Presence Abs. of	
	Tempers,	4.8.8.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.
	Tempers of Wate	848448888 48282755 6004000000000000000000000000000000000
	Helghl etaVI	86.00 p. 4 8
RO-	Nitrates	0.000000000000000000000000000000000000
NITRO- GEN A	Nitrites	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0
IC TEN.	-sug	
ORGANIC NITROGEN.	Dis- solved.	
QN.	Total.	
	Sus- Sus-	2000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
TROGEN AS	Pioniminos -sid b'vlos	844445555844884444444888 854445558848884544444888 855588488888888
NITROGEN	T'ot'l	### ### ### ### #### #### ############
	FreeAm- monia.	1.05 1.05
ED.	BySuspd	
OXYGEN	By Dis- solved	$\frac{\omega \overline{\alpha} \overline{4} \overline{4} \overline{\omega} \overline{v} \overline{\alpha} \overline{\alpha} \overline{v} \overline{v} + \underline{\alpha} \overline{v} \overline{v} \overline{c} \overline{\alpha} \overline{\sigma} \overline{v} \overline{v} \overline{v} \overline{v} \overline{\alpha} \overline{w} \overline{v} \overline{v} \overline{v} \overline{v} \overline{v} \overline{v} \overline{v} v$
Co	Total.	$\frac{\pi F \tilde{\omega} \tilde{w} \tilde{\omega} \tilde{w} \tilde{\omega} \tilde{w} \tilde{w} \tilde{w} \tilde{\omega} \tilde{w} \tilde{\omega} \tilde{\omega} \tilde{\omega} \tilde{\omega} \tilde{\omega} \tilde{\omega} \tilde{\omega} \omega$
.ən	Chlorin	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ON ION.	pepued ens-	© 14 © 29 - 12 0 0 4 4 0 ∞ 0 1
RESIDUE ON EVAPORATION	Dis.	23.2. 23
RE	Total.	460
	Odor	None None None
	Color.	တွင်းတွင်တွင်းတွင်း ကြောက်တွင်း ကြောင်းတွင်းတွင်းတွင်းတွင်းတွင်းတွင်းတွင်းတွ
APPEARANCE.	Scdi- ment.	V. Little Little Little Little Cons'd Little V. Little
AP	Turb'y.	Slight " Distinct None Distinct Distinct Distinct Decided Slight None Slight None Slight Decided Slight None Slight None Slight Nuch Slight Decided Slight Nuch Slight Much Slight
OF	1899 Examination.	May 9 June 6 13 13 13 14 15 16 17 18 19 19 19 19 19 10 10 10 10 10 10 10 10 10 10
DATE OF	1899 Collection.	May 57 7 10 10 10 10 10 10 10 10 10 10 10 10 10
No.	Serial	11100 889 987 887 887 887 887 887 887 887 887

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, NORTH OF JACKSON ST., JOLIET, ILL.

EDWIN O. JORDAN,	University of Chicago.
JO	
Report	

No. of Bac.	Cubic Centi- meter.	339, 600 230, 600 230, 600 250, 600 115, 600 60, 600 60, 600 110, 600
	Present To .sd/.	
ernter e, C.	Temper	
rature er, C.	Temper	
10 11	dgioH dsW	
NITRO- GEN AS	Sitrates	
	pended Sitrites	
NNIC	solved.	
ORGANIC NITROGEN.	Total.	
	p,pud sng	80 88 4 6 8 4 8 6 5 4 8 4 8 6 6 7 8 7 8 8 6 6 8 8 8 8 8 8 8 8 8 8
20 .	Albuminoid b'vlos	· · · · · · · · · · · · · · · · · · ·
NITHOGEN A	I'10'l'	- 20 20 - 20 20 20 20 - 20 - 20 - 20 20 20 - 20 20 20 20 20 20 20 20 20 20 20 20 20
1	FreeAm- monia.	<u> </u>
ED.	Bysuspd	8 4 8 8 8 6 8 8 7 8 4 8 8 7 8 9 6 9 6 9 8 8 7 8 9 8 8 7 8 9 8 8 7 8 9 8 8 7 8 9 8 8 7 8 9 8 8 9 8 8 9 8 9
OXYGEN CONSUMED.	By Dis-	6 2 2 1 1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2
25	Total.	<u> </u>
.əu	Срјоци	6.8 6.4 8 5.8 5.7 5.8 8 8 8 8 8 9 7 7 7 7 7 7 7 8 8 8 8 8 7 7 7 7
ON TON.	papuad -sug	\$68184168168888888846888548888888888
RESIDUE ON EVAPORATION	-sid beytos	8 8 8 8 8 8 6 6 8 8 8 8 8 8 8 8 8 8 8 8
RE	Total.	9.99.89.89.89.89.99.99.99.99.99.89.89.89
	Odor	QB.SS
	Color.	Ν
APPEARANCE.	Sedi- ment.	Much Little Cons'd Cons'd Little Little Little Little Little Much Much Much Cons'd Much
1.	Turb'y.	Decided Distinct Decided Nuch Secided Nuch Nu
0.0	1890 Exami- nation.	May June : : : : : : : : : : : : : : : : : : :
DATE OF	1899 Collec- E tion. na	
Zo	faires o.s	1

TABLE 85.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, SOUTH OF TOWN, JOLIET, ILL.

-
Z.
-<
RD.
-
0
7
\circ
die
=
WIN
~
痉
parties.
Squared.
0
_
-
)
0
00
0

University of Chicago.

1									_		_							_		
No. of Bac.	Cubic Centi- meter.	11	100.4.ca	610.00X	850,000	760.000	530.000	180.000	1.760.000	120.000	830,000	170,000	355,000	390,000		2,515,000	370.000	2,540,000	280,000	120.000
	Presen	:	:	•				:	:						:					:
	Temper if Aio	•						36.	24.	6:1	19.	133	<u>x</u>	8.	17.	<u>x</u>	<u>x</u>	-3	<u></u>	
	Temper of Wate		:	:			19.	21.	15 13 13		£5	32	; ;	: :9:			£	?}	17	-8.
	Heigh Wat	:	:	:	:	:	:	:	:	:	:	:		:		:				
ro- r As	Nitrates	0.0	<u>-</u>	0.	0.	??	7.	. 15	9.	φ.	0.	0.	· 왕	.05	0.	.035	0.	.005	7.	0.
NITRO- GEN A	Nitrites	0.	310.	610.	1.00	30.	.036	0.		0.	.044	0.	0.	0.	0.	.003	0.	0.	0.	0.
ORGANIC NITROGEN.	pəpuəd -sng	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
RGAL	Dis- solved.		:	:	4	:	:	:	:	:	:	:	:	:	:	:		:	:	:
OZ	Total.	<u>:</u>	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
	E Papid	-:	o. T	જાં	966	1.4	1:1	1.9		1.94	8 .	09.	86.	1.46	1:1	1.0g	:	1.26	۲-	66.
NITROGEN AS	Albuminoid Silvios	e;		_	.704	_	•		1:4								•		86.	
NITRO MMA.	T'10T	33 G	0.23	65	1.7	∞ ∞	3.12	2.96	4.8	96.	1.52	1 84	3.08	5). To	3.5	2.04	2.12	1.6	1.68	1.8
	Freedm- monia.	7.75	9.	∞.	6.3	14.	16.	13.	16.	13.	4.68	13.3	13.6		14.8	11.6	14.6	16.2	16.2	18.4
ED.	Bysuspd	6.5	10.4	16.3	13.7	∞ 35	30	€. €.	13.	12.7	9.	6.8	14.8	9.91	11.4	16.3	9.3	00.	٠ <u>ت</u>	11.1
YGE	By Dis-	15.7	J.	??	10	30	∞	9	T	∞	03	00	_	_		33	7	9		00
OXYGEN CONSUMED.	Total.	20.2		10.	??	10	-	∞	4	10	∞	9.	∞	9	7	10	9	-		C:
.əu	Chlorh	60.2	65	74	-84	94.	112.	87.	113.	100	825.	90.	102.	134.75	111.	115.5	117.	128.	113.	109.
ON.	pepued -sus	106.	ž Ž	3300	306.4	55	54.8	152.	5.4.	64.8	158.4	114.	104.	150.	.02	160.	158.	57.	40.	40.
RESIDUE ON EVAPORATION	Dis- solved.	454.																		
RENE	Total.	560.	500.	712.8	733. A	584	622.8	504.	509.2	597.6	565.6	617.	586.	692.	546.	662.	.099	583	478.	484.
	Odor	Gassy	:	9.5	9.9	9,9	9.9	9.9	9,9	9.9	9,9	3 9	9.9	9,9	9.9	9.9	9 4	7.9	9.9	9.9
	Color.		:		71	∞.	Muddy	· 	က့	9.	10.	ıc	ः		4.	-:	c)	c;	છુ	₹.
APPEARANCE.	Sedi- ment.	Much	,	Little	Cons'd	9.9	w •	*	Little	**	9	**	Cons'd	9,9	Little	Cons'd	99	9,9	,,,	Little
AP	Turb'y.		Distinct		Decided		*		Decided	:		Much	:	* 9	_	Much		*,	:	9 *
OF	1899 Exami- nation.	May 4		667	June 5	÷		96 ,,		01 .,		16 31		Aug. 7	Ŧ1 ,,	:	966		_	30
DATE OF	c- E	T T	333	<u>ج</u>	5 Ju	<u>@</u>	19	96	3/11/8	10	~	634	31	7. 7		21	000	4 Sept	11	100
DA	1899 Collection.	May	:	9.9	June	9,9	:	:	July	:	9 4	:	,.	Aug.	,,,	9.9	9.9	Sept.		9.9
No.	Serial No.		95	67	97	127	160	202		268	305	340	380	_	_	495	525			638

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER-KANKAKEE RIVER, WILMINGTON, ILL.

	0
	hicago
ź	(7)
DA) JO
OR	ty.
	ersit
j	ver
7.	ni
=	
1	
5	
) I (
<u> </u>	
lavi	

No. of Bac.	per Cubic	Centi- meter.	7.00	805 ×	7.650	008	006.6	0001	000	9.400	7 050	1,550	20:1	2 650	009	950	(K) (1) (1) (1) (1)	12,200	15,000	100	1,200	200	005,1	900,	4,560	201.2	0.30	#,·100		2,000	19 500	25,700
	lo	'sqv															•		:			:	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	•		•		0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0			
11111.6 . C.	rere Tiv	Temi To	İ	5	96	50	66	= x	00	26.	16.	<u>x</u>	12	6	20.	17.	16.	16	100	10.	2-	10	<u>x</u>	1		· Ł-	- 10	. <	. 7		; c	15.
		Temil W 10			54	3	95	33	3	27.	0.00	6	99	- 22	6	6	10	. Co))	. 62	. 6	=	0		0 10	÷α	i i c	. 7	9	0	. i.c
10 ;	ots:	IoII W	65	10	10		833	-	1.6	. T	6.3				-			-		000	2				. 6	210	. ox	; us	000	000	C	30
NITRO- GEN AS	sə1	Nitra	! -	2.5	_	_			0.5	_	_	0.		0.50				_	100		_		_	_		-		· G	-		50	3.25
Z	J	Mitri	035	800	0.08	000	800		90.	900.	0.	00.	900	8	00.	00.	100	000	0.	00.	0.	0.	0	0.	000	0.08	3	3	03	. 012	.012	.0021
Organic Nitrogen.		solve sus busq									:																					
ORG. NITR		BIOT.								•	:	•																				
	Am	p,pud ·sng	420	0	.024	80	300	979	800.	.040	.136	800.	.176	0.88	.136	.072	80.	21	0.14		0.88	960	960	050.	003	016	061	0	800.	190.	0.	0.0
HEN AS	Albuminoid	Dis.	.336	.64	9.	525	.31	888	424	.368	.280	++.	979	288	.280	855	.948	24	25		91.	.136	. 168	128	.178	.2	192	218	956	272	181	155
NITROGEN AS	Albu	l'toT	98.	.64	.624	9.	312	.56	. 133	.408	.416	.448	84.	.376	.416	.352	328	.36	256		218	533	1964	.184	.186	.216	926	2.0	196.	.3386	184	.28
		Freek		.085	.082	20.	.089	.074	.112	.064	.052	.0:	80:	890:	.076	80:	10:	.058	810.	100.	10.	. C##	910.	:03	<u>eto:</u>	.058	100	.03	.018	ţiO.	100.	100.
SN IED.	pds er.	BySu	_c;	00	3.	3.4	e5	35 50	33	65 63	3.4	7	رى ب	33	1.6	90		Ţ	G)	0.		1-	σ.	φ.	ж.	X.	-	10	0.		33	7
OXYGEN CONSUMED.		By Di	14.2	17	16.4	311.4	50.00	5.1	က တ	G? →	00.00	30	<u>ي</u>	10	6.8	7.4	9.9	6.6	9.9	6.5	53.50	4 33	9.4.6	÷	55	7.4	×	3.7.8	7.5	4.4	φ. ×	į
ပစ္ပ ——	.Iı	310'l'		<u>}-</u>		.11.8	- 2	×.	3		00	0	00	00	∞ 4.	7.0	2	00	6.8	9	8	ω.	10	4.6	6.	00	00	00	5-	4.8	œ	{-
.91	tiro	СРГ	-	1.65					ب ب ب				33.7	4.1	3.7	33.0	4 6	3.0	4.9	30.00	₩ ∞	85 00	4.6	∞. ∞.	4.3	3.6	2.6	e:	3.6	3.3	e: ∞:	3.5
ON MON.	 pə	enS ottoq		233	-		85		55.6	<u>x</u>	51.	57.	100	59.	38	30.	E	33	64.	4	33.	. 22	21.	12.	23	36.	4.	4	15.	0	11.	16.
RESIDUE ON EVAPORATION.	-8	ovios	267.6				260.8			366.	258.	275.	243.	251.	219.	250.	231.	2330	252.	258.	251	254.	256.	248.	275.	990.	319.	294.	291.	530.	301.	274.
Eva	l.	sto'I'	309.6	343.8	300.	332.	290.		359.6	341.	300	335.	342.	310.	\$ 5	300	295	314.	316.	262	276.	985	.280	360	277.	326.	316.	296.	300	330.	345.	290.
	Oaor		None	9.9	9 9	3.9	10	9.9	79	9.9	9 9	9.9	9 9	,,,	9 9	9 9	9.9	9 9	9.9	9.9	• •	:	4	9 9	7 9	* 9	9 9	9 9	99	9 9	99	9.9
		Color.	10	33	တ္	7	171	चर्चा	60	Muddy	e;	?}	_	0.	0.	0.	9.		25	ಲ್	93	કરં	टाः	ণ!	: :	ಣ	ಣ	<u>.</u>	ıc	٠٠;	ાર 	.37
APPEARANCE.	Sodi	ment.	Little	:	9 9	9 7	9 9	9 9	9	9 9	-0 -0	*9 *\$	10	3 9	Cons'd	3 3	Little	9 9	99	V. Little	-0.	Little	9.9	V. Little	9.9	Little	9.9	V. Little	4	Little	9.9	93
IV.	1	Turb'y.	Distinct	None	Distinct	Decided	None	Distinct	Derided	None	Decided	Much	Decided	Much	Decided	Much	49	9 9	Decided		9	12 15	99	Slight	9.	Distinct	Much	Decided	Slight	Decided	Much	Slight
OF	1899	Exami- nation.	5 June 6	::	5:1 .,	36	fuly 3	10	77	च्या कर		lug.		<u>-</u>	0.5	Sept. 5		61 ,,	96 ,,	Set. 2	** 10	** 13	53		iov. 6	** 13	30	3)ec. 4	***	61 ,,	25.
DATE OF		tion.	June 5 J	21	19	96 ,,	11y 3.1	01		63	23	13			3		,	30	66 33	st. 20	J	9	33	300	19 :	· 13	07 ,,	. 27)ec. 4 D	11	X.	. 26
,07.	[8]		100 1	4	161	10 E	239 .11	971	30.00	341						ESS NeE	. 200	. 019		714 Oct		162		T S	P 4			1010	p	100	11112	1142

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO

SOURCE OF WATER-ILLINOIS RIVER. MORRIS, ILL.

Report of EDWIN O. JORDAN,

University of Chicago.

65.000 70.000 140.000 140.000 345.000 351.000 434.000 466.000 568.000 372.000 175.000 175.000 175.000 270.000 100.000 30,000 755.000 130.000 782,000 115,000 720,000 250,000 640,000 No. of
Bac.
Per
Cubic
Centimeter. Presence or Abs. of Coll. Temperature of Air, C. -45566 - 665 - 655 Temperature of Water, C. Helght of Water. GEN AS Nitrates NITRO-0.000 Nitrites eng-sns ORGANIC NITROGEN. solved. -siq Total. p,pπd -sng Albuminoid Am NITEOGEN AS AMMONIA. $\mathbb{E}_{F_{0}}^{F_{0}} \otimes L^{F_{1}} = \mathbb{E}_{F_{0}}^{F_{0}} \otimes \mathbb{E}_{F_{0}}^{F_{0}} = \mathbb{E}_{F_{0}}^{F_{0}} \otimes \mathbb{E}_{F_$ [,10,] 42. 17.7 13.4 2.3 3.1 16.5 13.5 4 2.3 3.1 16.5 13.5 ONYGEN CONSUMED. By Dis-Total. Chlorine. క్రిప్రాల కోటా కోస్ 4 0 కోర్లు 4 అక్కించ్రులు అంది. - 4 4 6 4 - కుప pəpuəd -sns RESIDUE ON EVAPORATION Dis-Total. None Gassy None Gassy None Gassy Odor Color. ડાં માં માં જે દાં જે મર્ગ છે. માં હો હો માં નાં છે છે. મું હો મું છે. મું છે મું છ APPEARANCE Little V. Little Little Little Little V. Little Sedi-ment. Distinct Slight None Distinct Decided Slight Decided Slight Distinct Slight Distinct Decided None Slight Much Decided Turb'y. Much Slight 1899 Exami-nation. OF Sept NOV 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 25 ... 26 DATE 1899 Collec-May Sune July
Aug.
Sept.
Coct. Dec ; ; ; Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-FOX RIVER, OTTAWA, ILL.

University of Chicago.

Report of EDWIN O. JORDAN,

No. of Bac.	Cubic Centi- meter.	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1
	Present lo .sd A	
rature r, U.	Temper il A To	តាត់តាំនាន់នាន់នាន់នាន់នាន់ក្រាត់ តាន់តាន់តាន់ គឺដែ <u>ជដែលដាក់</u> ជ
	isqmsT isW lo	
	Helgh JaW	
0- AS	Mitrates	မေးက - မျှင်ဝင်နိုင်တို့ မှင်ဝမ်း ခုနှင့် ချွန်းမှု မေးဝင် မျှင်နှာ့ မေးက နေးမှုံးမှု
NITRO- GEN A	Solivites	2
	bended Sus-	
ORGANIC NITROGEN	-si(l solved.	
ON.	Total.	
	E Sus	8. 19. 20. 20. 20. 20. 20. 20. 20. 20. 20. 20
AMONIA.	Plourinoid	888888888888888888888888888888888888888
NITROGEN AS AMMONIA.	Torr	**************************************
1 3	Freedm- monia.	######################################
ED.	Bysuspd	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN CONSUMED.	By Dis- boylog	$\cos \infty \alpha \sigma + + \cos \sigma + \sin \sigma \cos \sigma + \alpha \sigma \sigma + + + + + \cos \sigma \cos \sigma \sigma \sigma \sigma \sigma + \sigma \sigma \sigma \sigma$
00	Total.	$ \begin{array}{c} \frac{\infty}{\infty} \cos \theta \cdot \phi \circ \phi + \frac{1}{2} \cos \theta \cdot \phi \cdot \phi \cdot \phi \circ \phi \circ \phi \cdot \phi \circ \phi \circ \phi \circ \phi \circ \phi$
ne,	Chlorin	$= \frac{1}{2} $
ON ION.	papuad sus-	
RESIDUE ON EVAPORATION	-siq solved,	8
REYA	Total	4 8 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	Odor	None contraction of the contract
	Color.	
APPEARANCE.	Sedi- ment.	Much Little "" "" "" "" "" "" "" "" ""
IV.	Turb'y.	Decided None Distinct Sight None Silght
OF	1890 Exami- nation.	Section Sect
DATE OF		8 1 2 2 2 2 3 5 8 1 5
D	1899 Collec- tion.	May June June June June June June June June
Zoz	Serial	######################################

TABLE 89.

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-ILLINOIS RIVER, OTTAWA, ILL

University of Chicago.

Report of EDWIN O. JORDAN.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-BIG VERMILION RIVER, LA SALLE, ILL.

University of Chicago.

Report of EDWIN O. JORDAN,

o. of Bac.	per Cubic Centi- meter.	1,000 1,00
.HoU	Presend Abs. of	
	Temperation To	
Temperature of Water, C.		
	idgiəH ətsW	
RO-NAS	Nitrates	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
NITRO- GEN A	Sofitin	0.000 0.000
NIC OGEN.	solved.	
ORGANIC NITROGEN.	Total.	
	m p,pud	600 600 600 600 600 600 600 600 600 600
N AS		:8558 :0549 :054
NITROGEN A	Albuminological filters of the state of the	5888888888888888888888888888888888888
Z	FreeAm- monia.	0.00
SD.	Bysuspd Matter.	
OXYGEN CONSUMED.	By Dis-	$\frac{\omega_1 \omega_1 \omega_2 + \omega_1 - \omega_1 \omega_1 - \omega_1 + \omega_2 \omega_2 + \omega_2 \omega_1 \omega_2 \omega_2 + \omega_2 \omega_2}{1000000000000000000000000000000000000$
03	.Into'l'	$4 \cdot 19 \cdot 6 \cdot 29 \cdot 19 \cdot 29 \cdot 4 \cdot 4 \cdot 4 \cdot 19 \cdot 19 \cdot 19 \cdot 19 \cdot 19 \cdot$
,9ti	Chlorin	中国もちゃらいとは本意のののはに対してのは、日本のは、日本のは、日本のは、日本のは、日本のは、日本のは、日本のは、日本
ON TON.	Dended Sus-	8778200086000000000000000000000000000000
RESIDUR ON EVAPORATION	Dis-	25
REEVA	Total.	466.1 46
	Odor	None
	Color.	
APPEARANCE	Sedi- ment.	V. Littile Littile Cons.d Littile Cons.d Littile V. Littile Littile V. Littile Littile V. Littile Littile V. Littile Littile Littile V. Littile V. Littile V. Littile V. Littile Littile V. Litti
AI	Turb'y.	Slight Decided Distinct Distinct Sone Slight Decided Slight Decided Distinct Much Slight Slight
OF	1899 Exami- nation.	May 93 June 13
DATE OF	1899 Collec- Extion.	Land to the state of the state
74(3)	-	104 June 1146
No.	Serial	44724724848557855785578557855785

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LA SALLE, ILL.

AN,
ORD
j
0
DWIN
田
of
cport
E E

University of Chicago.

No. of Bac.	per Cubic Centi- meter.	2000 2000 2000 2000 2000 2000 2000 200
	Presen lo .sd A	
	TeqmeT TiA to	
	TeqmeT olsWate	
10 1	dgiəH ətsW	
RO-	Nitrates	
NITRO- GEN A	Setinin	8.5.6.8.8.9.2.8.9.2.9.9.9.9.9.9.8.8.8.8.8.8.9.9.9.8.9
ORGANIC NITROGEN.	solved.	
ORGA NITR	Total.	
	Bus-sus	1128 1149 1159 1169 1176
EN AS	Albuminoid Dis- Div'os	656 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
Feet 14	T'10'T	86.00
Z	FreeAm- monia.	6. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
ED.	Bysuspd	ιως η περιούς ος τη τη ες τη ες τη τος
OXYGEN CONSUMED.	By Dis-	
ဝိ	Total.	
.əu	Chlorin	3121.01.000.300.300.300.300.300.300.300.300
ON TON.	pəpuəd -sus	\$\frac{1}{2} \frac{1}{2} \frac
RESIDUE ON EVAPORATION	Dis- solved.	4 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
RE	Total.	43.35.00.00.00.00.00.00.00.00.00.00.00.00.00
	Odor	Gassy None
	Color.	
APPEARANCE	Sedi- ment.	Little "" V. Little V. Little V. Little Little "" Cons'd V. Little
AP	Turb'y.	Decided Distinct Decided Distinct None Slight None Slight None Distinct Slight Distinct Slight Much Decided Much Slight Decided Much Decided Much Decided Much Slight
OF	1899 Examination.	88 88 3 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4
DATE OF	1899 Jollee- I tion.	May July 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
,o.Y.	Serial	2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

TABLE 92.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Source

Į.
H
_
á
ī
1
4
1
Y
-
ER
Y.
H
1
002
OI
Z.
Ę
LI
Ξ.
24
E
1
-
-
E
0

University of Chicago.

Report of EDWIN O. JORDAN,

No. of Isac.	Per Cubic Centi: meter.	11,100 12,500 12,500 13,500 11,100
	Presence Abs. of	
	Temper,	
ature r, C.	Tempera	
	id2joH ota7/	
RO- N AS	Nitrates	<u> </u>
NITRO- GEN A	Nitrites	595 - 21 - 25 - 21 - 25 - 21 - 25 - 25 - 2
NIC GEN.	popuged -sns	
ORGANIC NITROGEN.	Dis-	
	Total,	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
90	m k bid sus	88.00.00.00.00.00.00.00.00.00.00.00.00.0
TROGEN AS	bioniminoid -sid -sid	\$\\ \frac{\partial \text{2.5}}{\partial \text{2.5}} \\ \f
NITRO	I'10T	8668868° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° °
	FreeAm- monia.	10000000000000000000000000000000000000
ED.	BySaspd	<u> </u>
OXYGEN CONSUMED.	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
23	Total.	
16,	Chlorin	101001111001111001111001111001111001111001111
ON ION.	papuad -sus	8. 4. 5. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6. 6.
RESIDUE ON EVAPORATION	-siu boylos	8.82 8.82 8.82 8.82 8.42 8.82 8.82 8.82
RE	Tiotal.	######################################
	Odor	Gassy None
	Color.	Muddy Muddy Sanatabara
APPEARANCE.	Sedi- ment.	Little "." "." "." "." "." "." "." "
APE	Turb'y.	Distinct Distinct Distinct Sight Decided Slight Slight Slight Distinct Nuch Slight Nuch Slight Nuch Slight Decided Nuch Slight Decided Nuch Slight Decided Nuch Slight Decided Nuch Slight Decided Nuch Slight Decided Nuch Slight Decided Decided Nuch Slight
Sa.	1890 Exami- nation.	R=-5-8-656800F885658F4568-075891-588
DATE OF		% May
DA	1899 Collec- tion.	Nay June Sept. 1
.oZ	Sorlal	86.82 88.88

TABLE 93.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, HENRY, ILL.

	- 4
	Ch. convers
	-
1	-
JORDAN	9
	,
Ċ.	
INDWIN	Uniworeity
-	
	_
Cont.	
-	
-	
_	
5	
_	
~	
-	
_	
noda	
13	

No. of Bac.	per Cubic Centi- meter.	28,500 28,500 28,500 29,500 11,500 10,500 10,500 10,500 11,100
	Presenc	
	Tempers,	498888888888888888888888888888888888888
ture r, C.	reqmeT etsW lo	
lo :	idgiəH ətsW	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
00'	Nitrates	400000
NITRO- GEN A	Ritrites	- 1
IC IEN.	pəpuəd -sng	
ORGANIC NITROGEN.	Total. Dis- solved.	
	n pud	SX
oo .	Sus.	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2
TROGEN A	Albuminoid biodimis-sid b'vlos	8888888
NITROGEN	[A ['10'I']	6 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
	FreeAm- monta.	1
ED.	Bysuspd	
OXYGEN	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
000	Total.	$\frac{-1}{4} \frac{4}{9} \frac{2}{9} \frac{1}{4} \frac{1}{1} \frac{2}{9} \frac{2}{9} \frac{1}{9} \frac{2}{9} \frac{2}{9} \frac{2}{9} \frac{2}{9} \frac{1}{9} \frac{2}{9} \frac{2}{9} \frac{2}{9} \frac{1}{9} \frac{2}{9} 2$
•əu	Chlorin	83500888428444448669888667288844886488884489888648888444444448888844888848488884888848888
NC ION.	pəpuəd -sng	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
RESIDUE ON EVAPORATION.	Dis- solved.	23.25.0.0 2.35.0
RESEVA	Total.	\$38.00 \$30.00 \$30.00 \$30.00 \$30.00 \$3
	Odor	None Gassy None
	Color.	Madda a a a a a a a a a a a a a a a a a
APPEARANCE.	Sedi- ment.	Little Cons.d Little
AP	Turb'y.	None Decided Distinct Decided Slight Decided Distinct Decided Slight Distinct Much Decided Control
OF	1899 Exami- nation.	20 June 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
DATE OF	1899 Collec- tion.	Lay under un
.oV	Serial	80 May 107 June 1134 213 213 253 July 283 July 283 285 253 Sept 650 .

SANITARY WATER ANALYSIS-PARTS PER MILLION.

[LL.
VERYVILLE, I
4
RIVER,
of.
-ILLINOI
64
2.3
L
WAT
OF
(2)
RCI
Sour

University of Chicago.

Report of EDWIN O. JORDAN,

		 \$3555555555555555555555555555555555555	22
No. of Bac.	Cubic Centi- meter.	2000 2000	7.05
	Presen lo .sdA		
	Temper	<u> </u>	25 25
	Temper	្រុំនៅនាំនាំនាំនាំនាំនាំនាំនាំនាំនាំនាំនាំនាំន	- 0
	иман Бия —-		5. 00 5. 00 5. 00 5. 00
-01 A A S	Nitrates	$\frac{1}{16}\frac{1}$	2,72
NITRO- GEN A	Sitrites	<u> </u>	90.
RGANIC ITROGEN.	Sus-		: :
ORGANIC	Total. Dis- solved.		
	n pyd	 	~ 35
N A'S IA.	mA biod b'vlos	\$^4\$\\ 4\$\\ 4\$\\ 8\$\\ 8\$\\ 8\$\\ 8\$\\ 8\$\\	
NITROGEN A	Albuminoid biodos-sid b'vlos	200 4 4 4 6 6 6 4 4 4 6 6 6 6 6 4 6 6 6 7 6 7	
TIN	sinom.	\$\frac{\phi}{86} \frac{\phi}{2} \f	e er
	FreeAm-	63	5 to 5 to 5 to 5 to
OXYGEN	By Dis- solved	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	0 % 6 %
Ox	Total.		7.7
,9t	СЪТогЪ	84-44-44-44-44-44-44-44-44-44-44-44-44-4	31.
ON.	papuad -sng	# # # # # # # # # # # # # # # # # # #	0.5
RESIDUE ON EVAPORATION	Dis-	######################################	350
RES	T'otal.	######################################	385.
	Odor	N N N N N N N N N N N N N N N N N N N	10
	Color.	डां जं नं क्षे क्षे क्षे क्षे क्षे क्षे क्षे क्षे	3.6
APPEARANCE.	sedi-	LTT1 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	:
API	Turb'y.	Slight Decided Sight Decided Sight Decided Much Decided Distinct Decided Distinct Decided Distinct Comparinct Comparinct Decided Distinct Comparinct Com	
F.	1890 Exami- nation.	ក្តុក្ស ស្គ្រាស់ ស្គ	25 62
DATE OF		8 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 - 5 -	26 6
Q	1899 Collection.	May June June Sept.	
No.	Infros	全型互易器器是是是自动各级合品的影響性的特殊的容易是是是是	115

TABLE 95.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, WESLEY CITY, ILL.

Report of Edwin O. Jordan,
University of Chicago.

No. of Bac.	per Cubic Centi- meter.	932,000 1,410,000 1,575,000 1,575,000 1,575,000 1,575,000 1,
	Presenc	
	Tempers,	28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
sture r, C.	Tempers	18 18 18 18 18 18 18 18 18 18 18 18 18 1
	Height eta'll	
NITRO- GEN AS	Nitrates	### 1
RIN	Nitrites	25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
GEN.	-sng	
ORGANIC NITROGEN.	-sid solved.	
	Total,	
	E Proud	98888999988899998989999999999999999999
EN AS	Albuminold I'10T	88.88.88.44.88.44.88.44.88.44.88.44.48.88.44.88.44.88.44.88.44.88.44.88.44.44
NITROGEN AS AMMONIA.	I'10T	20 20 20 20 20 20 20 20 20 20
	FreeAm- monia.	4.55 123 120 120 120 120 120 120 120 120 120 120
N ED.	BySuspd Matter.	64444 6 6 6 6 6 6 6 7 7 8 7 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9
OXYGEN	By Dis- solved.	అయ్టార్గాళాచించిన చివ్వార్లు మార్జులు ఉంది. ఆయ్టానుని కోవేనాట్లో నానే కోవవేచినకోకుకుకుకుక్తి
000	Total,	လွှာလေးလွှာလွှာလွှာလွှာလွှာလွှာလွှာလွှာလွှာလွှာ
16.	Chlorin	######################################
ON.	bended Sus-	<u> </u>
RESIDUE ON EVAPORATION.	Dis- solved.	888 888 888 888 888 888 888 888 888 88
RES	Total.	33.5.1.1.4.1.4.2.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3
	Odor	None
	Color.	Multiple
APPEARANCE	Sedi- ment.	Little Cons'd Little Cons'd Little Cons'd Little Cons'd Little Cons'd Little
AP	Turb'y.	None Decided " Distinct None Decided Slight Distinct Decided Distinct Decided " " " " " " " " " " " " " " " " " "
OF	1899 Examination.	29 June 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
DATE OF	1899 Collec- F tion.	May 29 June 7 " " " " " " " " " " " " " " " " " "
.oV	Serial	1120 1130 1130 1130 1130 1130 1130 1130

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, PEKIN, ILL.

JORDAN,	of Chicago.
Ö.	sity
EDWIN	Univers
10	
Report	

No. of Bac.	per Cubic	Centi- meter.	120.000	512,000	129.000	205,000	55.000	2.030.000	500,000	1,435,000	470.000	580.CEO	1.940.000	989,000		90.000	020.000	310,000	240,000	120,000	500,000	430,000			000.000	000.000	000.000	380,000	140.000	10,000	5,000	20.800
re or flo')		Pres.				:			:	:			:	:	:					:		:	:	:			:				:	
orithe .U.			33	.05	31.	30.	3.5	٠	8	33	32.			Si 8	. 66	365.	70.	3.5	. 153	<u>×</u>	ें. हिं	<u>01</u>	<u>.</u>	9 =	- 10	16.	2 16	: o.		-	0.5	27
ature r, C.		M lo	∞.		26.25	24.	.5ç	200		86 E	20.	9.6	Si 3				3	38.	.03	15.	<u>:</u>	≘i:	<u>.</u>			. 0		C				
101	:НЯ э1в		3.5	3.6	17	က က (၁)	si,		_:		. 1	<u>.</u>	· .	. 10	÷ 11	c	-		9.1				_; -	, sC	-	· G	, <u>-</u>	-6	10	ei.	S) S	e 22
-01 V A S	89	NHTE	00	Τ.		- Sec. 1	200	5.		1.45	G 6	દિ ૧	٦ - ا	- 19	3.10	3	ıc	69	64 .	6	- -	2 :	,	-	100	_	.55	18	1 555	10	음: 	~
NITRO- GEN A	80	MILLIN	80.	104	33	887	25.	.26	20.00	si s		61.	+ 3	77.0	100	0.03	.056	.076	.07	.076	91.	<u>ज</u> ं ।	20.00	0	: 19	-	: ::	.034	.056	1011	900	240.
GEN.		puəd sns		:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:	:	:						:	:		
ORGANIC NITROGEN		SIU SOTOS		:	:	:	:	:	:	:	:	:	:	:	:			:	•	•	:	:	:					:	:	:	:	
	1 1	b'baq stoT	:	: %	: ::	. 10	:			200	:	: :	:	:	:	00	£ 53	:	: &	·	: :	:	: 02		<u>2.</u>	<u>∝</u>	6)	61	: 3	: 83	: 200 200 200 200 200 200 200 200 200 20	0.0
As 1.	oid Am	-sng		_	_	_	_		_	368	_					_	_		_	_	_	_	_	_	_			_	-			-
TTROGEN AMMONIA	uminoid	-si(I b'vios		_						.432											_					_					212	
NITRO	Alb	l'ioT	.36	27.	7	10. 53.0	<u> </u>	35.5	9	× 1	00.	201.	9.	:=	666	3 5	9.	.59	13	9.	3 (in in	00.	100	10	51	96.	.3.5	<u>ac</u>	<u> </u>		210.
		4991'4 mom	.64	3	.92	7	φ		97.	200	00.	+ ;	, n	6. 6.	204	.65	.336	.218	οó	344		× 6	1 26	3.5	19	2.56	33	1.76	1.46	- 8	32. c	01 0
EN IED.	of pds	Bysus	0.	7.	90 +	٠٠. د د د د د د د د د د د د د د د د د د د	-: 0	.; 0		20 C	_	25.0						-1	7.	9	-		0 0	2 -	Si	15	9.9	9	?? ??	-	£ :	1
OXYGEN CONSUMED.		id val	1 9.4	_			z. ≀		ė,	o r	4	9 11				2.0					7.6			000	6.9	6.7	7.5	5.2	10	5.0	- 0x	
3	l. I	BIOT'	8	- 6	27		<u>+</u>	<u>ه</u> د	oc o		0 0	-0	Ċċ	_	2 16	6 0	G		10			_		. 5	00	(-	9	5.	20	10		s l
16.	_	СРИ	15.	2 15.	=	123	= :	20	7.7	- C	- 40	200		27.	11.	= =====================================	200	50.	£.	533	61	53	67.	293	30	20	48.	:8:	<u></u>	36.	% 5°	-0.4
0N 10N.	pa	puad sus	18.	103.	× +	38	33	£ ;		23.5				210		9.00	11.	37.	689	26.	8:		20	2 6	2.	23	16.	1.	10.	∞:	± ÷	?
RESIDUE ON EVAPORATION	.b	aatos sid	356.	336.4	318.4	207.6	550	3552.x	200	- T-S-	0.110	.)(.)-[000	260		36-1	336	360.	3.00	372	9	.511.	411	400	390	361.	388	350.	367	370.	366	101
REYA	1.	niol'	37.1.	459.6	366.1	358.	379.2	200	X	386	535.	515	Ç.);	501. A16	2710	394.	380.	397.	386	308	406.	39%.	. 50 t.	22	411	388	101	361.	377.	33.50 3.50 3.50 3.50 3.50 3.50 3.50 3.50	155 155 155 155 155 155 155 155 155 155	. (2)
	Odor		None	•		: :	• •	•		9 4		- 4				:	9.9	:	:					9.9	9 9	9.9	9.9	• 9	9,	3.9		
-		Color.		.e.	ıc.	- j. 1	اخا ،	ugu s	년. 1	ر اد اد	9,-	ų.	÷::	· -	-	Muddy	?	.65	23	- !	6		. ·	200	?}	-	- c:	???	er:	89	د زد	•
APPEARANCE.		ment.	Little	* *	9 9	9		. :		9 7	: :	Transfer). Little	amr	1 11+1100	Little		V. Little	p,suo,)	Little	e ;	: ;	: :	:	*			•	9 +	:	: :	
AP		Turb'y.	Distinct	*	Decided		: ;	None		Decided	4 12 2 2	JUKIN.	DISTINCT	THE SELECT	- Tireba	Distinct	Slight		Decided	4 4	-	Slight	; :	Dietlnet	Decided	Much		Slight		Much	: •	
30.5	1890	Collec- Examition.	May 25	:: 31	6 June 7	+1	:	6: 1	July	* :		95	Nun.	: :	1	3 5	Sept. 6	.:	(17	:	Oct. 1	= :	2 !	1 30%	6	16	:	Dec. 2	oc :	= :	÷ :	
DATE OF	068	Collec- I tion.	May 24	. 30	June 6.	: :	•	:	July 6.	27	2	£ .	A112.		6.1	1 2	eritin u	6	× 1×	98 ,.	ort. 30	01	X o	02 3 3	さい シング	- 10			Jer. 5		<u>6</u> 2 : :	X.7.
Vo.	[13]	.ms	55	{-		137	521			8	200			2.2	23.	282	3000	613	525	693	1330	071	53	2000			000		tures	1053	<u>×</u> :	60

TABLE 97.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER -- ILLINOIS RIVER, HAVANA, ILL.

Z
-
< 0
-
0.00
ORD
\sim
7
0
_
WIN
-
200
ED.
)—(
5-3
print.
Egung
0
~
-
نې
2-4
0
eport
C.
-
a)

University of Chicago.

No. of Bac.	Cubic Centi- meter.	68.800	
	Presen To .sd A		
	ragmaT riA lo	888535348888888888888888888888888888888	
	reqmeT of Wate	<u> </u>	
tot a.re	ngieH otsW		
RO-NAS	Nitrates	4.24.48.88.82.41.88.85.80. : : : : : : : : : : : : : : : : : : :	
NITRO GEN	Nitrites	9 = 8 = 1 = 1 = 1 = 1	
GEN.	Sus-		_
ORGANIC NITROGEN.	Total. Dis-		
	g sng	844511100644885188 1448884111111111111111111111111	
EN AS	Albuminoid b'vlos	= = = = = = = = = = = = = = = = = = =	
NITROGEN AS AMMONIA.	Tot'l	8884628444644656888888	 }
4	FreeAm- monia.		2
ED.	BySuspd Matter.	000 0000000000000000000000000000000000	
OXYGEN	By Dis-		>
Co	Total.	<u>w-cacacanoren cacacanoren ar-xar-</u> <u>wur-anamerro cacacanoren ar-xar-</u>	
ıe.	Chlorin		
ON ION.	bended Sus-	\$\$\$\alpha\text{8}\text{8}\text{8}\text{8}\text{1}\text{8}\text{9}\text{9}\text{1}\text{8}\text{9}\text{9}\text{1}\text{8}\text{9}\text{9}\text{1}\text{8}\text{9}\text{9}\text{9}\text{1}\text{9}\text{9}\text{9}\text{1}\text{9}\text{9}\text{9}\text{1}\text{9}\text{9}\text{9}\text{9}\text{1}\text{9}\text{9}\text{9}\text{9}\text{1}\text{9}\text	202
RESIDUE ON EVAPORATION	Dis-	28 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	.010
REN	Total.	888 888 888 888 888 888 888 888 888 88	000
	Odor	NO N	
-	Color.	N Baisia aroromonos – generomos generomos – generomos V	
APPEARANCE.	Sedi- ment.	V. Little	
41.	Turb'y.	None Decided Distinct Decided Distinct Slight Much Decided Distinct Much Decided Distinct Much Decided Much Decided Distinct Company C	
OF	1899 Exami- nation.	S S S S S S S S S S S S S S S S S S S	2
DATE OF	1899 1899 Collec- Exami- tion. nation.	the many many makes the market makes the	
		778 May 1778 May 1772 June 1772 May 177	
.o.Y	Serial	25	7

TABLE 98.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-SANGAMON RIVER, CHANDLERVILLE, ILL.

University of Chicago.

Report of EDWIN O. JORDAN,

No. of 13ac.	per	Centi- meter.	8.400	11.600	1,830	3,100	8 350	2,100	9,900		1.200	1,900	11.500	4.200	1.300	2,000	3,200	1.500	10,800		2.900	(C)	1,200	6.400	1.900
ce or Colf.	nə lo	Pres.					-								:										
		Tm9T To	24.	C 3	300	83	33	655	.6~	-66	66	21.		16.	33	33	55	œ	-	16.	7	=	T.	_	- 1
		Tenij	33.	26.5			107	26.	35	.36.	27.	5	19.	į-	6.	· · ·	00.00	10.	10.	2	18.	oci	33	4	÷į.
10 1	Б1в	1911 77	∞ ∞	œ.	 		· ·	33	3.6	ıÇ.		Low	:	:	9.9	4 0	•	:	_	_	7.1	_	-71	1.6	3.6
RO- N AS	Vitrates		.e.		_	_				_		_		_	_		_	_	_	_					
NITRO- GEN A	sə	Mitrit	90.	<u> 190</u> .	.056	.018	.016	.00	.048	910.	10.	.006	300	С.	<i>ج</i>	.005	-:	:006	.006	.006	£.0.	0.	900.	100.	.015
GEN.		sus	:	:	:	:	:	:		:		-:		:	:		:	:	:	:	•	:	:	:	:
ORGANIC NITROGEN.	I -	Solve		:	:	:	:	:	:		:	:		:	:	:	:	:	:	:	:		:	:	
3%]'	r bad	:	:	.:	:	:	:	:	:	:	:		:	:	or.	-	-	· ·	:	:		:	6	9
oo	id Am	p,pud -sng	.112		_		_	_		_		_	_				_				-		_		
GEN A	Albuminoid	Dis-	33	.24	152	c;	.128	807.	1964	. 932	. 136	197	<u>:</u>	.000	.056	980.	101	. 128	.088	.088	000	088	126	.136	800.
NITROGEN AS	Albu	Totyl	.432	67.c.	.352	304	27.5	.208	:33	.264	.368	.261	£.	.218	.136	.381	3033	.152	.176	880.	216	.168	.136	. 15g	0.24
		FreeA		.0 <u>.</u>	.034	.062	.055	.014	80.	.038	10.	.0220	:0:	S	.008	0.	07.1	.026	.062	30.	860.	.056	o: 0	.108	.161
ED.	a pd	susya Matte	تن. تن	1.6	30	00	5.4	83 83	30	€. 5.	3	2)	03	©.5 ∞	- I	6: 1	1.4	9.	e.	0.		00	??	6)	e:
OXYGEN CONSUMED.		By Dia	9.6	5.1	4.6	Ci.	1.1	-	50 10	3.6	os oc	3C.	3.1	25	31 30	2.6	2.6	3.6	1:	∞. -	3.6	31	5)	₩.	2.6
000	l.	RIOT	15.1	~	8.1	6.	6.8	4.3	· - 5	.9	30.	5.	5.4	5.6	3.5	4.5	÷	1.2	1.	∞. -	4.6	65	00	63	₩ 90
*ət	ца	Срјо	2.35	3.5		3.6	- ;	-	6.4	5.6	1.5	4.6	1.7	00	1.6	 	7.	6.4	7:0 00	6.2	9		6.	. č	5.
ON ION.	pa -	bende Sus	l					F.			•	162.	.86	13%	33	56	.99	5.	19.	16.	46.	ص	16.	% %	73.
RESIDUE ON EVAPORATION.	- p	si(I ovios	216.	240.	3	287.2	354.4	290.	218	300.		202.	271.	38	25.58	312.	288	293.	285	304	386	355	300	286	263.
EVA	1.1	RIOT	356.8	380.	141.	421.6	395.	355.	434.	370.		361.	373	(E)	350.	338.	351.	338.	301.	320.	334	331.	316.	311.	336
	Odor		None	9 9		9 .	9 3	9.9	10	9 9	9 *	* 9	:	:	-d -0	;	:	:		. 9		:	:	:	*
		Color.	??	9.	0.	00	02	9.	-	0	Muddy	:6	0	.e.	() ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ? ?	-	_			0.	:0:	3)	<u>.e</u> .	0.	υį
APPEARANCE.		Sedi- ment.	Much	Little	Cons'd	, .	Little	V. Little	Little	9 .	Cons'd	Little	Cons.d	Little	V. Little	Little	9 4	9.9	7 9	9 9	9.9	V. Little		Little	9.9
1.		Turb'y.	Decided	-	9 .	:	:	Distinct	Much	9 *	:	Decided	Much	Silght	Distinct	9 9	Slight	Decided	Slight	Decided	Distinct	Slight	•	Decided	Much
OF	1890	Exami- nation.	une 1	coc	· · · · · · · · · · · · · · · · · · ·	niy 6	13	.115. 3	10	:	ept. 1,	-	:: 21	\$i	et. 5	61 ,,	1.	10v. 2	01 ,,	" 16	.66. 33	Dec. 1	1-	· 1 1	56
DATE OF	0881	0	y 31 June	ne 7	<u></u>	Jy 5 J	22	12. 3.A	c.	16	31 Sept	11. 13	S	34	L. 40	x	33		α	10	33	0 6%	ω.	13	ନ
			90 May	116 .lune		is Ju	. 0	11 1. 868	. 3	*	· + 400	356 Sep	6 19	:	5 Oct.	•	· ·	5 Nov	50	:	71	3	6 Dec	; O	3.
.07	[8]	i102	C	11	X -	100	000	955	* *	173	5.0	65	1 × × 3	715		- - - -	び、文	233	555	35	185	100	1056	1090	1126

TABLE 99.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-ILLINOIS RIVER, BEARDSTOWN, ILL.

Repo

EDWIN O. JORDAN,	University of Chicago.
ort of 1	

No. of Bac.	per Cubic Centi- meter.	9.300 9.
	Presend Abs. of	
	reqmeT ria to	9 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	Temper of Wate	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 9
	dgiəH ətaW	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
30- N AS	Selentiv	
NITRO- GEN A	Nitrites	\$\frac{1}{2}\text{3}\text{2}\t
IC TEN.	pəpuəd -sng	
ORGANIC NITROGEN.	Dis-	
OX	Total.	
Tr.	E Sus-	08 0.00 0.
NITROGEN AS	Albuminoid Dis- Dis- Divid	821-23-488888888888888888888888888888888888
MITINAMIA.	I'10T	8.8.8.4.4.4.4.4.8.8.8.8.8.8.8.8.8.8.8.8
73	FreeAm- monia.	11.00
N ED.	Bysuspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN CONSUMED.	By Dis-	$\overline{\bigcirc}$
Cos	Total.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
.əu	СБРоті	07-0012888888888884044488888888888888888888
ON.	penged -sns	<u> </u>
RESIDUE ON EVAPORATION	-si(I solved	2
RES	Total.	88 88 88 88 88 88 88 88 88 88 88 88 88
	Odor	None in the second of the seco
	Color.	
APPEARANCE.	Sedi- ment.	Much Little Cons'd Little Nuch Little Cons'd Little Little Cons'd Little Cons'd Little Cons'd Little
AP	Turb'y.	Decided "" Distinct Decided "" Slight Much "" Slight Much "" Decided Much "" Decided Much "" Decided Much Decided Much Decided
3 OF	1899 Examination.	May 59 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
DATE OF	1899 Collec- tion.	May 5 June 1 Jun
.o.V	Serial	### ### ##############################

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER-ILLINOIS RIVER, KAMPSVILLE, ILL.

University of Chicago.

Report of EDWIN O. JORDAN.

No. of Bac.	per	Centi- meter.	8,490		7,750	009	006	2.950		550		710	340	4.400	3.300		520	5,100	1.300	5.300	1.800	19, 100	4.3(0)	3,500		23.500	
		Prese						•	•																		
		Temp of A	<u>∞</u>	33	5	:::::::::::::::::::::::::::::::::::::::	35.5	37.	31.	25	350	35	600	×	90	19	6	11.	53	o.	. ic	23	10	6	_	200	:
		qm9T s7/ To	.55	.56.	100	.667	500	2. 2.	15.	33	65	65	60	6:	17.	16.	- 1	1%	17.5	21	2	3	-	30	7	2.	:
	ogu Zpi	Hely 17.	91.	17.1	16.9	16.	15.9	15.9	16.5	5.5	15.3	15.1	15.3	15.6	15.3	15.3	15.3	5.5	20	15.6	5.7	16.	15.7	15.6	× 10	19.	
RO. NAS	89	Mitrat	.83	F	.49		1.4	16:	1.13	7	10	13	50	10	100	9.	16.	50	7	.C	18	00	:0	1.0	8	1.15	
NITRO GEN A	<u> </u>	Mitte	.43	.031	.07	***	91.	80.	80.	0.045	-	8	.07	980	.016	.034	120.	.03	10.	980	23	90.	031	10.	18	030	:
NIC GEN.	_	-sus		-		:	:									:	:										:
ORGANIC NITROGEN.	-	Sid					:		:	:	:				:		:	:									:
		b'bnq stoT		€		·:	.: 0	23	:		98	99		1	×:	??	.: 	+(2	эc	2)						
AS.	oid Am	-sng	<u> </u>	,	80. 9		_		•							_	_		_		_			÷	÷	06.	
TEOGEN A	Albuminoid	-sid	.33	3.	.336	.31	हुं.	67:		86.	.95	.38	35.	63	33	. 19		0)	÷;	53	30	0.0	121	200	G	.16	:
NITCOGEN AS AMMONIA.	Alb	['101'	.64	.464	.416	.35	33.	.368	.376	280	.395	7	358	395	37.5	304	.311	.314	.261	311	1.072	.661	36	5.5	3	.36	:
		FreeAi		.068	.05	99.	80.	.044	.03	.058	.068	188	.156	.366	.358	256	.168	. 198 861	100.	× .	12.	9.	1.398	1.312	1961	37	
EN.		Bysns	c.01	c.	4.8	2.1	63	2.6	3.6	e 5	9.	}	1.1		-	with	week	=	_	under	कुर	G1	X.	IC.	61	C:	
OXYGEN		By Dis	2.0	2-	3.5		4.6	3.6	00 T	10	16.4	6.1	6.1	6.	6.4	6.4	6.6	30	6.	6.4	6.6	5.3	10	3)	5.6	5.6	:
23	1.	rio'l'	18.6	o.	1-	6.2	1	6.9	00	\- -	{-	6.8	-1:	-1	6.8	6.8	{-	6.9	{-	6.X	<u>-</u>	×	6.3	65	×	6.5	
,91	ıi1	СРГО	80. 80.	6.5	00	7.5	30.	21.	∞ •							13											
ON.		puad sus	518.	221.6	76.	30	55	22.	264	36.	39.	39.	.40.	104	300	50.	œ.	300	× ×	10.	41.	31.	20.	0	30.	62.	
RESIDUE ON EVAPORATION	.h	sid evios	308.	8.032	9.1%	352.8	342.	383.	200	306.	290	275	316.	215.	57.67 67.67	26×.	3336	326.	3/16.	342.	320.	324	328.	358	320.	33	
RES	.1	втоТ'	7:36.	502.4	357.6	356.	355	310.	100	315	320	314.	356.	346.	320.	318.	316.	354.	351.	352.	364.	358	348.	358.	340.	341.	
	Odor		None	* *	9 +	7 9	*	;	:	:	:	:	:	9 .	:	uir +	:	•	9.9	:	9.9	9 .	3 3	;	. 9	**	
		Color.	63.	9.	~	ಣ	er;	.6.	7		18).	.č.	***	16.	15.	زن	ic.	<u>. I</u>	??	ा		10.	33	Muddy	10	33	
APEARANCE	;	Sedi- ment.	Much	p.suo.	Little	9 .	;	V. Little	Much	Little		9 9	•		Limbe	*	:	:		3 9	9.	4.4	9,9	;	9,9	9 9	
11.		Turb'y.	Decided	:	:	:	Distinct	Decided	Much	Decided	:	Distinct	Much	:	•	:	9 9	Distinct	*	Much	Decided	Much	:	-	Much		
) F	1400	Collec- Exami-		66.	11. 6	2	100	50	27	3	ut. 1	X	444		657	t. ?	-,	02	100	17.	16	~~~	c. 1		10	553	
DATE OF		c En	31 June	=	5 July	, ()	. 92	2.11			30 Net		:	- 03	- 2	4 Oct.	-	x	. 6	1 Nov	10	300	29 Dec		63	02	:
L)	1800	Colle	May:	June	July	:	:	Aug.	:	9 9 9	99	Serpt.	:	7	ু ৷	Det.	:	:	*	Nor.	, ,	71	*	Dec.	1 99	3.	
Serial Zo.		25	1.3	276	325	:,6;5	341	150	510	545	5.40	17:0	(96)	1111	7.15 (メート	ic x	75%	910	696	3.5.	1001	1069	1007	1138		

STREAMS EXAMINATION-SANITARY DISTRICT OF CHICAGO.

3	
-1	
_	
-	
_	
-	
Z	
FTON,	
\sim	
md	
F	
-	
40	
24	
_	
IIIA	
9	
20	
EE	
ER,	
line.	
_	
-	
_	
_	
m m	
2	
158	
OIS	
NOIS	
NOIS	
~	
~	
Ţ	
Ţ	
Ţ	
Ţ	
Ţ	
1111	
1111	
1111	
1111	
1111	
1111	
ATER-ILLI	
ATER-ILLI	
ATER-ILLI	
WATER-ILLI	
OF WATER-ILLI	
OF WATER-ILLI	
OF WATER-ILLI	
-ILLINO	

University of Chicago.

Report of EDWIN O. JOHDAN.

No. of Isac.	Cubic Centi- meter.	2000 2000	-50,000
	Presen		
	raqmaT riA lo	######################################	-
	reqmeT of Male	्रित्रा श्रेष्ठ संस्था संस संस्था संस्था संस्थ	
	dgiəH nsW	: ::::::::::::::::::::::::::::::::::	-
RO- N AS	Sitrates	あっては、	ç.
NITRO	Nitrites	200 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0.30
	solved.		
ORGANIC NITROGE?	Total.		:
	F p,pud	66888888888888888888888888888888888888	. 250
A. A.S.			0/1
NITHOGEN AMMONIA	Albuminoid Pis- Olv'd	25	
Z	FreeAus. ntonia.	"qq : 58998845888888888888888848484899999	.
	Bysuspd		
OXYGEN CONSUMED.	By Dis-	1-11 00 00 1-00 00 10 10 10 10 10 10 10 10 10 10 10 1	۰ ۲
000	Total.	ု တယ်လွှာတွင်တွေလ ႏုတ္တန္း တွင်းသည်း ႏုတ္တန္းတွင်တွေတာ့ တွေလတ္တတ္တည်း မြင့္ချေပးသည့် တွေလာက္တတ္တည်း မြင့္ချေပးသည့် မြင့္ခြင့္သည့် တွေလာက္လည္း မြင့္ခြင့္သည့် မြင့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည့္သည	0.0
ne.	Chlori		
on ION.	pəpuəd -sng	80558888888858588589848588-85789+848	198.
RESIDUE ON EVAPORATION	Dis- solved.	2888	512
REE	Total.	00 05 0500 05	- DQ-
	Odor	None : ::::::::::::::::::::::::::::::::::	
	Color.	×	
APPEARANCE	Sedi- ment.	Cons'd Nuch Little Nuch Cons'd Little	
Υ	Turb'y.	Decided "" "" Distinct Decided "" "" Distinct Decided Nuch Distinct Decided Nuch Distinct "" "" "" "" "" "" "" "" "" "" "" "" ""	
OF	1899 Exami- nation.	May 10 May 10 June 15 June 15 June 15 May 10 June 15 May 10 June 15 May 10 M	85
DATE OF	1899 Collec- E: tion. na	0724 2 1 2 8 1 2 5 8 2 6 5 2 8 8 2 1 1 1 1 2 1 2 1 2 1 2 1 2 1 2 1	
Zo.	Serial	22 May 1177 Jun Jun 1177 Jun Jun 1177 Jun Jun Jun 1177 Jun Jun Jun 1177 Jun	

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER MISSISSIPPI RIVER, GRAFTON, ILL.

JORDAN,	of Chicago.
Report of EDWIN O.	University

No. of Bac.	per Cubic	Centi- meter.			0000	96.000	31,000	16,000	9,000	000:	000,61	0.900	OOM-T	1.300		006	1,400	 	000.4	2000	4.500	1.500	900	999 Vin	007.5	6.300	1.700	1.100	200	3,100	1.),(147)
ce or Coll.								:	:	:	:	:				:	:				:	•	:	:							
ature , C.	per Alr				26.		530	31.	<u>.</u>	51 S	.50	2 2	3	000 G1	g;	3.				80	ි දි	G) G		2 1-	=======================================	1-	21	9:	D	.; - C	
ature or, C.				:	333	ê	₹	10	26.5	: : :	. 000	88	ο ο ο ο ο ο ο ο ο ο ο ο ο ο	26.		.90	<u>.</u>	3,5	: : 3:	<u>x</u>	16.	16.	. # . #	0.0	<u> </u>	্	\ 	ec:	er:	_ c	•
10 1	igh Sate		17.	10	16.1	16.7		:		9:50	20	- 32	6.1	5.6	6.	٠ زه	က :	त्रं न स्	- 00	7		ه ود د ود	. T	2 10	9	2	6.5	- co	÷ ::	7:-	
30- V AS	891	Mitra	<u> </u>		ે !~	36	.5.	.45		500	; <	000	. O	0.	0.	0.	0.	ن د رو	15	0.		0.0		e re	: 03	(3)	15.	0.	-	- ×	
NITRO- GEN A	sət	intiN		900	015	.005	.01	.012	8.	0.8	3	00	9.0	00.	.006	0.	0.	900	000	900.	0.	0.0	, c		. 9	3	0.	0	0.	9.6	
NIC GEN.		bene						:	:	:	:				:	:	:	:			:	:	:					:			
ORGANIC NITROGEN.	-S	Tota		:				:	:	:	:					:	:				:	:	:						:	•	
		p,pud	9.	:	7.68	8	80	: 75			++	94	800	96	84	:	:	: 20 00 20 0	16	1.5	36	36	:	9	: : : : : : : : : : : : : : : : : : :	-		??	56	101	
AS.	5	o'vios esus	<u> </u>			_	_				_				_	•							_			_					-
NITROGEN AS Ammonia.	Albuminofd	-sid	1		. 352	_	_	_	_		_					•	_				_									200.	
NITE	TV I	[,10[,	re;	1.13	1.1.			_		-		-		-		•	_	_		_	-										
		Free	60.5	70.				•	•		•	•		•		.03 .03 .03				•	0.28	38	50	50	.036					9 2	
GEN IMED.	pds	Alos Bysn Hatt	0.0	2. 7. 2. 7.	11.5	6 10.4	00.2	∞: ∞: ∞:	⊃. ∵.	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	3 -	6 3 6	\$ 00.00 T	ري ده ده	50 50 50 50	9.1	0.00	2 2 - 2:3 2 - 5	000	× 1.4	67.1 67.1	x =	7. → 	* OU.	∞ ∞	701	3.1	Service Contract Cont	-	ni Si oc	
ONYGEN	-81	Total			22.5 11.			5.9 14.				0)				3.6 8.	200		2	.2	. 1	υ. υ. υ.		0.8.0	3.8	0.8.0	œ	9.6.9.	.2 10.	= x0	
ne.	ilio				9 63	9.1	∞.	6.1	× :	ے <u>ب</u>	- 1	10	00	6:0			1	- 00	, α.	10.	9.5	رة د د) id			G)	3 6 1	o≥ cc	3.4	x & x	:
7 %	geq s.	nen Su	336.	. SS	38.38	86.4	59.6	55.6	31.6			91.0	62.	083	54.	E	9.0		3.5	92.	· 章:	8 8		 		2	11.	%I.	 		
RATIO	.be	Alos	140.	G	159.6	14.8 13	5.6 3	55.	20.1	2) 4	0.00		×	57. 2	.89		÷ 3			×.	⊋;		: 5	90	ب	1	. 28		. 23		
RESIDUE ON EVAPORATION		Total	l	- 3	898.4 13	1.9 1	5.9 11	417.6 15	9.6 14	23.00		_																			
		10[]	e 476.	- 2. i.ī	96	153	जि	4	XX 6	303	-00-	276	G.	të.	3	: E	S 8	6 1	ं हें व	왕	-	\$ \$ \$ \$ \$	§ €	50	50	55	1-	17	<u>x</u> 8		
	Odor		None	, ,,	9 .	9 9	9 9	9 1		: :	9,9	9	1 7	9 9	9.9	: :	9 9		:	3	3 :	: :	4	13	43	9 9	:	:	9 1	11	
101		Color.		:		<u>.</u>	οc.	ж. 	7.	ر م	Ç –	- 10	e.	65	==	0,1	S :	S -	?}	-	- e.i .	T. 0	jes	· 67	8	ιc	9.	9.	£} :	ing eri	
APPEARANCE.	Sedi.	ment.	Much		9.	**	* *	Cons'd		Consid	nemo	9 0	Little	19	9 9	9		:	Cons'd	99	Little	Cons d	Little	7,7	p.suo.)	Little	9.	:	: :	: :	
A.		Turb'y.	Much	Decided	Decided	**	. ,			: :	*	Much	Decided	Much	9 9	Decided	40 - 41	Much		9 7	Slight	Much	Much		*	:	:	:	**	: :	
(See				53.5		œ	15	÷13				3 67	60	Ξ		₹ 72.0		x =		230	9	_ ;) <u>_</u>	16				15)
DATE OF		1	10 May		- Parent	;	,, 11	: :	: " : " : "	5 July 6		26 1	2.1119		16 44	÷ :		1dəs 9	30		4 Oct.	: : :	-	; ;	33	39 (20)	29 Dec	,, G	13	: :	
1).4	1899	Collection.	May	: :	;	June	:	:		7 III ?		:	1112.		:	:		Sept.	2	:	Oct.	: :			.,	;	:	Dec.	:	: :	
.07	lai-	108	क्ष	218	1 4	115	145	200			21.	125			7	512			333	100	-	0 37.5		935	15	166	9601	1058 1	9891	X 16	1.7.1

TABLE 103.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM ILLINOIS SHORE, ALTON, ILL.

Z
4
a
~
=
0
-
0
\cup
Z
25.4
\equiv
6.3
12
4
0
2
)ŗ
0
2
60
9
2

University of Chicago.

No. of Bac.	per	Centi- meter.		3.470	17.400	99.90 8.000	14.500	11,000	39.000	5.000	5.500	3.000	5.500	5,500	9,500		1.200	006		3.000	2.300	2,200	1.700		2,600	6.250		2,700	1.800	008:1-	12,700	
to 95 Coli.		Pres. o						:	:	:	:	:	•	:					:	:	:	:	:	:	:	:	:	:	:	:	:	
		qm9T a 10		16.	30.	16.	30.	21.	33	27.	28.	31.	33	??	88	30.	œ.	6:	કો કો	30.	19.	16.	· ·	20	19.	10.	<u>2</u> 2	13.	?		·	:
ature er, C.	er ete	qm9T W lo		<u>×</u>	21.	18.	19.	19.		. 36.	.68	.88	30.	39	36.	.98	 28.	30.	- 		18.	15.	16.	<u>x</u>	18.	≘	6	10.	ထင်	*	0	:
10 t	911 Q.	i9H W		17.5	16.5	16.1	17.	14.4	13.5	15.3	15.8	ર દે. દે	6	-T	7.1	£ .5	3.1	6.2	3.6	4.	က်	∞ ?:	es .	-: -:	6. T	ලා ලා	3.9	5.4	∞. ∞.	ಬ	60	
RO- N AS	sə	Mitrat	ಚ	13		1.4	تن	2: 2:	ri i	- -	ಀ	.35	∞.		.15	4.	.05	Γ.	ાં	ಀ	77	.05	3)	ig.	٠.	<u>ાં</u>	.15	ಲ	7:	सुर	. 55	
NITRO-GEN A	sə	Mitrit	.024	.03	.038		900:	.014	.01	.016	 99:	810.	810:	.038	9.	.0 .	.13	10.	.076	.013	.0075	.00	0. 2.0.	800.	<u>ම</u>	600:	10:	860.	.013	800.	.013	:
HC BEN.		sng		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
ORGANIC NITROGEN.		SiU solve	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
ōZ	1.	rota	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	
	d Am	p,pud -sng	.15	99.	.376	.224	.256	.104	. 193	880.	.072	920.	840.	104	:	960.	80.	.124	.168	.152	.136	.208	ં	960.	80.	.232	.16	. 232	.088	:	308	
TROGEN AS	Albuminoid	Dis-	.26	. 28 82:	.264	.3396	596.	.336	. 288 882	.352	.336	.336	: : : : : : : : : : : : : : : : : : :	184		196.	1994	816.	. 192	કરું	÷	.152	3	527	.256	.264	.208	.208	.208		.193	:
NITROGEN AMMONI	Albu	Torr	.41	œ.	. 64 10	.56	.52	7	8 † .	7	£0¥.	395	23	886.		.36	÷08:	.372	.36	.352	.336	. 36	- j e	جن وي	.336	.496	.368	44.	966.		₹.	:
		A9917 inoni	.025	840.	.065	9+0:	.078	070.	.061	.056	.063	60.	.038	860:	.114	60:	.03 4	.072	.03	840.	.062	.048	820.	910.	3 3	104	£60.	980.	.324	.13	.17	:
ED.	pd r.	Bysus	3.9	∞ 3)	11.9	7.6	-1	1.6	က က်	0.	9.8	<u>က</u>	3.1	3 3	0.7	3.6	હો	0.	9.	9.	-:	7	351	1.5	0.	3.	3.6	9.	9.	<u>∞</u>	1.9	:
OXYGEN		By Dis	10.	9.4	∞ ;	7. (3)	·-	∞ e3	ت. 	œ.	5.6	5.6	10	67:	4.9	5.1		8.9											· ∞	6.3	6.3	:
Coy	J.	rota			19.9						14.2	9.8	 	(- {-	14.6	~ 00	??	6.8	7.6	∞ ∞	6	G	т С	9.8	7.6	8.4	8.6	15.	8.6	∞	?? ∞	
re.	ıi1	СРГО	1		æ.	ಣ	9		જર	કર <u>ે</u>	4	÷	6.	15	12	70	6.	{~	9	6	{-	91	9	0	Ö	=	10	10	5	0	0	
ON ION.	pa	pengd Sus	342.		667.2			168.	188.	6. 6. 6. 6.	122.4	٠٠ - :	6.5	56.	#00°	183.	35.	76.	95.	106.	64.	. 80	64.	20	21.	<u>.</u>	70.	.89	36.		16.	:
RESIDUE ON EVAPORATION.	.k	Solve	214.	189.6	195.2	232.8	189.6	208.	182	178.4	181.6	204.8	254.	236.	243.	168.	215.	178.	191.	210.	188.	178.	186.	204	230.	184.	204.	179.	210.	300.	.33e.	:
RESEVAL	-1	rio'l'	556.	688.8	862.4	588.	436.4	376.	368.	277.6	304.	276.	316.	292.	643.	351.	240.	254.	.986	316.	252.	.528	000	254	251.	308.	274.	217.	.346.	217.	252.	:
	Odor		None	9.9	"	9.9	* 9	• •	4	;	9,9	9.1	:	:	•	* >	. 9	9.	:	:	9 9	9,9	9 .	9 9	3	9 9	3.9	7.9	. ,	;	"	
6		Color.		:	Muddy	io.	: :	9.	ie.	∞.	ic.	.05	æ	≈;	89	5.0	-:	9	-	e.j	7:	7	-	 	<u>ن</u> ج	က္	66°.	\$? -	9:	+.	ಣ	•
APPEARANCE.		Sedi- ment.	Much	:	9 9	9.2	Cons.d	9.9	9.9	Little	Cons'd	Little	*	9.9	Consid	* ,	Little	;	:	9.9	;	:	•	4 9	9.9	Cons.d	9.9	9.9	Little	10 10	ud e	
AF		Turb'y.	Much	Decided	*,	;	:	**	y.	:	3,9	3.9	Much	Decided	Much	:	9.9	Decided	Much	7,7	:	:	٠	:	**	9.9	:	:	:	4.	:	
DATE OF	1800	田口	May 13	35	31 June 1			33 33				9 8	*	1119.	,	11,	9.9	Sept.				Oct. 5	+1 ,.	,, 19	96 ,.	Nov. 2	6 ,,	;	Dec.		4 4	:
DAT	1000	ర్	May 10		31	June 8	15	· ·	87	July 5		61 ,,	96 .,	1119. 2		16		Sept. 6		08	27	Oct. 4	" 11	,, 18	55	Nov. 1	000	15	66 ,,	Dec. 13		
No.	[T	Seria	135	55				186	215	248	584	319	356	397	135	167	7.5	584	620	65-1	- 169		173	808		r -		975	1097		-	:

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM ILLINOIS SHORE, ALTON, ILL.

University of Chicago.

Report of EDWIN O. JOHDAN,

No. of Bac.	ret Cubie Centi- meter.	2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 2.1.00 3.1.00 3.1.00 4.5.00
	nosord lo sdv	
	rioquioT' TiA lo	[
athre Pr. C.	Temper of Wate	<u> </u>
	dgiəH ərr//	(2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 + 2 +
30- N A S	səjrajin	<u>ਫ਼ਖ਼ੵਫ਼ੑਫ਼ਫ਼ੑਖ਼੶ਜ਼ਜ਼ਜ਼ਫ਼ਜ਼ਜ਼ਫ਼ਜ਼ਜ਼ਫ਼ੵਜ਼ਜ਼ਫ਼ੑਫ਼ਜ਼ਜ਼ਸ਼ੑਖ਼ਜ਼ੑਜ਼ਜ਼ਜ਼</u>
NITRO- GEN A	sorbitiv	8
NIC OGEN.	solved. Sus- pended	
ORGANIC NITROGEN.	Total, Dis-	
	m sng	844289944448444484444444444444444444444
N AS		9-78-88-28-88-88-88-88-88-88-88-88-88-88-88
NITROGEN AS	T'10'T L'10'T L'10'T L'10'T L'10'E L'1	+888864444588886552888888888888888
N	Freedm.	899998889989989989989989989989989999999
N ED.	ypress. Bysusper	10 51 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
OXYGEN CONSUMED.	By Dis-	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
000	Total.	10 40 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
,91	Chlorin	$\begin{array}{c} w \cup w + y \cup y \cup w + y \cup + w \cup y \cup y \cup y + y \cup y \cup y \cup y \cup y \cup y \cup y$
0.N 10.N.	pepqeq Sus-	######################################
RESIDUE ON EVAPORATION.	Dis-	8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
RE Eva	Total.	######################################
	Odor	None of the state
	('olor.	N
APPEARANCE.	Sedi- ment.	Much Cons'd Little Cons'd Little Cons'd Little Cons'd Little Little Little
1/.	Turb'y.	Much Decided Much Much Nuch Since Since Si
G C 33	1899 1899 Collect Examition.	20 May 13 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
DATE OF	1839 Collection.	Nay
70.	Intros	三記を記述を受ける場合の名称を記念が必然を見ることです。 これを表示を表示する。 これを表示を表示する。 これを表示を表示する。 これを表示を表示する。 これを表示を表示する。 これを表示を表示する。 これを表示を表示する。 これを表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を表示。 これを表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を表示を

TABLE 105.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, ALTON, ILL.

1%	
4	
0	
20	
-	
0	
-	
~	
0	
20	
IN	
5	
-	
~	
-	
Carrie	
0	
_	
E	
$\overline{}$	
0	
-	
(1)	
20	
1	

, in	Chicago.
ORDAN	of
.10	ity
o.	niversi
KDWIN	Univ
10	
ort	

No. of Isac.	per Cubic Centi- meter.	13.750 27.500 22.500 22.500 22.500 22.500 22.500 3.000 22.500 23.000 23.000 25.000
	Presend Abs. of	
	Tennper TiA to	
ature D., Te	Temper	
	Helgh Mate	:
RO-	Setrativ	- 18 0 18 0 4 14 14 15 15 15 16 0 0 0 0 15 15 15 15 15 15 15 15 15 15 15 15 15
NITRO-GEN A	Nitrites	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0
IC FEN.	pepued sus-	
ORGANIC NITROGEN.	Dis- solved.	
5Z	Total.	
	a p,pud -sns	88 98 98 98 98 98 98 98 98 98 98 98 98 9
EN AS	Pioniminoid Dis-sid	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
NITROGEN AS AMMONIA.	T'10'T	85 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
	FreeAm- monia.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0
N ED.	Bysuspd Matter	でいる での 4
OXYGEN CONSUMED.	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Cox	Total.	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
.ən	Chlorin	$\begin{array}{lll} \text{usy} = & \text{usy} \text{usy} \text{usy} + + \text{usr} \text{usy} + \infty \cap \infty \text{usy} \text{usy} \\ \text{usy} = & \text{usy} \text{usy} \\ \text{usy} = & \text{usy} \\ \text{usy} = - & \text{usy} \\ \text{usy} =$
on.	pepued -sus	266.000
RESIDUE ON EVAPORATION	Dis.	88888888888888888888888888888888888888
RES	Total.	2011
	Odor	None .
	Color.	:: : : : : : : : : : : : : : : : : : :
A PPEARANCE.	Sedi. ment.	Much Cons'd Little Cons'd Little Cons'd Little Cons'd Little Cons'd Little Little Little Little Little
AP	Turb'y.	Much Decided Much Decided
OF	1899 Exami- nation.	28. 28. 28. 28. 28. 28. 28. 28. 28. 28.
DATE OF	1899 Collection.	May 10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
No.	Serial	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

TABLE 106.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM MISSOURI SHORE, ALTON, ILL.

Report of EDWIN O. JORDAN, University of Chicago.

No. of Bac.	per Cubic	Centi- meter.		6.500	10,750	95,500	19,500	15.(HH)	9,500	9,000	8,500	9,500	5,000	000.0	3,000		200	£		3.300	3,900	1.650	1.400		1.600	3,700		(C) (c)	6.100	30x	6.300
re or Colf.				:		:						:							:	•		•	•	:							:
ornite D	riet Tiv			16.	.30°.	65	ري اي	÷;	35	\$ 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	300	3.	34.	2; 3;	\$ 1.00 \$ 2.00 \$ 3.00 \$.30	ر د د د د د د د د د د د د د د د د د د د	65	•	.;; ;;	19.	16.		30.	6.	10	01	23	3	-	0.
ernte r, C.				: ::	- 	- - - - - -	16.	2.	5.7.	.36.	30	∞ (2)	30.	30	26.	26.	3.	 S:	24	30.	∞ <u>`</u>	15.	16.	<u>x</u>	<u>∞</u>	1	6	10.	œ	-	0.
	ighi Sate			17.5	16.5	16.8	13	14.4	13.5	15.8	15.8	10.00	c.	ر: دن	·	رة زن	-: :::	Ø. €1	3.6	÷	90 90	∞. ≎ì	?? ??	2.1	6: -	67	3.9	5.0	00 00		
NITRO- GEN AS	səm	NITE		9.	16. 16.	£.	o:	7	ıς	7	÷.	e }	.55	0.	0.	с. —	0.	.0.	<u></u>	:0:	0.	<u>:</u>	(원):	ς.	3.	0.	-	33	10	=	
	<u> </u>	Mitr	.00		012	3	3.	7 0.	.00	.00 .	100.	.002	0	0.	0:	c:	0.	0.	0.	100.	9.	(X)	C	.007	0.	. COD.	300.	.003	0.	100.	1001
ORGANIC NITROGEN.	·s	vios ng neu	:	:	:	:		:	:	:	:	:				:			:	:	:	:	:	:	:				:	:	
ORGA	-s	IO		:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:
	1	o'bnq 10T	:	:	: :	9	: ∞	ः :	े देश	:	:		÷1	ু: ু))	: ?}	:	. 9	: e)	:	:	: œ	:	: 2)	×.	: «	:	:	:	:	: :
202	An .	sng	_	20.0	_							_	_	_	_			_			_	_	_	_						80.	
TROGEN A	Albuminoid	sid 'vios		.53	304	30	57	. 208	1.2.	23	•								_		_	_								37	-
NITROGEN AMMONI	Alb	'10T	<u>.</u>	7.7	1.36	ટ <u>ેં</u>	€	89.	.216	.464		887.	.392	œ.	.344	.376	3963.	.376	. 1.51	<u>∞</u>	36.8	3000	.336	368	**	355	.36	.336	₹	.301	. 153
		Free.	.03	80.	.co.	.058	180	.056	.078	.073	<u>. 108</u>	£	390.	10.	ō.	90.	.013	0.18	030		.026	$\frac{1}{\infty}$	3	.038	$\frac{\infty}{0}$	110.	.045	.053	370.	.011	100.
ED.	pds 1er	Bysu		15.3													0.	9.	1.6	00 01	-	5.	9.1	7	7	- 13	03	\$? ·	33		÷ la
OXYGEN		RA D	10.	9.6	6	∞	<u>.</u>	ci.	re.	6.	9	100	10	10	:C:	10		9		}~	∞	{ ~	L	œ.						∞ ∞	
000	al.	10'I'	-	56. E.	3	?? 6: 6:	χ. 	17.9	15.6	16.	17.5	13.6	∞ ∞	∞ 1G	∞ i~	6.	G !	6.9	7.6	9.6	. 69.	9.1	9.6	9.4	7.6	φ α	9.6	-	<u>x</u>	=	∞ 3.
es.	loriı	СИО	1.1	7:	<u>~</u>	÷;	9.	- 1	?}	-	-: -:	33	3.6	3.8	00	2.6	3.4	3.4			٠; خ				 or.		03	6)		?? ??	
ON.	pər	ns ns	419.	1133.	1349.6	77.9.6	393.5	000 0000 0000 0000 0000 0000 0000 0000 0000	326.	181.6	461.6	151	138.	56.	116.	17.	50.	-1	125.	-0	×.		:96:	55	o€ -}	3,5	Ē.	8:	334	 	34.
RESIDUE ON EVAPORATION		alos i(I	170.	316.										<u>%</u>		140.	181	oc.	89	200	55.	- -	200		176.	166.	155	146.	155.	150.	167.
RES	J.Is	10T	586	1118.	7	55.00	536.	156.	476.4	321.6	6.55.6	3390.	308	. XX	300.	317.	211.	6.00	3500	- - - -	240.	105	 -:	0000	: -			236.	136	3	108
- Administrative	Odor		None	9.	:	•		9 10	9 9	• •	0 9	10	* 9	* 9	;	•	:	9 9	:	. 9	:	:		• 9	:	9 4	• •	:	:	4	**
		Color.			ः	9.	wit.	<u>ဗ</u>	ıĠ	o.	ıc	ಟ	ic.	ଟଡ଼	**	÷	8.	0.	-	E	ची.	rc.	***************************************		: ::	6	2	:S:	9.	-	7.
APPEARANCE.	Sedi.	ment.	Much	0.9		9 9	9 9	p.suo()	9 9	7.7	Much	Little	p.suo.)	Little	7 9	p,suo,)	Little	2.7	9 9	7 9	9 9	Cons.d	Little	:	* •	Cons.d	9 9	,,	Little	:	9.9
AP		Turb y.	Much	Decided	•	:		• •	**	:	:	;	Much	:	:	*	. ,	Decided	Much	99	:	;	;	:	**	* 9	:	9.9	:	**	:
	6681	Exami- nation.	r 13	55	le 1	C:	20	33		1,0		8	67	00	10	-	<u></u>	t. 7	+	2.	œ		7	1:0	36	≎1 .:	10	16		1000 1000 1000	33
DATE OF			May	:	Jun	:	:	:		July		:	* 9	ZIIV 3	:	:	:	2 > 6 11	:	:	:	Oct.	:	:	:	VOV.	:	9.9	Dec.		8 7
DAT	6681	collec- tion.	May 10		:	une 8	:	(F)	· :	July 5	27	: 19	96	Aug. 2	6 .,	9 ,.	(PC +1)	Sept. 6		÷	:	Oct. 4	= :	× :	:: :2	Nov. 1	30	9	58	.:	077
No.	Ini:				£		11:0	2×-	(C)			3.5.	859	100 A	135	170			-9		697	~				-	0%6	975	1030	-	11933

STREAMS EXAMINATION-SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM MISSOURI SHORE, ALTON, ILL.

Report of	Report of EDWIN O. JORDAN,	University of Chicago
Report	of	
	Report	

No. of Bac.	per Cubic Centi- meter.	11.12.000
	Presence App. of	
ernta .O.,	Temperative To	
	Temper otavi lo	~
	dgi9H ots7/	<u>ເລື້ອກາຈະແຕ່ເຮັນ</u> ພາການພາພານພາຍພາພ ເຂົ້ອກາຈະແຕ່ເຮັນພາການພາພາຍພາພ ເຂົ້ອກາຈະແຕ່ເຮັນພາການພາພາພາພາພາພາພາພາພາພາພາພາພາພາພາພາພາ
0. A S	Nitrates	: 16 16 78 0 + 10 10 16 10 10 00 00 00 00 00 00 00 00 00 00 00
NITRO-GEN A	sətirtiN	
EN.	pəpuəd Sus-	
ORGANIC NITROGEN.	Dis.	
62 —	Total.	
	ens budy	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
TROGEN AS	Torr Dis-siu	24 - 25 - 25 - 25 - 25 - 25 - 25 - 25 -
NITROGEN	I'10T	2111 21128887 85584888888888888888888888888888888
	FreeAm- monia.	00000000000000000000000000000000000000
ED.	Bysuspd	ರಾಪರ್ವಾಧಯ 400 ಭಜ್ಞಾಣದ
OXYGEN	By Dis- solved	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
Cox	Total.	యటిక్రిగ్రాఖలన్యలనునునిగా అవారలలను అలకు చెక్కా మంటలు ఆమార్థ్ కుగ్కాశంలో అశ్యనాగా కాకు తను
.ən	Chlorin	က — — — ေ တွင္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္တစ္
ON.	pəpuəd -sug	868548488888888888888888888888888888888
RESIDUE ON EVAPORATION	Dis- solved.	842556668265555777222225588874877333
RES EVAF	.fetoT	170 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	Odor	None
	Color.	— ကန်က်လုံလုံလုံလုံလုံလုံလုံလုံလုံလုံလုံလုံလုံလ
APPEARANCE	Sedi. ment.	Much Cons'd Cons'd Little Cons'd Little Cons'd Little Cons'd Little Cons'd Little Cons'd Little
AI	Turb'y.	Much Decided
S OF	1899 1899 Collee- Exami- tion. nation.	22. 33. June 1. 25. 35. 37. 37. 37. 37. 37. 37. 37. 37. 37. 37
DATE OF	1899 Collee- tion.	May 10 Ma
.o.V	Serial	8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. .

Source of Water-Mississippi River, 400 Yards prom Illinois Shore, at Chain of Rocks, Punping Station St. Louis Water Works.

ORDAN,	of Chicago.
-	
DWIN O.	'niversity
OI E	س
Keport	

No. of Bac.	per Cubic Centi- meter,	1000 25 1 1 2 2 1 1 2 2 2 2 2 2 2 2 2 2 2 2
	Presen lo .sd A	
	riemper niz lo	######################################
	Tennper Diraki	: : : : : : : : : : : : : : : : : : :
	Heigh Unite	: : : : : : : : : : : : : : : : : : :
NITRO- GEN AS	Sitrates	i in the state of
	Nitrites	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
ORGANIC NITROGEN.	solved.	
ORGA	Total.	
	n nud	82522222222222222222222222222222222222
A. A.	b'vlos -sug	
ITROGEN AS	Albuminoic Dis- Dis-	2
NIT.	E 1.10.L	
	FreeAm- monta.	7-4-4-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-6-
EN MED.	BySuspd	
ONYGEN CONSUMED.	sid val	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
-	Trotal	$\mp 478778888888888848998444999999999999999$
16.	. ——— Поти	4 4 6 1 5 10 10 4 6 10 10 10 10 10 10 10 10 10 10 10 10 10
0.N 10.N.	popuad -sus	28 28 28 28 28 28 28 28 28 28 28 28 28 2
RESIDUE ON EVAPORATION	Dis-	
REVA	Total.	F
	Odor	NON
	Color.	
ř.	0:5	
APPEARANCE.	Sedi- ment.	Much Cons'd Much Cons'd
1.	Turb'y.	Decided Nueh Nueh
) F	1809 1899 Tollee-Exami- tion, nation.	
DATE OF	c. Ex	1
D,	1509 Collection.	May May May Mune
No.	Serial	

TABLE 109.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, AT CHAIN OF ROCKS, PUNITING STATION ST. LOUIS WATER WORKS.

Report of Edwin O. Jordan, University of Chicago.

No. of Isac.	per Cubic Centi: meter.	16.100 16.100 18.100 18.100 18.100 18.100 19.100	006.4
	Presend Abs. of		
	Temper		20.
ature er, C.	Temper	::: ::::::::::::::::::::::::::::::::::	00
t of	dgiəH ətsW	:: 1889 489 888 5 + 11 mg 1 + 6 6 10 4 10 10 10 10 10 10 10 10 10 10 10 10 10	9.7
RO- N AS	Setrativ	: :: :: :: :: :: :: :: :: :: :: :: :: :	<u> </u>
NITRO- GEN A	Setititiv	00000000000000000000000000000000000000	90.
GEN.	bended -sus-		
ORGANIC NITROGEN.	Dis-		: : :
	Total.	2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	2.96
As.	Sus.		
NITHOGEN AS	Albuminoid Silvios	######################################	•
NITH	[1,10,L	8-4558-4 Exercise & 888 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	
	Freedin- monta.	80.00000000000000000000000000000000000	•
EN MED.	Bysuspd	$\infty \infty$ ω	
ONYGEN CONSUMED.	-sid val	∞ 5- ∞ 0 5- 7- TC ○ 20 4 61 4 4 TC 4 TC 6 TC TC 4 4 4 60 T	0 vc
	Total.	######################################	
116.	Chlorin		<u> </u>
ON TON.	Pended Sus-	2105 2048 2048 20418 20418 20418 20418 20418 20418 20418 20419 20418 204	342
RESIDUE ON EVAPORATION	solved.	88.50.00.00.00.00.00.00.00.00.00.00.00.00.	276.
RENEVA	Total.	2330 1750 1 6 1750 1 6 1750 1 6 1750 1 7 1760 1	618.
	Odor	None	:
	-	: : : : : : : : : : : : : : : : : : :	: : ରାଡ଼
μĵ	Color		
APPEARANCE.	Sedi- ment.	Much "" Cons'd Much Little Cons'd	Much
AI	Turb'y.	Decided Much	
OF	1899 Exami- nation.	May. Jume 152 Sept. 138 Se	16
DATE OF	1899 1899 Collec- Examition.	84444444444444444444	Dec. 15
.o.Z	Serial		1033 1104 D

TABLE 110.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, AT CHAIN OF ROCKS, INLET TOWER, ST. LOUIS WATER WORKS.

University of Chicago.

Report of EDWIN O. JORDAN.

No. of Bac.	Cubic	Centi- meter.	•		11,100	14.300	69.000	24.000	97.00	000,02	12.0(N)	8.500	13.000	5.150		6.550	4.450	3,000	11.000	1,500	8.900	8.900	1.800	200.20	5,100	1 800	OUF 6	8.500	
ce or		Pres.																	•	•		•	•	•					:
		r jo ituə _L ,		:		27.	چې	33	20.	35.	15	36.	37.	.66	35.	33	32.	38	- 100	8	200	.18	∞.	19.	eş	4	<u>} -</u>		:
ature E. C.		TemiT			:	22.5	54.5	.96	÷ €	55.	26.	29.	.666	.98		5.00		29.	55	19.	16.	15.	17.	17.	G	9.	00	0	
10 1 (10	gh gh		:		25	20.7	&. €.	8.0%	20.4	633	20.6	16.6	14.3	19.9	13.9	ž C:	7.5	8.6	8.6	9.9	ۍ س	4.5	30	٥٥.	ත. හ	6.5	10	4.6	:
RO- N AS	ธอ	Mirat			£.	£.	_:	-	10	9.	.35	.45	.45	ું	坦	.45	3;	. 15	G3	ಚ	10		ુ	0.	£.	ુ:	7.0	. 15	:
NITRO- GEN A	sə	nanz			.0075	.002	90.	900	0.	300.	0.	.00:4	0:	0.	300:	.0035	.01	0.	900:	.0034	900.	0.	0.	0.	300.	0.	0	300.	:
NIC OGEN.	_	sus		:	:	:	:	:		:	:	:		:	:		:	:	:	:	:	•	:	:	:	:			:
ORGANIC NITROGEN.	-	si(I 97/los	:	:	:	:	:				:		:	:		:						:			:			:	:
	1 . 1	b'baq glo'f'	6	95	176	800	632	224	824	792	82	808	7.3	684	744	: 50	: 38	28	880	336	256	252	214	172	221	11%	136	39.5	:
A8	noid Am	b'vios	<u> </u>			_	_	_	_	_			_							_		_	_					.018	:
NITROGEN A AMMONIA.	Albuminoid	-sid																						_				0. 11.	•
NIN A		I'10T	1							_								_	_	_				_	_				:
	-tu	Assid	5.														2 .03	₩.	00. 12	<u>2</u>	99.	10.	0.	30.	.G	<u></u>	30.	301.	:
OXYGEN CONSUMED.	pd	Bysus		Ξ	₩.	18.	31.	21.	≃:	33	.4 5.8	53	33	<u></u>	53				6 3.5	33	52 53						31	6 3 3	:
OXYC		sid va	∞	10	3 10	9.14	10	7	10	300	25	6 -1		\$	9	_ 	23	53 CI	- - -	-	-0	35	6 1	00 00	6 1	00	÷ ∞	~	:
	1	R10T	-		3 31.	-			-										200					-					:
16,	nin	Сріо	6.	4	ш.	ಣ	ಣ	9	ci	90	4	- Ji	10	44	<u>ن</u>	-3	} -	ξ	00	6.	10.	10.		~	12.	50	13.	15.	
ON TON.		opuəd sus	3260.	2730	2907.2	2866.4	1100.	3009.6	2196 8	3085.6	2624.	2381.	2516.	2080	1877.	1536.	:17:	889	×6.1	661.	571.	513.	386.	386.	385	156.	358	382.	:
RESIDUE ON EVAPORATION		sid earlos	210.	240.	201.8	221.6	256.	216.4	232	200.	.30€.	195	311.	158.	198	138	301.	205.	216.	0888	25. 25.1	255	27%	2306.				294	
RE] . [nio'l'	3470.	3030.	3112.	30.88	4356.	3456.	2728.8	3285.6	2821	2576.	2760.	2278.	5002	1734.	.8/	1094	1110.	894	825°	798.	662.	(38.5)	.699	420.	616.	676.	:
	Odor		None	• 9	9.9	•	9.4	9 9	9.9	, ,	9 9	9.5	9.9	• •	9 9	9 9	1 9	7.9	10	9 9	11	9.0	9 9	:	. ,	9.9	9 9	9 9	
		Color.				6)	≎}.	under	-	-	93.	**:	0	?)	₩.	0.	0.	18.			એ	ા	??	23	?≀	. 15	10		
APPEARANCE		Sedi- ment.	Much	:	:	:	**	9,9	9 .	7.7	**	3.9	10	:	9.7	7.9	*,	:	;	7	:	*	Cons.d	. 9	Little	Cons'd	4 9	Much	
A.P.		Turb'y.	Decided	:	;	*	*	9 11	*.	7,	Much	10	*	*	9.9	9 9	* 4	. 9		9.9	70	,,	9 9	:	:	9 .	:	:	
E4	00%	Exami- nation.	- A	15	300	ne 1	16	33	30				7	62	$\frac{\infty}{}$				10	553			201			11	-	16	:
DATE OF	-	- Ex nat	8 May		:	1 June	**	;	1. 6	13 July		20 2	3 Aug.	., 0	:	: -	1 Sept		**	**	; x	5 Oct	,, ()	3 9	2 Nov	,,	Dec		
DA	1800	Collec- Exami-	Apr. 2	May 1		June	.,	::		July 1		2	Aug.	1	- ;	::		1500	1	e s	= = = = = = = = = = = = = = = = = = = =	Jet.	07	26	Nov.	**	Jr,	Dec 15	
Zo.	11:	iroz	=	31	6:3	94	100	193				373	101	113	176	515	560	591.	635	(7,7)	100	743 (No.X	850	501	555	0.21	_	

TABLE 111.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 400 FEET FROM MISSOURI SHORE, AT CHAIM OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

Report of Edwin O. Jordan,

University of Chicago.

No. of Bac.	per	Centi- meter.			20,700	11,400	57,500	10,500	11,500	33,000	14,000	10,500	16,000	2,700		5,150	8,400	7,300	2,900	7,100	€,300	0,000	2.800	4.400	3,300	7,300	13,500
		Preso Abs. o		:		•		:	•	:		:		:		:	:	:		:	:	:	:	:	-:	•	
		qm9T A 10				27.	28.	35.	.02	83°	27.	36.	37.	65	33	35	33	25.	ි. ද	18.	23	18.	19.	c;	+		٠.
ature r, C.	er ete	qm9T 3W lo	:	:		22.5	24.5	.98	25	27.	56.	20 30	83	8	55	37.	27.	53	19.	16.	15.	17.	1-	6	6	œ	0.
tot .19	q2	təH W		:	33	23. T	24.8	30.8	20.4	23.9	9.0%	16.6	14.3	13.9	13.9	9.7) () ()	9.9	9.9	ۍ ښ	4.5	00 00	က	3.9	6.3	5.5	4.6
RO-	80 	Mitrat		:	.375	٠.	-		ī.	9.	.55	÷	.15	15	.55	2	હ ું	:33	w.	.83		33.	0.	왕.	દ્ય	.15	.15
NITRO- GEN A	sə	Mitrit	:	:	8.	98.	900.	.0062	9.	90.	8.	0.	0.	0.	.0024	013	0.	.003	0.	.00	0.	0.	0:	8	.003	0.	.002
IIC GEN.		sns	:	:	:	:	:	:	:	:		:	:	:	:		:	:	:	:	:	:	:	:	:	:	
ORGANIC NITROGEN.		Dis		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		
OZ	1	T'ota		:	:	:	:	:	:	:	:	:	:	:		:	:	:	~	:		:	:	:	•		
	d Am.	p,pud -sng	1.04	:8: 	1.36	1.488	1.464	1.248	1.14	1.75	1.06	.612	1.06	.764	888.	88.	.56	.38	308.	.256	. 232	.056	.176	1+6.	.076	.248	9.
NITROGEN AS	Albuminoid	Dis-	.16	.14	* **	.112	.176	. 152	.16	.216	.16	808.	80.	.136	. 152	.13	.13	.136	.07:	.104	880.	.184	960.	960.	.061	£.	₹.
NITRO	Albı	I'10'I'	1.3	+:-	∞.	1.6	1.64	+ :-	1.3	1.94	1.33	82.	1.14	6.	1.04	-	89.	.53	. 38	.36	.32	.24	.272	.34	11	3.3	6.4
		FreeA	90:	0.	9	890.	.064	990.	980.	.076	920.	.078	10.	990.	.058	.082	.027	.038	.036	.036	.03	.034	10.	.018	.013	:03	.07
ED.		BySus	15.8	33	18.	17.2	32.5	23.5	13.4	15.4	υ. ∞	13.8	16.9	12.9	15.2	14.4	7. 7	4.6	£.5	30.00	3.6	+	3.0	3.2	1.3	4.2	3.4
OXYGEN		sid ya solve	5.4	4.	∞ π	16.7	.9	6.4	5.1	3.4	3.1	∞ ∞	1.4	4.6	63 63	4.6	-	4.3		6.5	7	د	3.1		4.2	2.6	3.
Con	1.	RIOT'	21.2	16.3		33.9				18.8						19.	↑ .	8.9	7.6	∞ ∞	· •	7	6.6	7.4	5.	8.9	5.6
ne.	ıiı	СРЈО	6.3	4.6		÷ €;	÷	m	33	+	+	÷		بر	1-		} -	10.	=	10.	13.	16.	14.	<u>c</u>	63	15.7	1
ON.	pa	bend	3325.	3340.	3660.	2089.2	279.2 4097.6	3487.2	2472.2	205.6 3257.7	1817.2	2533.	2741.	2356.	2035.	1.158.	1051.	905.	730.	570.	516.	378.	396.	528.	162.	124.	482.
RESIDUE ON EVAPORATION		Sid Solve		274	210.	233.6	279.3	8.96%	238.8	205.6	200.8	192.	211.	268.	201.	210.	215.	255	262.	276.	272.	290.	304.	278	318.	318.	314.
RE	۱۰.	Tota	3560.	3614.	3900.	2322.8	4376.8	3744.	2700.	3463.3	3018.	2725.	2952.	2624.	2236.	1668.	1266.	1160.	992.	846.	788.	.899	700.	706	480	742.	7.96.
	Odor		None	91	7 7	**	9 4	11	11	**	7,7	7.7	9,9	9.9	9 9	1.7	9.9	9.9	9.9	9.9	9 9	9 9	9.9	9.7	9.9	9.9	77
		Color.			•	લ્યું	ં	7.	7	ıe	ಣ	e0:	0:	દરે.	ં.	0:	0.	-:	0.	. I5	.15	<u>ુ</u>	<u>.1.</u>	ાં	<u>61</u> .	.33	-1
APPEARANCE.		Sedi- ment.	Much	9 7	7.9	9.0	77	. ,,,	9.9	9.9	9 7	9.1	9.7	9 9	7 9	9 9	9.9	9.9	9.9	9 9	3.9	Cons'd	77	Little	Cons'd	9.0	Much
AF		Turb'y.	Decided	**	**	97	;	:	3 9	99	Much	*	7.	;	:	11	91	;	;	:	**	9.9	,,	,,,	"	*,	3
OF	1000	Exami- nation.	May 1	15	128	June 1	" 16	11 23	30	g		98	3 Aug. 4		** 18	3.5 (C)		ii 15	33	68	Oct. 6		160	Nov. 3		Dec. 1	" 16
DATE OF	1000	Jose Collec-E tion.		May 12			122	667	53	July 13 July		: 32	Aug. 3 A	-	17	T 67	31 Sept	Sept. 11	13	288		20	26	Nov. 2	13	30 1	Dec. 15
No.	[T	Seria	10 A	30 N	65	95 Ju	151	194	227		_	37.1			17	516	561			1,00	_	-				10335	

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

Report of EDWIN O. JORDAN,

SOURCE OF WATER-MISSOURI RIVER, FORT BELLEFONTAINE, WEST ALTON, MO.

ort of Edwin O. Jordan, University of Chicago.

No. of Bac.	per Cubic Centi- meter,	8.000		SS.4.	00% e1	5 600	1.600	(S)(C)	9.100	0.06,02	4,000	10,000		200		0.00	0000	6 600	0000	18.200	
	Presenc 10 .sdA		:	:	:													•		• • •	:
	Tempers		. S	31 6	: : : :	€	1.	G1	169	19.	.55	÷	23.	19.	10	\ \ -	-	ıc	0	1	
ornite r, C.	TequeT otsW lo	29.	818	33 8	900	200	<u>x</u>		3	38	19.		2	31	_	9	15	0	6	<u>-</u>	:
	Helghi otaw	13.3		xo s	ر د د د د	9	20.	٠.	20	4.6	£.	¥ -	£.	4	6.7	10	10	7	0	63	:
NITRO- GEN AS	setaull/	1											-							. S.	:
	Nitrites	0.	0.0	30	. E	000	0.	0.	0.	0.	.03	<u>.</u>	0.	300.	300.	0.	0,	00	8	100:	:
NIC GEN.	solved.	:	:	:	:						:			:			-			:	•
ORGANIC NITROGEN.	-siu	:	:	:					•	:		:									:
	Total.		:	: ::		9	??	16	77	3		34		14	.: 80	9	7.	20		10	:
81.	-sus	1 .	-		200	-				_	-	_		_	_			_			:
NITROGEN AS	Pion Pion Pion Pion Pion Pion Pion Pion		128	500	000	. 136	.072	.07.	.032	176	101	.136	- O.	.03	380.	.07	070	.05(0	0	
NITE	[4] ['10'T		889	1.42	536	432	.344	.228	.256	. 288	.264	 	.216	.176	968	. 248	2963	30.4	1961	.136	:
	Freedm- monta.	.058	0.58	0.00	360	800	.020	.038	30	810.	030	850.	.036	980.	30.	035	0.05	056	.012	-	
EN (ED.	Bysuspd			00	21.2				4	4.7	2.4	G?	4.6	03 33	·-	63				9 -	:
OXYGEN CONSUMED.	By Dis-	!	ಕ್ಷಾ ಕ	N C	9 50	600	9		33.33	্য						90					:
၀ ၀	.lato'l	19.9						-1	{	- E		φ. #	x	÷	6.	+	÷	5 6	6.	American Caris	
16.	ilaofil')	5.5	χο υ σ: ι	0 0 - 0	10.6	12.1	13 6	15.3	15.6	13.9	17.4		83 83 80	27.		16.8					:
0.N 10.N.	pepuad-suS-	3280.	2366.	-+++	991	863	700.	364.	563	493.	100	191	388	350	62×	464	45%	803	787	218.	
RESIDUE ON EVAPORATION	-sitt .baylos	240.	198	906	276	275	1986 1986	304	314.	307.	317	310.	301	310.	2063	321.	× × × × × × × × × × × × × × × × × × ×	330.	316.	358	
RE	Toral.	3520.	1361	1905	1270.	138	908	. 66%	200	Q	1356.	771.	795	699	924	-188- -188-	816.	0.38	× 500	909	
	Odor	None	: :	:	:	:	;	:	:	:	7 9	:	;	:	9 9	:	:	*	:	:	
	Color.	e3 3	Sinday.	ج ج	-		.15	ici	<u>- 61</u> .	92	<u>.e</u>	- 2	<u>??</u>	3)	=	35	0.	0	-	Muddy	
APPEARANCE	Sedi- ment.	Much		:	:	:	:	p.suo.)	Much	:	:	:	p.suo.)	Much	:	Consid	Much	:	:	:	
11.	Turb'y.	Much	. :	:	:	Decided	:	:	:	Much	:	:	:	:	:	:	9.9	:	:	:	
OF	1899 1899 Sollee-Exami- tion, nation,	Inly 28	507.	? ×	15	(Z.)	6.	et. 6	10	20	 	Nov. 4	01	:	24	ec. 1	xx:	16	23	S	
DATE OF	1899 Collec- Etion. n	17:	7. 7	r.	-	12	J. (2)	0 0	27	51	56	7.	5.	91	23	30 D	-1	=	12	×.	
	Col tio	S.July	2017.0	101		:	*	Det.	:		:	10%		:	:	:	T Dec.	:	:	:	:
7.07	Infros	33	W. 15	35.50		100	-	110	0 = 1	37	4630	[1]	500	376	INT	57	965	101	1.39	153	

TABLE 113.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 YARDS FROM ILLINOIS SHORE, JEFFERSON BARRACKS, MO.

Report of Edwin O. Johdan, University of Chicago.

No. of Bac.	Per Cubic Centi- meter.		24,100	53.00	1,000	000 H 6	9.00	13,000	15.500	002.8	6.750	3.200		37.38	10,100	20.690	31,100	2.600		3,600	1,000	2.000	11.900
	Presenc				:			:	:									:	:			:	:
	neqmeT att. to				ج وي				:				:	:					:	:			
	Tempers) 2 2 3 3		9 20	200	30.	2. S	G X		30.		<u>:</u> 1-	16.	30.	19.	6.	ъ. Б.	<u>∵</u>	ee.	÷)	ë.
	idgiəH ətsW		\$ \$ \$ \$: : 3;	20.55 10.05	23.55 23.55	19.8	16.2	e: -	13.0		10	က မ က မ	- m		ص ن:ن	33.5	÷	5.4	9.9	∞. →	9.1	.c.
NITRO- GEN AS	Nitrates	0.	 	.75	.65°	. 1	10	12		00°	÷:	.05 650		; C		0.05	0.	₩.	ः	.co.	€.	.35	₹.
	Nitrites	800:	.005	50.	0.0	20.	.003	0.	0.	000	900	900.	88	38	38	0.	ο.	900.	800.	30.	800.	10.	300
ORGANIC NITROGEN.	solved. Sus- bended		:		:			:	:	:		:	:	:			:	:	:	:	:	:	:
ORG/ NITR	Total.		:	: :	:	: :	:	:	:	:	: :	:	:	:	: :		:	:	:	:	:	:	:
	p,pud -sng	∞		. 13	.536	038	.792	.524	.576		288	.336	30 0 0 0 0 0	0.000	935	166	. 956	. 288	.16	. 184	· 참:	80.	. 208
N AS		oj oj					_				973	104	104	1840	168	152	144	113	176	248	63	216	400
ITROGEN AS	Albuminol Piori Divios	- 0.5		_	_	_	_	_	_	:			.352										
7.	FreeAm- monia.	800					_			046		001	013	024	032	018	33	062	036	054	0.94	043	013
r.	Bysuspd Matter.	11.6	ಣ ರ	ক	70	2 9	. 9	<u>.</u> د ويد	<u>ာ</u>	· -	6	9	-	· ·								_	
OXYGEN CONSUMED.	By Dis- solved	8.30	1-		?? -	7 52	63	030	00 c	300		∞	9.0								∞ 		
Con	Total.	17.5	85.00 80.00	57.50	17.6	8.9	17.8	16.7	16.4	0 00	16.9	9.4	7°.6	00 oc	7:00	9.	9.3	8.3	6.5	14.	9.1	6	9.4
.en	Chlorin	00 co	्रे~ ©] क च	3.55	න ය දෙන	100	5.7	∞ 			6.1	6.9	69.	9.0	تر. ن	6.4	≎} ∞	6.7	9.9	οο - 1	©} ∞	6.8	6
ON ION.	papuad -sug	1946. 1581.	2490. 3150.8	2613.2	1618.4	2513.2	1803.2	15-13.	1829.	971	664	735.	362.	130	87.	. . #	150.	124.	136.	73.	16.	23	7.6
RESIDUE ON EVAPORATION	Dis- solved.	209. 181.	218. 2490.	266.8	202.	201.6	192.8	207.	217	180.	196.	207.	192	193		208	182.	196.	170.	184	214.	202	217.
RE	Total.	2155. 1765.	2708.	2880.	1850.4	2714.8	1996.	1750.	2046.	1151.	860.	935.	554.	322	277.	303.	322	320.	396.	257.	230.	284.	296.
	Odor	None	9.9	,,,	9 9 9	99	9.9	3 9	; ;	9.9	"	;	; ;	,,	3.9	9.7	3.7	:	9.9	3 9	90	9.7	9 •
	Color.		40	9.	4: -	ini	eci.	တဲ့ ေ	٠. 	ર જ	0	Muddy	-05	- 7	. 00	58	13.	ः	.25	ıc.	च	ಲ್ಟ್	. 3
APPEARANCE.	Sedi-	Much	::	;	, ,		-	93	; ;	,,	9.9	*	11 110-101	n silo	Little	9.9	3.9	Cons.d	ittle	9 9	9.9	7 9	3 9
PPEA						~							2	3	1			0					
	Turb'y.	Much	Much		Dobidod	:	Much	9,9	: ;	7.	"	3,	7.7	3.9	9.7	93	9,9	77	91	,,	7.7	"	7.7
OF	1899 Examination.	May 3	6 June 7	57	30	15	÷.		Aug	300	196			300	ct. 7	15	0~ ,,	Nov. 4	" 13		Dec. 8	15	233
DATE OF	1899 Collec- E tion.	~ =	- the	- es	29	\$ T	<u></u>	53	ಣ <u>∓</u>	11-	: 33	31 Sept			9		19	භ		23	-	- T-	
		14 May 38	10 June	;	9299 cc	299	*	366	407 Aug	180	518	21	598 Sept	2080	oll Oct.	785	6	905 Nov	12	,, 10	1062 Dec	35	33
No.	Serial	""	====	131	S is	র না	334	33	¥;	7	27.5	551	550	2 5	- 10	1-	000	8	ਨ	1001	106	1092	1133

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-MISSISSIPPI RIVER, EAST OF MIDSTREAM, JEFFERSON BARRACKS, MO.

	Č
;	8
JOI:DAN.	Phicarro
	_
÷	ĵ.
<u>.</u>	+
EDWIN	University
01	-
eport	
4	

No. of Bac.	Cubic	Centi- meter.	22.650	35,000	26.500	18.500	12.000	34,000	17.000		7,500	12.100		6,800	008,2	42,000	8,400	10,600		14,000	0.000	6.600	⊋ Z ;		4.500	1 700	1.700	6.000
10 99 ('0]].		Pres.			:							:	:			:		:										
ature G.		TempT	:			()2	:			:		:	:		:	:			:				:	:				
		TuoT W lo	.96	.98	800	26.	36.	28.	œ;	88	??	66	65	30.	30.	30.	19.	17.	16.	 03	19.	19.	6	6.	<u>22</u>	6.5	33	e.
10 J	ця 918	Hei W	23.5	22.8	33	20.5	22.9	33.6	19.8	16.2	14.3	14.4	13.9	∵ . ∞	7.0	6.3	6.7	5.3		ಕ್ಕಾರ್	¢? co		***	5.4	6.8	oc.	4.6	ان دن
NITRO- GEN AS	sə	Nitrat		.8	1	pant .	pand	rci	£.	.45	£.	.36	e.	rci	ಬ	.G.	≎?	<u></u>	ું.	88.	0.	0.	.095	:3:	<u>(C)</u>	e.	e:	M
		MITTE	00.	30:	900:	0.	800.	0:	0.	0:	0.	.055	900:	0.	1005	.003	.000	.005	.003	0:	0:	.003	.005	800.	.020	.003	010.	.012
ORGANIC NITROGEN.		eng Sus- eagles	:	:	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	
ORG. NITE	_	gioT'		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	•	:	:		•	•	:
	Am	p,pud ·sns	1.568	1.624	1.208	.873	:	1.232	.661	.632	. 595	.83	.504	.56	87.	288	16.	. 216	.96	318:	. 192	. 199	.181	.272	-	181	::	-
EN AS NIA.		D'vios	192	.176	192	808	:	208	176	891	168	24	183	16	16	51	ि	181	91.	168	85	128	136	168	21	256	176	91.
NITROGEN AS AMMONIA.	monta.		.76	30	7	.08	-		_					_		0.5			.43			_		_	_		200	
7.	FreeAm- monta.			.054	part	pant		08	 8	07.2	.e	044	058		<u></u>	01 #	0.55	035	 	- 810	970	1770	0.11	920	00	160	044	100
SD.	a	Matte	9.	32.3	-	9.	6	_	-	10	6	23	-	00	9	20	_	೧೯			6.	-	\$5					
OXYGEN			44	5.00	9.4	5.4	9.4	50 63	÷.	£ .	2.3	10	4.6.1	5.4	∞. •	∞. ∵	7.5	7	∞ 0;	∞ ≀~	6.7	7.1	·-	σċ	(~ 00	0C	οc	7.1
000	. [втот	65	38.1		16.		\$7 \$7	17.9	16.7	17.9	16.2	<u>.</u> ;	17.5	9.1	7.6	9.5	9.7	с: С:	-6 -	7.6	ලා වැ	0.00	?? ∞	12.8	-6:	<i>σ</i> . ∞	oc :
ıe.	LĮJ!	Срјо	4	ugi										6.1	9	9	6.6	9	5.6		∞		{-	īĊ.		эc		C.
ON.		bend Sus	2620.	3367.2	3975.2	1921.6	208.8 2309.6	2432.	2016.	1614.	2204.	12:28	1118.	855	811.	424.	.386	130	161.	300	1.0%	.93	168.	154.	126.	. 99	76.	202.
RESIDUE ON EVAPORATION	.fi	SOLVE	225.6	255.2	364	216.8	208.8	181.6	192	205.	232.	152.	1	194	201	306.	191.	300	- 182	308.	194	:00:	<u>2</u>	170.	180	202.	2003.	201
RES EVAL	ŀ	giol'	2845.6	3622.4	3239.2	2138. #	518.4		2208.					1049.					312.									
	Odor		None		3 9	7 9	9 9	9 9		0 9	9 0	9 4	9.9	9.9	9 7	9.9	, ,	9.9	.,	9 9	9 9	9 9	9 .	9.	9.9	9.9	3 9	:
	Color.			??	THE .	चं.	7.	ıc.	œ.	~;	0.	સં	ಣ	0.	0.	.E).	Τ.	7	ec.	ಬ್	.25	. I.	৽	.es.	.e.	**	જ	**
APPEARANCE.	Sedi- ment.		Much	**	9.9	:	:		**	:	9.7	7,9	:	**	. ,	:	Cons.d	:	Little	Cons'd	Little	Cons.d	:	Little	:	2 9	*	p.suo.)
AP	Turb's		Decided	Much	**	Decided		99	Much	. ,	:	9.0		9.	*	7 9	:	*	99	9,9	7,9	**	9,9	9.9	**	;	**	***
E.	800	Exami- nation.	ne 7	1-	55	30			33	₹1 21				56		6						22	V. 4	13	C.			£5
DATE OF		p)	E.June	91	200	5000	6 July	,, 1	21 "		S.Aug.	,, .[]	.,	35	31,Sept.	30	33	*	6 Oct.	: 2	19 61	36 44	3 Nov.	11 **	33	7 Dec	*	21
DA	DATI 1899 Collec- tion.		June		9.0	:	July	9 9	1912	**	1112	:	7.	6 10	3	Sept.	67	33	Oct.		9 4	63	NOV.		67	Dec.	1 ,,	2)
Sortin No.				155	197	230	261	3/11)	335	367	1	× +	W.	518	552		67.1	(30)	752		953	863		916	()()5			134

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of EDWIN O. JORDAN, University of Chicago.

No. of Isac.	per Cubic Centi- meter.	181.000 181
	Presenc	
ature,	Temper	50.
ature er, C,	Temperature T	
	Heigh Wate	: 888888888888888888888888888888888888
RO- N AS	Nitrates	○ 1- 1- 1- 1- 1- 1- 1- 1- 1- 1- 1- 1- 1-
NITRO- GEN A	Nitrites	000 000 000 000 000 000 000 000 000 00
IC FEN.	pəpuəd	
ORGANIC NITROGEN.	Dis-	
0%	Total.	
92	p,pud -sng	1.04 1.288 1
NITROGEN AS AMMONIA.	Pioniminoid b'vios	6.25.48.48.48.48.48.48.88.88.88.88.88.88.88.
NITRO	[A 1'10'T	:: 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1
	FreeAm- monia.	0.05 0.05
EN (ED).	BySuspd	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN CONSUMED.	By Dis-	ი⊱იი+410+44440000+000000000000000000000000
000	Total.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
re.	Chlorin	4444446916161616161616161616161616161616
ON ION.	pepuəd sns-	230. 250.
RESIDUE ON EVAPORATION.	Dis-	28.50 28.50
RE	Total.	28.20 28
	Odor	e do not not not not not not not not not no
	Color.	
APPEARANCE.	Sedi- ment.	Much Cons'd Much Cons'd Little Cons'd Little
APS	Turb'y.	Much Decided Much Decided
OF	1899 Exami- nation.	May 3 June 7 1 1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
DATE OF	1899 Collec- E tion.	21 181 38 5 5 5 8 8 8 8 8 11 1 8 1 1 1 8 1 1 1 1
.o.V	1	15 May 112 June 156 " 231 July 331 " 336 " 336 " 449 " 484 " 553 " 553 " 710 " 710 " 710 " 710 " 710 " 710 " 710 " 710 " 710 "

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-MISSISSIPPI RIVER, WEST OF MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of Edwin O. Johdan, University of Chicago.

No. of Bac.	Cubic	Centi- meter.	26.250	18,000	12,000	40,000	21,500	31.000	27.00	9,500	34,000	13,000		9,(50	33.900	37.800	91.600	9.700		14,700	00x.17	99,100	19,000		19,400	3.700	3.200	1,800
ee or Coll.		Pres.													:			:	:			:			:			•
эппв Э.О.;			:			.30°.				9:											:				•			:
ature er, C.		Trent W to	26.	.00°	٠ ج ج	.32.	26.	×	æ,	(%	23	.055	.005	E	.30	30.	19.		16.		<u>s.</u>	15	<u>S.</u>	Ġ.	23	···	÷	0
To t			23.5	∞ ?}	÷	20.5	6.53	33.6	8.6 <u>1</u>	16.2	1.3	H. H.	13.9	₹ %	10.	6.3	6.7	10		90	??		-				9.1	
RO- N AS	sə	Mitrat	1.1	16:	9.	19		1.	6:	ಖ	£.	c:	ıĢ	? -	?}	-:	100	5	5	CC.	??	3)	57	~:	9.	e)	(6).	325
NITRO- GEN A		MININ	10.	920.	8.	0.	.003	0.	1000	0.	<u> </u>	0.	.00°	9.	000	100	9.	300.	300.	(90)	.00°	300.	301.	9.	900:	200°	000	<u>\$</u>
NIC GEN.		sns	:	:	:	:		:	:		:	:	:		:	:	:	:	:		:	:	:	:	:	:	:	:
ORGANIC NITROGEN.	-	SIU SUVE			:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	-		:	:	:	:	:	:-
0/		b'baq	ژي :		<u>.</u>	:	:	456	556	36	318	:	56	396	<u>x</u>	30%	304	-:	25	36.8	300		36 1			36	398	
AS	oid Am	-sng	_		_	_		_	_	_			_			_					_		_			_	_	
NITROGEN A	Albuminoid	-sit[_	91.176		_	:	8.224				:						_	_	_	_		_	_	_	·	111	
NITI	Al	['10']		.19		_	•	99.1				:		_				_	_	_	_	_		_	_	_	T.C.	
		Коет Іпош		.062	•			·	·																			
EN MED.	Þd	Bysus	2 31.9	131.4	19.6	13.6	3 14.7	5 22.5	2 5.5	× 13.2	3 16.1	3.14.2	9.13.8	5 14.7	- co	30. 4 10. 4	5 1.4	55	3.0		6)	- ×	5.5	3. 4.5	6 4 2	9.1.6	2.0	
OXYGEN CONSUMED.		sid ya erfoz	6.	8					3.0	÷	2 1.8	30	500				_	5.7		. S.	7.	₹. + €	5 +· C	6	8 6.0	7 1.9	20	63 10
5	.1	rioT'	8	.5.	8	$\frac{1}{2}$	6:	œ; ∞;	<u>∞</u>	$\frac{\infty}{\infty}$	$\frac{1}{\infty}$	-	==	<u>c:</u>	œ	00	6.	c:	α	တ်	?-	6.	(-	œ	5		1-	∞ ~
.ou	na.	old")	-	33	33	33	33	-	10	+.	IG.	10	1-	9	9	1-	0	G.	=	2	C.	=======================================	27	9	=			
ON MON.		sus sus	3273.9	3029.6	3250	5550.X	2564.8	2519.2	2641.2	2470.	2844	2336	1803	1285	934.	690	393	138	40×	<u>*</u>	402	3338		× ()	332.	297	297	388
RESIDUE ON EVAPORATION		sid							198.7	300	214.	SINO.	194.	102	313	201	236.	216.	260	266	100	25K	276.	250.	245.	202	251.	258.
Bya	.1	nto'l'	3532.8	3248.	3181.2	2459.	2770.8	2715.6	2831.	2670.	3058.	2536.	3001.	156.	1146.	500	X 25.00	684.	728.	684	641.	596.	600.	- S	577.	562.	551	646.
	Odor		None	9.9	:	1,	:	9.9	9 9	:	9 9	:	:	:	:	:	:	*	9 9	:	,,	9.9	:	9.9	9 •	9.0	9.7	3
-		Color.	100	^)			with		~	~	_		~		Muddy			٥١.	0)	0)	- 19	10	21	15	÷:	2)	2)	-
E.	L	[00]		_			_		- :			_	**	_	NEW.	_	_	_		_								
APPEARANCE.		Sedi: ment.	Much		•	4.	7.9	;	:	:	:	:	:	:	:	:	p.suo,)	Much	Cons.d	:	:	:	Much	p.suo.)	:	:	:	Much
14.		Turb'y.	Decided	Much	:	Decided	:	:	Much	:	77	:	9 *		* *	:	7.7	Decided	Much	**	:	:	:	:	:	:	:	:
).F	1800	Ton. nation.	1- 51			OK:	ulv 7	1.	::	٠٠٠ ١٠٠٠	15. 4		× :	96	Sept. 1	·:	1.	667)ct. 7	100	(F)	:	10v. +		- 07	S . 3		: 23
Date or		E BE	6.111116	16	33	000	6.111	-	100	-	31112	=	1-	125	31 Se	×	23		-	-07			-			-		
1),	1800	Conec	June	*	7.	:	HII		99	9.7	1112.	:	:		**	Sept.		9.5	Ort.	:	9 9	9.9	1001		:	Dec.	:	**
.0%	[12	Reth	133	157	15.6	000	2633	27.50	1000	360	110	0.1	10,1	521	554			7111	151	ICX.	Colonial States	153	71.75	210	1001	1065	1003	1136

TABLE 117

Much Decided

Turb'y.

1899 Exami-

1899 Collec-

Serial No.

DATE OF

nation

tion.

Decided

July

9

July ...

Much

尼部於中亞家勢

Aug.

Sept

::::

Sept.

Much

- 685-1882-

Oct. : :

Oct.

Nov

ಹಾಗ್ ಜಿ

Dec

1096 1096 1137

Much

18は11118

June

-550588

June

May ::

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO,

SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISSIPPI RIVER, 100 YARDS FROM MISSOURI SHORE, JEFFERSON BARRACKS, Mo.

O. JORDAN.	iversity of Chicago
EDWIN	Cnj
J O	
Report	

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-ST. LOUIS, MISSOURI, TAP WATER.

	Chicago.
OAN	of
JORI	rsity
0	rei
EDWIN	Uni
Jo	
Report	

No. of Bac.	per Cubic Centi- meter.		3,450	6.500		8.000	1.500	4,500	000.1	1,600		002	600			1.400		670	2.790	600	730		086	0.00	OUG	310
	Presen lo .sd A					:	:	:	:	:	:	•	:	:			:	:								
	remper riv lo		38		.00	:	:						:	:				:								
	Temper of Wate		8.8	5.	.98	38	33										:									
	Heigh Mate				:		:	:		:		:	:	:	:			:								
NITRO- GEN AS	solerill	1	ان بر انت																							:E
	Setinin		.00.0 .00.0	.0062	.001	.001	100.	9700.	0.	0.	30.	0:	100.	.003	0.	.005	9.	0.	0.	0.	300.	0.	.e.c.	0.	.014	0.
GEN.	-sns	-			:	:	:	:	:	:	:	:	:		:	:	:	:	:		:	:	:		:	<u>:</u>
ORGANIC NITROGEN.	Dis-		: :		:	:	:	:	:	:	:	:	:	:	:	:	:			:	:	:	:			:
OZ	Total.		: :	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:		::
oo.	B Sus		# E E E	300	.048	ਰ.	.016	800.	910.	800.	80.	% 0.	800.	190.	850.	30.	300.	.020	50.	.01	10.	0.00	500.	30.	.1.	010
NITHOGEN AS	Pionimoid Dis- Dis- Dis- Dis-	3	176	192	.256	. 232	308	.160	144	. 160	. 136	.136	.073	190	104	104	960.	101	.088	966	960.	.072	.113	80.	.072	.008
NITTHO XXX	I'10T		8,8,	ा	.304	373	- 224	.168	91.	. 168	.216	.216	80.	.138	. 152	861.	10.1	. 138	.113	. 136	.168	. 128	3	.104	657	120.
	Freedm- monia.		18. S.	.016	10.	ろ.	390.	.00.	.016	900.	970.	.036	.013	÷00:	310:	800.	: :0:	.008	.012	810.	010.	.031	.012	800.	.000	.008
ED.	Marter. Marter.		0.6	00				ાં	رن ب	હેક	©: ©:		9.	-	9.	_:	ei.	ιċ	0.	7	er;	}		-	77	9.
OXYGEN CONSUMED.	By Dis-		1-1-		÷:	10		5.4	53 53	31.	3.4	4.6	1.6	-	÷	ος ος	3. ₹	4	+	∞ •••	4.6	₹ •	5.1	-	∞. ∞:	<u>د</u> د:
000	Total.		0 t-	00	∞ ∞		Į.	4.	5.6	5.4	5.6	5.6	ت. زن	† :	4.6	±. ∞.	4.5	4.00	-	4.2	7.6	<u>.</u> ت	5.5	±.	4.1	7
,91	Chlorin		10 17 10 17									9	7.1	7.1	<u>د</u>	9.6			11.7			<u>;</u>	15.3	15.1	5.1	15.3
ON ION.	pepued sus-	_ I	30.4																							
RESIDUE ON EVAPORATION.	Dis-	s	307.2 264.	200	216.	239.	238	: ::::::::::::::::::::::::::::::::::::	216.	250°.	330°	190	200	2 2 3 3	237.	270.	268.	28 1.	258	285	280	270.	272.	278	280	317.
RE	Total.		337.6	396.	504.	378	370.	338.	354	988	336.	365.	390	311.	328.	346.	35-1.	333	355	326.	318.	332.	300	330	313.	341.
	Odor		None	9 9	9.9	9 9	9 9	9 9	9 9	10	9 9	9.9	-9	9 9	9.9	99	9 9	3 9	9 9	9 9	9 .	9.9	9.9	9 9	9	-3 -9
	Color.		Muddy	9.	**	ıĢ	803	ıĢ	.e.	ri.	Muddy	9 7	.05	.G.		?	<u></u>	?}	ા	-12	ं!	-:	া:	15.	??	.22
APPEARANCE.	Sedi- ment.		Much	3,7	**	•	*	Cons'd	Little	79	7,7	9.9	;	7,	99	**	:	V. Little	Little	9 9	V. Little	Little	9 9	**	V. Little	Little
. 11	Turb'y.		Decided	7.	Decided	7,7	Much	:	:	:	*	*	:	9.	9.9	**	**	**	77	**	*,	,,	**	,,	9 9	7.9
OF	1899 Exami- nation.		June 7	- FG	:	el ylu	555	38	ug. 4	12	::			6 ,,		666 ,.	ct. 7	100	06 11	22	iov. 4		166 19	8 8		33
DATE OF	1599 Collec. E		June Ja	3 61	530	July 14.5	21	3	1112. 3.1	11 - 11	17	553	" 31 Sept	Sent. x		3	Oct. 60		61 ,.	96 .,	6	=	3	-	7	231
No.	Serial		9.11	19:	公公	XXX	3333	365	406	416	- X	515	550			202	-	-	3C	3	-	116	1000	-	•	1132

TABLE 119.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-SANITARY CANAL, KEDZIE AVENUE.

Report of EDWIN O. JORDAN,

11		1
No. of Bac.	per Cubic Centi- meter.	940.00 1,060.00 2,060.00 1,270.00
	Presend Abs. of	
	Tempers,	8. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5. 5.
r, C.	Tennpers of Wate	22
	Height etaW	
30- N AS	Setrativ	0000
NITRO	Setititi	800 800 800 800 800 800 800 800 800 800
N. N.	pəpuəd -sns	
ORGANIC NITROGEN.	Dis- solved.	
OZ	Total.	
	g p,pud	44.6 4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0 4.0
HEN AS	Diodiminoid Pistonia	6.6.6.4.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6
NITROGEN AS AMMONIA.	I'10T	20.00.00.00.00.00.00.00.00.00.00.00.00.0
	Freed m. monia.	
as og	BySuspd Matter	60 60000-00 60 40 6000
rGE	.bevios	0 t- 1-0 10 0 0 t-
OXYGEN CONSUMED.	Total.	α α α σ σ σ σ σ σ σ σ σ σ σ σ σ σ σ σ
ge.	Chlorin	9.8.8.8.4.4.8.8.4.4.6.9.9.7.7.6.9.9.2.7.7.6.9.9.2.7.7.6.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9
ON ION.	pepuəd gns-	26. 116. 10. 26. 24. 22. 25. 25. 27. 27. 27. 27. 27. 27. 27. 27. 27. 27
RESIDUE ON EVAPORATION	Dis- solved.	250. 272. 176. 176. 174. 174. 175. 175. 175.
RESEVA	Total.	251. 202. 202. 206. 180. 166. 179. 179.
	Odor	Gassy Gassy Gassy
	Color.	<u> </u>
APPEARANCE.	Sedi- ment.	Little Much Little V. " Little
AP	Turb'y.	Much Distinct Much Distinct Much Decided
SOF	900 1900 illee-Exami- on. nation.	30 May 22 23 24 June 5 25 25 25 25 25 25 25 25 25 25 25 25 2
DATE OF	1900 Collection.	Apl. 30 May 8 14 21 28 June 4 118 118 118 25
No.	Serial	1752 Apl. 3 1789 May. 1 1816 2 1850 2 1889 2 1892 June 1 1952 1 1992 2

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, BRIDGEPORT.

University of Chieago.

Report of EDWIN O. JORDAN.

No. of Bac.	Der Cubic	Centi- meter.	670.000	160,000	200,000	5~0.000	510,000	510,000	630,000	370,000	110,000	340,000	1,100,000	240,000	1,800,000	1.700,000	600,000	830,000	160,000
ce or Colf.		Pres Abs. o							•	:				•					
		lmoll'	e i	+	×.	£.		?;	32	= :	0)	6.5	Ť	**	<u>25</u>	=	<u>+</u>	16.	13.
		dm9T	0.	50	10.	0		<u>ښ</u>	0	· ·	21	0.	0	œ		œ	-	21	=
10 t	011 U	ioII iV	4.8	5.	٠,٠	4.6	63 70	10.	∞. •	3!	3.6	÷	5.		4	41.	9	10.	÷
NITRO- GEN AS	ธอ	Mirat	0.	5	:6	-:	0.	0.	0.	.05	<u>.</u>	.05	0.	.112	ಆ	0.	2.3	0.	0.
	_	Nitrit	0.	900.	.00.	.013	.001	.00.	870.	800	910.	.008	-:	.088	.017	.05	0.	.005	.018
NIC GEN.		-sng		:	:	:	:	:	:	:	:	:	:	:	:	:			:
ORGANIC NITROGEN.		Dis-	:	:			:	:		:	•	:	:	:	:	:	:		:
		b'baq stoT	97	333	:	:	:	.35	.: 91	22	:	.: 92	.:. 91	3:1	: ::	91	<u>.</u>	*1	89
AS A.	oid Am	p,vios	_	_	_		_	_	-				-		_	print	_		· ·
NITROGEN AS AMMONIA.	Albuminoid	-sid						3 1.0.1			~		_	_		<u> </u>		_	
NITR	I AI	Totyl						2.36	-							≎ §			1.08
		reed. Sinom	1					9.4											
EN MED.	pc oq	BySusi	8.5	90 90 	3 8.6	9.7	3 5.6	0C	6.3	3.3	~	35	£	9	50	{-	2	S	7
OXYGEN CONSUMED.	p	By Dis									÷.		{~	5.	5.	=	X	{-	9
3	"	Total						17.3			Merry								
'ou	ıiı	Срјо	88	116.	689	141.	213.	98.	33	88 88	9.	5.	33	8.	£	93.	-18	90.	60.
ON ION.	pa	opnod sng	42.	31.	85	34.	ŝ	33.	£	65	10.	25	3.	34.	35.	.10	30	57.	99
RESIDUE ON EVAPORATION	-1	Solved	104	330	314.	.178.	634.	375.	.380	341.	374.	285	257.	358	356.	430.	523.	386.	.062
REEVA		Total	446.	431.	3.13.	513	636.	408	414	387	384	291.	2.55 5.55 5.55 5.55 5.55 5.55 5.55 5.55	295	35 X	470.	562.	437.	356.
	Odor		Gassy	9.9	29 4	9 19	9.9	9.9	6 4	*	*,	9.0	9.	* *9	9 9	9 +	9.9	* *	*
		('olor.	ಣ	00	Muddy	7 7	3.	9	25	Muddy		<u>∞</u>	unife	e de la constante de la consta	-	7.	10.	03	20
APPEARANCE.		Sedi- ment.	Little	9.9		7 7	9.9	7 7		Little				7 7	*	Cons.d.	Little	Cons.d.	Little
AI		Turb'y.	Much	9.9	*	0 00		9 0					:	*	4 6	* *	9 0	9 9	9.
) F	Const	Exami- nation.	n3	0		000													23
DATE OF		r- Ex	olJan.		:	4.	; x	5 Feb.	., !!		. 9	5 Mar.	?1	. 0	9	AL	6	. 9	:
DA	1000	Collec- tion.		•	:	:	:	Feb.				Mar.			:	Apr.	.:	:	:
.oX	[13	alaos		1197	1237	1246	1301	1327		1397					15.59	1596	1634	1673	1712

TABLE 121.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LOCKPORT.

-
Z
4
-40
RD.
70
pered
0
0
\cup
Z
-
5
2
-
区
~
_
0
تىپ
Or
-
Prestig
0
Rel
_

No. of Bac.	per Cubic Centi-	meter.	445.000	830,000	S00.000	000,000	000 068 390 000	1.750.000		1,350,000	8 000 000	7.500.000	1,140,000	1.610.000	1.250.000	1.850.000	1.800,000	610,000	650,000	1.160.000	1.500.000	1 100,000	180,000	130,000
	resend s. of			:	:	:	:														_			
oture O.	mper iiv 10	9Т o		:	:	α) ()	2	30				70	55.	63	25					25.5		000	31.
	mper Nate		હ.	0.		o . c		0	0	· c	16	0	9.5	5	0	15.5	14.5	20.	15.5	19	18.55	<u>«</u>	5	(3)
	lelghi ets7/	1		:	:	:							•	:		:		•	:		:			
RO-	rates	NIK	-:	0.	90	· ·	0	0	, LC	178	-	9	0	0.	0	.25	.05	0.	0.	0.05	300	0	0	0.
NITRO-	trites	- 7	0.	.003	200.	0.0	10.	.032	960	115	015	0	0	0	.016	100	0.	0.	300	0	0.	.002	0	0.
ORGANIC NITROGEN.	papu -sn	S		<u>:</u>	:	:								:	:	:	:	:			:			
RGA	Dis- lyed.		:	:	:	:							:	:		:	:	:	-		:			-
02	otal.)T	:	:	:	:		:			:	:	:	:	:	:	:	•			:			:
700	p.p				7.48									-									_	
NITROGEN AS AMMONIA.	Albuminold -si		90	1.73	 	7.50	33		528	1.64	1.05	.86	1.36	16.	.56	중 증	.56	85.	.74	11	.57	99.	98.	.83
NITRO	I'1	οT	3.36	03.0 4.0	00.00 00.00	. c.	50.00	2.23	1.88	2.04	2.16	4.30	∞ ??	1.68	8:	1.48	1.44	6	1.59	7.	1.28	1.28	1.68	1.6
	mkəs .aino		11.8	ص م م	13.0	2000	6.6	, x	4.4	5.4	© ∞	5.6	11.4	∞ 7	9.4	00	10.6	3.00	12.8	18.3	.08	10	.06	24.
N ED.	rtter.	BY	8.3		7.0	7.5	6.8	00 33	9.4	·-}	6.9	25.5	5.0	8.9	∞ ∞	10.2	10.1	<u>ئ</u>	6.4	6.3	8	5.4	8.6	6.
ONYGEN	Dis-	s BA	19.2	ر د د د د د د د د د د د د د د د د د د د	0. 8 0. 8	7	8:6	=	9.4	9.6	11.2	<u>~</u> :	5.5	9.11	9.5	7.	80,00	∞ ∞	i~ ∞	8.7	6.5	8.6	7.1	8.8
Con	otal.	T	27.5	કો :	25 X	18.6	16.6	19.3	∞. ∞. ∞.	16.0	18.1	19.2	₽.1%	18.4	18.3	17.9	16.9	15.2	15.1	14.9	17.3	14.	\$5 \$5	<u>∞</u>
'əu	plori	o	112.	. 60	107	152.	86.	98.	58.	79.	74.	129.	175	115.	123.	54.	86	88	117.	150	268.	106.	219.	232.
ON ION.	pəpus -sns	ed	105.		32.5	30.	63	45.	73.	30	÷.	13.	13.	30.	65.	97.	 	28 28	4.	17.	56.	35.	67.	16.
RESIDUE ON EVAPORATION	Dis- ived.		410.	388.	i o	538	350.	398.	258.	384.	362.	505.	610.	:000:	531.	303.	386.	.90f	436.	490.	526	432.	625	650.
RES	otal.	T	515.	413	.000	570.	352.	440.	331.	404.	-50+	514.	623.	5:20.	596.	-100f	407.	134.	.440.	507.	586	467.	692.	.999
	Odor		Gassy	9 ;	; ;	9,9	9.9	7,	9,9	7 0	9,9	**	• ,	*	*	**	;		99	7	**	9 9	9.9	*
	Color.		Muddy	9 ;	: 3	9.0	7.9	99	83.	<u> </u>	Muddy	16.	9.	က့	77.	7.	ব	ಣ	77.	7.	771	4.	Muddy	.45
APPEARANCE.	Sedi- ment.		Much	Cons'd.	Tartle		Cons'd.	Little						7 9	Much	Cons.d.	Little	9 9	;	* 9		•,	Cons'd,	99
AP	Turb'y.		Much	: :	:	*	* 9	* *	9 0	7.9	"	9,1	. ,	;	:		:	:	39		*	9 9	7 9	7
OF	1900 Exami-	пастоп.	an. 3	5,	0 0	30	Peb. 6		13	<u>्</u> र	87	Apr. 3		:	•	Hay 2		15	?} :		une 5	<u> </u>	61 ,,	:
DATE OF	ర		Jan. 2 Jan	∞ <u>:</u>	33	:	5.	50			97 ,,	Apr. $2 A$	φ.	16	00 00 00 00 00 00 00 00 00 00 00 00 00	30	May	:	• • • • • • • • • • • • • • • • • • • •		:	•	. ,,,	June 25
.o.Z	Intros	3		1199	596	1303		-	1493	1555	1563	٦.		1675	1717			18	1856	1895	19:33	1957	0000	3035

TABLE 122.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of EDWIN O. JORDAN,

SOURCE OF WATER-DESPLAINES RIVER, LOCKPORT.

Chicago.	No. of Bac.	per Cubic Centi- meter.		88,000	2.700
sity of		Presen			
University		Temper TiA to			. 8 . 9.5
		Temper	0.0	0 +	00
		Heigh DiaW		-	
	ITRO- GEN AS	Mitrates	.45	က် က	파 뉴
	NITRO GEN	Nitrites	88	000	020

No. of Bac.	per Cubic Centi-	meter.	2.600	14,200	88.000	10.900	2.700	1.900		20,000	24,000		10,300	2.000	3,800		2.600	9.200	3,300	5.200	30.000	2,500	1.600	6.500
ce or Golf.	gesen or esem			:		:	:	•		:	:		:	:	•	:				:	:	:		:
eture O,	nper TAIr			:		∞ -	0.5	-13	30 50 50			85 101		: :33		33		% %			35.55		183 161 161	31.
ature D , re	nper Wate		0.	00	10.	0.	0.	0.	0	0.	0	0	10.	6:	6.	14.5	13.5	15°	33	16.5	30.5	24.0	20.5	20.5
	Mate otsW		- parent	, , , , ,		:		:	:	:		:			:		:		:	:	:	:		:
NITRO- GEN AS	вэзв.	HIN	.45	<u> </u>	ဗ	=	÷.	1.55	<u>કુ</u>	9.	£.	ಣ	0.	ः	ì-	e;	.e.	œ.	.e.	-	æ.	0.	0.	R. S.
Nri G1	sətir	NIC	.003	8 5	188	.026	10:	.015	910:	.0 .0	810.	8	810.	800.	10.	0.	0.	.65	0.	900	0.	8	0.	<u>ව</u>
IC SEN.	pəpı		:	:	: :		:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
ORGANIC NITROGEN.	-sie			:	: :	•	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
02	tal.	oT		:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
	An	pud	101.	90.	0560	.016	0.	.033	<u>₹</u> 01.	<u>=</u>	. 192	395	. 136	0.	.016	900	.208	- 1#	£30.	.048	308	. 136	90:	.033
TROGEN AS		Alos	.376	96F.	900	288	:30+	.361	. 22.	??????????????????????????????????????	∞ ?:	364	813.	æ.:	.416	.336	:3G	٠;	27:	3000	86.	:22:	×.	- S
NITROGEN AMMONIA	I I	10T	.48	. 256 256 256	35.55	.30-1	301	306	×	355	+	.656	2	388	.435	22:	52	.34	<u>x</u>	1.	爱.	108	:311	386
	mya .eine.			990.									.(558	.e.	<u>:</u>	- 3	. O	(5.1	169	(5)	3 o.	32O.	980	960.
ED.	rer.		4.	म् व		1.6	ŝ? -	re.	??	7	 	÷.	9	iG.	:	_	3.0	9	ĸ.	*****	ن ن	₹~ :	ů.	3.5
OXYGEN CONSUMED.	-sic	I VI los	 	عا يد ما يد		с: :	8.6	- ∞	6.1	φ. ∞	∞ 35	9.	ž	₹7. 0C	0.	<u>-</u>	oc .	₹ <u>-</u>	-	œ.	30 30	رة دن	ος 	ж Э
000	tal.	o.T		ic ic si o						с: :	·			8.50									0.3	
.or	ilorir	СР		ත ත			ic.	6.2	~	ئة زو	ci	:		∞ <u>.</u>							?? x		- C:	œ
ON ION.	pəpi -sn	rsd IS	20.	∞ -	10.	 		6.	36.	9	106.		 	er.		 	5	.30	~~	**	£	<u>2</u> 2	œ	36.
RESIDUE ON EVAPORATION	red.	AJOS (I	.869	350.	306.	254	288	. SEC.			14.	16:1	533	335	368	308		366	385	3968	37.	360		×
RE	tal.	0.17	718.	200 x	+000	278	223		<u> </u>	ος -1	.500°	226.	3		2. 2. 2. 3. 3.	152	£30.	396	394	400.	32	3.5	346.	374.
	Odor				Gassv	:		:	:		:			:		:	:							
	Color.)C.	esi <u>r</u>	<u> </u>		₹~	-	- The	기 .	ic.	ಚ	9.	7	in.	ıĊ	G	i.	₹.	.33	चां ।	်း	50°	ie.
APPEARANCE	Sedi-	ment.	Little	:	Little	9 9	1	**	Cons.d.	Little	Consid.	Little	:	4 9	9,9	* 9	9	***	**	9 3	Cons'd.	Little	d d	0
IV.	Turb'v.		Slight	:	Much			:		:			:	:	9 7		:				Much			Distinct
OF	1900 Exami-	nation.	2 Jan. 3	6 4		08 . 68												:	?}	æ :	June 5	:	19	36
DATE OF	1900 1900 Collect Exami-	tion.	Jan. 2.1	بر بر : :	: :			Mar. 5 Mar.		GI :		Apr. 24		16			May	:	:	ši :	June J	:		:
No.				<u> </u>	1270	1305	13:55	1458	14:41	27.0	1561	15.55	1637	1676	1717	1754	1787	X X	1891		1924	19890	1000	2033

TABLE 123.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DRAINAGE CHANNEL, BEAR TRAP DAM, LOCKPORT.

JORDAN,	University of Chicago.
EDWIN O.	^T niversity
Report of F	-

(
No. of Bac.	Cubic	Centi- meter.	1,700,000 380,000 3,000,000 10,000 3,000,000 1,650,000 3,800,000 1,360,000 1,360,000
ce or Coli.			
sture ; C.			2.8.8.8.8.8.8.8.2.8. 2.8.8.8.8.8.8.8.2.8.2
ernre C. C.		meT	で で で で で で で で で で で で で で
t of			
RO-NAS	sə	Nitrat	viw vi o o o o o o - o - o - o
NITRO- GEN /	sə	Nitrit	20000000000000000000000000000000000000
N.		puəd	
ORGANIC NITROGEN	-p	Dis solve	
ORG		RioT	
	Am.	p,pud -sns	17.6 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0
ITROGEN AS AMMONIA.	lbuminoid	-siq	25. 25. 26. 28. 28. 28. 28. 26. 26. 26. 26. 26. 26. 26. 26. 26. 26
NITRO	Albu	['10T	4:000000000000000000000000000000000000
		FreeAr	8.5. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
ED.		Bysus	# 0 69 69 69 69 69 69 69 69 69 69 69 69 69
OXYGEN		By Dis	40444044400 000101-1-4 010010
Con	!-	Total	F-04F-1004-0-0-1-1000-0-10-10-10-10-10-10-10-10-10
re.	ıiı	СРІО	5.6.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.
ON ION.	p	epued -sns	<u>0.4.4.4.8.8.4.8.4</u>
RESIDUE ON EVAPORATION	•1	Dis-	186.08.08.05.05.05.05.05.05.05.05.05.05.05.05.05.
RE		Total	204 210. 210. 220. 230. 230. 230. 230. 230. 230. 23
	Odor		Gassy
		Color.	- झुं राष्ट्रां राज्य झुं लंबा स
APPEARANCE		Sedi- ment.	Little V: V:
ΨV		Turb'y.	Much Decided Slight Much Distinct Decided Distinct Decided
3 OF		1900 Exami- nation.	Apr. 17
DATE OF		1900 Collec- tion.	Apr. 16 May May June
.07	I I	Seria	1719 1719 1719 1719 1719 1830 1930 1931 1931 1931 1931

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-DESPLAINES RIVER, W. BANK, N. JACKSON ST., JOLIET.

Report of EDWIN O. JORDAN,

No. of Bac.	Cubic	Centi- meter.	600,000	1,050,000	830,000	1.210,000	500.000	410,000	630,000	970,000	2,050,000	1.050,000	1.750.000	380.000	110,000	1.120.000	40.000	90.00	350,000	440.000	180,000	850,000	630,000
ce or Coll.				:	:	:		:				:	:	:		:		:		:	:	:	
ature ., C.			9.5	<u>∵</u>	0	÷	-16.	=======================================	33	-13.	=======================================	***	=	=	<u>2</u>	16.	30.	21.	<u>₹</u>	<u>ક</u>	16.	<u></u>	33
ature O. To.		Temil To	0.	10.	0.	4.5	0	0.	0.	0	0	÷	::	۸.	∞i	22.	80		30.	÷	33	.54	21.
lo t	dg ore		•							:			:				0	:		:			
RO-NAS	89	Nitrat	0.	0.	<u> </u>		<u>С</u>	.e	23	??	ıĢ	0.	0.	-:	S.	.0.	.60	.C.	0.	ıc	0.	0:	0.
NITRO- GEN A	J	Mitrit	:00:	300.	SE.	10.	920.	10.	190.	.050.	910:	980	.035	80:	<u>x</u>	.035	800.	910.	0.015	.032	300	30:	10.
GEN.		puod sus		:	:	:	:	:	:	:	:	:		:	:	:	:	:	:	:	•	:	:
ORGANIC NITROGEN.		siQ solve		:	:	:	:			:	:	:	:	:	:	:	:			:	:		
0%	l.	Tota		:	:	:	G)	?)	30	9	and .	9	9	6		:	•	:	9	6		:	÷
	1 Am	p,pud -sns	9.	_	_				_											_			
NITROGEN AS AMMONIA.	Albuminoid	Dis-	2.8		_																		
NITRO	Albu	I'10T	3.4	ون د.	2.3	1.38	.68	£.		œ.	.c.	.68	.46	=	33.	. 296	.31	27	.33		.36	.352	.376
		FreeA	8.11	=					\$ i		1.16	-	en:	35	900	-		e i	00	3.6	-	50	FC.
SN (ED.	pd 	Bysus	1.9	00	-3	9	9	<u>{</u> -	7.6	63 63	\$?	5.6	-	1.3	} <u>~</u>	9.6	63	e5	10.	1-	3	30	-
OXYGEN CONSUMED.	·pa	By Dia	26.7	<u>о</u>	œ.	1-	200	•••						10.	4	9	10	9	4.5	10	6.1	6.2	3-
000	1.	Tota	28.6	∞	16.	x. :-	10.2	10.6	5.0	œ.	() ()	11.6	¿-	1- 0.5	5.6	G:	6.5	6 5	9	6.1	6.6	\ \-	7.5
, *ət	ri1	СРІО	113.	110.	95	38.	Ξ.	€} €}	08 -	=		45.	41.	30.	33	41.	30.	27.	31.	40.	-		58.
ON TON.	pa	puod sus	38.	. 699	6.	88	115.		110.	<u>ان</u>	23	22	-	=	55		5	· - 5		88	_	٠:	95.
RESIDUE ON EVAPORATION		siU solve	506.	124	426.	295.	196	203	236.	217.	226.	301.	268	27.3.	1357	276.	215	211.	226.	232.	00 to	231	260.
RE	.1	nioT'	5.11.	190	133.	381.	31	220.	316.		· 257	386.	283	283	.3 IO.	300	1551.	20 20 20 20 20 20 20 20 20 20 20 20 20	210.	262.	270	237	285
	Odor		Gassy	**	:	9 9	9 9	* *	Slight	Gassy	9.9	9.9	9 9	4 9	49 0	*	* *	:	9 0	9 9	9 9	? .	9.9
		Color.	Muddy	***	ಟ	50	.12	-:	Muddy	;	दर्	:35	<u></u>	-	ଦୀ	es.	e:	≎∶	e.:	.s.	89.	ej.	e0;
APPEARANCE.		Sedi- ment.	Little	v	:	Much	Little	.,	Much	Little	:	:	:	*	:	**	**	**	9 9	•	. 9	:	:
AP.		Turb'y.	Much	:	• •	:	7.0	9 9			:	:	9.		Decided	Much	Distinct	Decided	9.9	Distinct	Much	Decided ,	Slight
OF	1970	Exami- nation.	Jan. 3			:		Feb. 6	. 13	30		April 10			May 2		15	. 25	. 29	une 5		. 19	: 35°
DATE OF	10xx0	ollec-Exami-	63	īx ;	10		666	-	©?	<u>6</u> :	.9°	-	9	:		May ?		:	. 2X	2000	1	X.	. 25
NO.	_	Lios	1162 Ja			. 193																. 66	· \$250
,			-		-		0.00	-		-	_	16	16	-	7	the same	-	orani orani	1	1	10	15	हा.

TABLE 125.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, JEFFERSON ST., SOUTH JOLIET.

of EDWIN O. JORDAN,	University of Chicago.
Report	

No. of Bac.	Cubic Centi- meter.	1.500.000 3.200.000 3.500.000 1.380,000 600.000 1.600.000 1.600.000 1.650.000 600.000 540.000 540.000	
	Presen		
	remper riv 10		
	Temper otsW lo	<u>्रक्ष क्ष क्ष क्ष क्ष क्ष क्ष क्ष क्ष क</u> ्ष क्ष क्ष क्ष क्ष क्ष क्ष क्ष क्ष क्ष क	
	Heigh JisW		
NITRO- GEN AS	Nitrates		
IN G	Nitrites	2000 2000 2000 2000 2000 2000 2000 200	
HC HEN.	popuod sns		
ORGANIC NITROGEN.	Dis-		
OZ	Total.		
	p,pud -sns	888.44.93.94.1.38.88.89.44.89.94.1.1.00.	
NITROGEN AS	Pioniminos -sid -sid	1888 1888 1888 1888 1888 1888 1888 188	
NITRO	I'10T	8658 6848 41588 8886 688	
	Freeding.	8.54.58.55.58.59.59.59.59.59.59.59.59.59.59.59.59.59.	
N ED.	BySuspd	4 w 4 w w w s s - w s s	
OXYGEN	By Dis-	ו מי מי היי מי מי מי מי מי מי מי היי היי	
Con	Total,	<u>e o si - o c c c c c c c c c c c c c c c c c c</u>	
.er	Chlorin	8289558488848644	
ON TON.	pepued -sns	8,16,8,3,19,6,5,8,0,5,8,0,5,8,0,5,8	
RESIDUE ON EVAPORATION.	Dis-	8.8 8.8 8.8 8.8 8.8 8.8 8.8 8.8 8.8 8.8	
EVA	Total.	25 25 25 25 25 25 25 25 25 25 25 25 25 2	
	Odor	Gassy	
	Color.	म्त्रे में सह में सह में सह में कि क्षा का का का हा का में कि का कि में में में हों में हुई में कि कि का का का हा का में कि का	
APPEARANCE	Sedi- ment.	Little Cons'd Little	
V	Turb'y.	Much ": " " " " " " " " " " " " " " " " "	
S OF	1900 Exami- nation.	25 Mar. 6 26 April 3 27 April 3 30 May 23 28 1 1 15 29 1 1 15 20 1 1 15 20 1 1 15 21 1 15 22 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
DATE OF	1900 ('ollce-	May 150 110 110 110 110 110 110 110 110 110	
.oX	Serial	1451 1451 1451 1452 1453 1453 1453 1453 1453 1453 1453 1453	

TABLE 126.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-KANKAKEE RIVER, WILMINGTON.

29
A
2
princip
0
_
<u>_</u>
\circ
7.
-
-
~
page
t
0
سه
-
0
_
-
0
K
_

No. of Bac.	per Cubic Centi: meter.	2.800 2.800 2.800 2.900 2.0000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.000 2.0
	Presen To sof	
	Temper	
	Temper of Wate	9000-0000-0000000000000000000000000000
lo 1	dgləH əts77	#####################################
NITRO- GEN AS	solithin selection in the selection is a selection in the selection in the selection is a selection in the selection in the selection in the selection is a selection in the sel	0.00
GEN.	popuod -sns	
ORGANIC NITROGEN.	Total. Dis- Solved.	
	F sus	40-00000000000000000000000000000000000
EN AS	Pionimoio p.vios -sid	88 44 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
NITROGEN	I'10'I'	8.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5.5
	FreeAm	99.00 90.00 90
EN MED.	Bysuspd	674 94 F-4 13 4 1 1 1 1 1 1
OXYGEN CONSUMED.	-sid val	α α α μις - α α α α α α α α α α α α α α α α α α
၁၅	Total	
.en	riaofd')	သည္။ သည္က သည္က သည္က သည္က သည္က သည္က သည္က သည္က
ON TION.	papuad sus	
RESIDUE ON EVAPORATION	Dis.	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
EV	Total.	
	Odor	
.:	Color.	$\frac{\mathbf{K}}{\mathbf{A}}$ $\frac{\mathbf{K}}{\mathbf{A}$
APEARANCE.	Sedi- ment.	V. Little V. Little V. Little N. Little Much Little Much Little Much Little V. Little Little V. Little Lit
1.	Turb's.	Slight Distinct Slight Decided Slight Nuch Slight Much Slight Much Slight Much Slight Much Slight
DATE OF	1900 Exami- nation.	25. 1an. 25. 25. 25. 25. 25. 25. 25. 25. 25. 25
DAT	1900 Collec- tion.	Jan. Mar. Mar. May. May. May.
.o.Y.	Serlal	28

TABLE 127.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, MORRIS.

Report of EDWIN O. JORDAN,

No. of Bac.	per Cubic Centi	meter.	000 400 1	1,300,000	1,230,000	215.000	320.000	520.000 1 000 000	310,000	800,000	1,000,000	000,040	225,000	60,000	55,000	13,000	#,000 %	000	640,000	110,000	8.000	24.000
	lo .				:		:	:		:	•	:			:	:	:	:	:	:		
oture .O.	rpers		4.11		ب ب ن	0	4.	ب س رو		0	<u>.</u>	တုံ မ	<u> </u>	15.		55 50 60 60 60 60 60 60 60 60 60 60 60 60 60	, 20 31 31	17.		? ?		30.
ature r, C.	npers State		0.0	: o :	00		, O. C.	0.0		0.	-	90	o တ	10.	15.	÷:	∞;	. 90	20	10	<u> </u>	<u>.</u>
	eigh Vate		ىن سى:	ن بن	6.0	 	6.6 6.6	ۍ د اب د	16.6		12.5	9.1.	- 1C	6.	4	7	44.0		4, 71 25 C	o. ∠	91 00 00	4.3
RO-NAS	setes	Nitr	.03		4. 7	<u> </u>	.65	م زه	9.99	6.	ος.	1.15		<u>و</u>	٠÷	.0 .0	4:	∞ a	3 5			.g.
NITRO GEN	səti.	Nit	1.12	020	8.8		.11	038	0.00	.034	.05	.018	900	0.	.034	.03:	4.		000	# G	3.68	िः
IC IEN.	pəp -sı				:		:	:_		:	:	:			:	:	:	:	:	:		
ORGANIC NITROGEN.	is-		:	::	:		:	:		:	:	:	:		:	:	:	:	:	:		:
OZ	tal.	O.T.		: :	:		:	:		:	:	:	:		:	:	:	:	:	:	:	
	An	pud	.692	96.	1.36	. 150 144	969.	. 448 676 676	.536	.328	.248	296	150	104	.032	.128	.035	+0·	980.	20I.	88	880
TROGEN AS		vios	1.008	\$ 87.	8} ?	# 886: - 886:	. 224	.304	144	. 288	.296	8 1 8.	0000	192	.352	. 168	272.	882.	184	502.	# 6: 6:	.176
NITROGEN AMMONIA	Albu	10T	7.7	1.02	1.48	65.	36	. 705.	688	.616	.544	110	4.7	296	.384	536	300 E	. 33 20 20 20 20 20 20 20 20 20 20 20 20 20	XX 5	300	-04. -06.	264
	-mA:	Free	7.9	4.4	1.84	2.44	1.04	7.96 	.36	1.008	1 8.	1.86	50.1	1.48	3.96	2. E	1.84	1.73	gel.	200 200 200 200 200 200 200 200 200 200	20.03	
N ED.	ter ter		41 H	0 02 0 03	9.11	ە ئەت	9.4	6; c	1	5.6	1.4	11 ,	0.1	1.7	C.S.	6.	o.	œ ,	- ,	0.0	25 2. 5	6
OXYGEN CONSUMED.	vpəv.	By I	3.4	6.4	गा १	5 .c	9.4	6.1	0 67	53.	<u>.</u>	φ. 0	20 e0	5.0		5.5	6.3	6.5	6.5	7.0	0 rc 0 o	6.9
Con		oT	5-0	મું ∞ કું ઝાં	0	010	00		60	00	8.4	11.6	_	7.6	7.6	6.4	5.0	8.9	0.0		. cc	
.9t	Jorin	чэ	60.			<u> </u>	11.	2 000 1 0000	916	13.	11.8	o. o.	9. 9.	15.	38	-82	19.	63	<u>∞</u> ;	- 1 -5		
ON ION.	qeq	uəd 1S	ထို		217.	4. 80 0 00	407.	64.	314.	140.	38	198.	30 .		6.	.12	9	6.	တ်	27.0		19.
RESIDUE ON EVAPORATION	ed.	ros	413.	348.	250.	2330	205.	250.	136	184	188	170.		2 5 2 5 2 0 2 0 3	296.	.308	330.	236.	300°	· · · · · · · · · · · · · · · · · · ·	148.	25.5 25.5 25.5 25.5 25.5 25.5 25.5 25.5
RE	al.	toT	420.	366.	437.	260	612.	314.	303. 450	324.	226.	368.	207 903	264.	30.5	239.	236.	242.	.50e	266	276.	247.
	Odor		Gassr	:	9.9	: 1		:			•								:	:		
	10101	Color	9.	ئە ئەن	<u>. 15</u>	: :: :: ::	₹ 7.	Muddy	من يو	600	o į	ಖ	şç o) es	7	: 25	-	35	۽ قد	કું.	is e	.63
APPEARANCE.	Sedi-	ment.	Little	Much	Much	Little		Little				Cons'd.	; ;	Little	9,9	;	9.9	;	3,7		, , ,	3
AF		Turb'y.	Much	; ;	**	,,,,	***	,.		"	**	:	, ,			Decided		Distinct			Nieting	
OF	1900	Exami- nation.	5 Jan. 8	= ×	:	30 Fob 8			Mar. 8	3		3 Apr. 4	;;	7 6 77	May	*	., 16	5		June 5	2 2	23 ,,
DATE OF	1900 1900	Collec-	1.	; ;	96 ;;	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		61 ,,	Mar. 7	61 ,,	4.9	Apr.	; ;	010	3000	May 7		18 31		June 4	;;;	36
No.	f Isi	rəS	1196 Jan	1209	1296	1307	1373	1400	1461	1506	1571	1602	1640	1715	1758			1854	1893	1938	1962	2037

TABLE 128.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-FOX RIVER, OTTAWA.

Report of EDWIN O. JORDAN.

	No. of Bac.	per	Centi: meter.	3.700	30.300	12,600	25,300		1.100	50,000	200,000	5 × 5	0.000	009 8	901:3	8.500	00x	3	002.3	6.200	9,600	3,600
			Prese Abs. c											-								
			qməT A lo	6	2)	333	-	+	31	{ -	16.	=	+	50	, c.	10	67	19.	??	.96.	. [≳	35.
			gm9T gW lo	oi	0)	23	÷.	ıc.	33	0.0	~	e4	7		2.5	19.	601	19	£3	33	.98	27.
	10 1	ogi kpi	gl9H 37/	65	500	65	****	-	-	20	-	7		÷	73"	00	32	es ru	33		3	5.5 10.
	RO-	รอ	Mirrati	?!	ù.	77.	œ	G:	9	x		16	0.	0.	.05	0.	3	12	10	0.	ο.	18
	NITRO- GEN A	<u> </u>	Minit	8	.016	10.	010.	10.14	.016	3	0.10	.025	.014	10.	00:	.000	0.	900.	.013	φ.	0.	. 036
	NIC OGEN.		epuod -sng -sng		:		:					:	:					:	:		-:	
	ORGANIC NITROGEN.	-	siq Sid		:	:	:	:	:	-	:	:	:	-		:	:	:	:	:	:	:
			p,pud	:	: ::	:		9	0.4	96	76	80	: 68		23	80		76	<u>∞</u>	- X	:080:	?; ?;
	A.S.	oid Am	-sng	1				_	_		_	_		_	_	_	_	_	_	_		
	NITROGEN AS AMMONIA.	Albuminoid	Dis-					_	-								_					
	NITR	AIR	['10T'	966	77.	s;	.336	384	.35	.584	1.13	967	.416	5.14		.504	.384	.488	+	.36	:335	. 256
			reedr Sinom	.078	990:	80.	E	.256	÷67.	.156	++2.	91.	3.	90.	.016	810.	910.	30.	-	810.	.038	.O.
	ED.	pd	BySusi																		9.	
	OXYGEN CONSUMED.	p	sid val solve	10	-	∞ ∞															6.5	
	03	-1	Tota	10	-4	=	7.4	χ .e	α re	15.2	19.6	133	37	9	12.9	14.9	8.6	©: ∞	52	?! ∝	6.8	5.7
	16.	ı[1	СПО	6.8	6.	6.	2.	 G;	65	5.6	0? 0?	≎ર	oc.	e2.	83	3.	_	??	 	9.	\$ } 00	
	0.N 10.N.	p.	bende Sus	0.	ci.	 	ος. ?)	36.		139	500.	97.	22	623	<u>∞</u>	36.	9	-	21		£	.05
	RESIDUE ON EVAPORATION	T	Solve	4.8	310.	35	33	170	-50g	-EE		:0x	348 348	273.	262.	568.	301.	565.	22.5	200	200	276.
	RE EVA	1.	nioT'	×.	315	300	3.0	196.	230.	304	131.	3(5)	(35) (35)	534	:380	304.	307	276.	314.	.3.31	X :	295.
		Odor																				
			Color.	.15	-	23	Maddy	es.	e.	- e	:0:	<u>~</u>	c.	+ .	7.	- -	7			7	S !	
	AFFEARANCE.		Sedi- ment.	V. Little		Little		('ons'd.	Little	V. Little	Much	(b.sio.)	9 9	*	;	Little	:	* * * *	:	:		:
	7.		Turb'y.	Slight	:	b -d	Much	•	:	Night	Much		Decided	Much	;	9.	Decided	Much	Slight	Decided	Distinct	Slight
	30 3	1000	Exami- nation.	Јап 5	22	5.		Mar. 13		×.	Apr. 4	:	X	:	May 2		9 :	* "	June 6	:	61 ;	37
-	DATE OF		Collec- tion.	Jan f	=	<u>x</u> :	= :	Mar. 12	9.	-	Apr. 3.		-)	2.1	May 1	X	:	? } ? }	June 5.	= :	X.	: :
	.07	[13	iros			1543					7						-X23			C#35	282	2020

TABLE 129.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, OTTAWA.

Report of Edwin O. Johdan.

No. of Bac.	per	Cubic Centi- meter.	10,000	225,000	173,000	000,076	10 (00	35.000	000,000	68,000	210,000	165.000	12,000	6.000	5.000	1.700	7.300	008.1	000	17.300	000 65	19, 100	17,000		
		Prese			:	:			:															:	:
		V lo	e;	ei e		er er	16.	200	4	7	(-	16.	=	. 	9	27	16	3	19	6	26.	56	33	:	
r, C.	919 911	Tempo R/I lo	c:	\$ £	: i =	; c	000	· 65	5.	ಣ	G	99	6.5	-11- -		.27.	6:	3	19	33	5.	5	62	:	
		giəH gV/	3.5	ر د د د	بر ن ن	910	10	10	5.	က	;	rc.	٠.	-	33	33	35	. 65	20		35	200	3.5	:	:
RO-	se	Nitrate	1.45	1.55	G #6	15	300	1.9	.55	9.	ල. ද?				ī.G	∞.	7	9	9	0	7	77	58.	:	:
NITRO-GEN A	86	Nitritt	.15		820	920	39	.034	.024	.032	890.	70.	90.	-:	.08:	80.	.18	9	.16	· 67:	200	9		:	:
AIC GEN.		-sus	:	:	:			:	:	:	:	:	:	:	:	:								:	:
ORGANIC NITROGEN.	-	-sid solve		:	:				:	:	:	:	:	:	:	:					:			:	-:
. —	1	[stoT	:	:	:		:	~	:	~	.:	:	~		~	:	~	_	~		-	-		:	-:
70	d Am	-sng	. 13(بن ورو	38	0	14	<u>300</u> .	.03	380.	80.	.10	.168	30%	.328	.23:	80.	=	.048	.01	.07	0	.010	:	-
NITROGEN AS	Albuminoid	Dis-	-63	1 6	336	.216	968.	.288	.192	ō.	.168	.176	.256	.368	.424	.248	.216	.274	300	358	. 192	308	.176	:	:
ORTIN CRA	Alb	I'10T	.76																					:	
		FreeAn	∞·	e} → 0	96.00	1.6	1.6	1.36	88.	.48	.4	1.14	1.04	.746	. 184	454	1.024	99.	94.	96.	€.	1.36	88.	:	:
ED.		BySusi	9.																					:	:
OXYGEN CONSUMED.		By Dis		v ix	- 10	30	7.5	7.3	9	3.4	4.6	0	9	<u></u>	<u>ن</u>	∞	~	5-	9	6.1	6.2	6.9	5.5		:
၁၅	•1	Total		10.2 2.3			7 .	∞	φ. •			} -	=		<u>≃</u> :	133	20			7.4	7.6	6.6	6.	:	:
:e:	ıiı	Срјо	36.		0 7	30	15.5	15.5	∞ —	٠ •	× 2		7.6	. S	5.6	10.4	15.8	17.	14.	16.	14.	21.	21.8	:	
ON TON.	pe	puəd sng	86		. %	6.	+	6.	08°	ص م	13.	30.	61	#	0	.00	4	10.	16.	88	6.	++	250 250	:	:
RESIDUE ON EVAPORATION		Solve Solve	406.	300 - 300 -	298	250	300.	310.	174	199	33.0	384	391.	410.	375.	252.	242	305.	246.	.540°	280.	214.	343.	:	:
RE Eva	.1	Tota	434.	532	350.	256.	314.	316.	254.	 	273.	414.	410.		375.	355	546.	315	262.	273.	.386	258.	274.	:	
	Odor			:	Gassv		:	:		:	:	:	:	:	:	:				:	•				
.;		Color.	85.	- -	Muddy	ें हें हें	16	ಀಀ	-Jr	7	.12	ကဲ	-	. 45°	7.	***	ग ु.	<u>.</u>	771	→.	w.	٠ ?	€:	:	
APPEARANCE		Sedi- ment.	V. Little	Little	:	;	:	99	Cons.d	Little	**	:	•	"		*	**	*	*	**	9	y,	7,	:	
1.	Į	Turb'y.	Distinct	Much	:		_	Much		3 9	:	Decided		_		Mueh	Decided	9.9	:	:	4.9	Distinct			
OF	1000	Exami- nation.	4 Jan. 5	: :		Feb. 6		Mar. 6				Apr.	•	;		May 2		., 16		5 June 6	13	61 :	1, 26		
DATEOF	1000	Collee-I	Jan. 4J		o (c)	Feb. 5	30	r. 5	22)			Apr. 3		17	<u></u>	May 1 N	00	15	e€. •••	fune 5J	,,	:	35		
No.	[B	Seri		1255				1455 N		1529				1683		-				1931 .	1964	1998	2036	:	

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER-BIG VERMILLION RIVER, LA SALLE.

University of Chicago.

Report of EDWIN O. JORDAN,

			- ,	
No. of	per Cubic Centi- meter.	4 in si ji	2000 2000 2000 2000 2000 2000 2000 200	000.09
	Presenc			
	Tenrpera of Air,			
	sraqmaT iatsW lo			
	Height 1918W		φ	
RO-	setartiz	401-801-888 661-861-888	+ 0 0 - 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	2.15 .65
NITRO-GEN A	Nitrites	800080008 800086	30000000000000000000000000000000000000	95.9. 4.5.
IC EN.	pəpuəd -sng			: :
ORGANIC NITROGEN.	Dis- solved.			: :
OZ.	Total.	• • • • • • • • • • • • • • • • • • • •		. 0 . 0 . 0
	a A B B B B B B B B B B B B B B B B B B	00889999999999999999999999999999999999		0.016
TROGEN AS	Albuminoid biolioid bivios	200000000000000000000000000000000000000	2 £ 8 8 2 8 8 8 8 8 8 8 2 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1	===
NITROGEN	A I'10T	80. 10. 10. 10. 10. 10. 10. 10. 10. 10. 1	\$45.55.55.55.55.55.55.55.55.55.55.55.55.5	-158 -664
	FreeAm- monia.	8 - 5 3 m 5 5 5 8 5 8 5 8 5 8 5 8 5 8 5 8 5 8 5	8 8 1 1 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	920
ED.	BySuspd Matter.	6 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	010-11- 004-014-0100-0-0-00-0-0-0-0-0-0-0-0-0-0-	= 19.0 = 19.0 = 19.0
OXYGEN CONSUMED.	By Dis- solved		1 00 00 01 + + + 00 00 00 + 01 00 + 00 00 00 00 00 00 00 00 00 00 00 0	5. 50 4.73
000	Total.	. 4 20 50 00 10 10 10 10 10 10 10 10 10 10 10 10		÷ <u>∞</u>
.91	Chlorin	왕왕왕 왕왕왕 왕왕학	:	55 55 55 45
ON ION.	papuad -sug	, o o o o o o o o o o o o o o o o o o o	<u> </u>	3. 5. 5. 5.
RESIDUE ON EVAPORATION.	Dis- solved.	266 266 266 266 266 266 266 266 266 266	#88888888888 #8688888888888888888888888	99.99
REE	Total.	200 - 100 -	2	- 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	Odor			: :
	8			::
ដ	Color.		$\frac{1}{(4)^{1/2}} \sin \frac{1}{2} \cos \frac{1}{2} - \frac{1}{2} \int_{\mathbb{R}^2} d^2 \sin \frac{1}{2} \sin \frac$	
APEARANCE	Sedi- ment.	Cons'd V. Little Nuch Little Cons'd Teans'd	Much Clittle Clittle Clittle Little Much Little	Much
N.	Turb'y.	Slight Nuch Slight Much Slight		Much
30	1900 1900 Collec- Examition.		Nay 827 13 83 82 14 15 15 15 15 15 15 15 15 15 15 15 15 15	ន្តអ
DATE OF	ec- E	»	Serial Se	150
U	1900 Colle tion.	Feb. : : : : : :	Apr	::
No.	Serial			2008 2008 7808

TABLE 131.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, LA SALLE.

Report of Edwin O. Joudan, University of Chicago.

No. of Bac.	Cubic	Centi- meter.	30.100	18,200	154,800	195,000	0000	195,000	53,000	200,000	000.85	320,000	120,000	260,000	360,000	65.000	16.000	25.000	4.600	8.500	1,800	5.700	8.600	25.500	3,900	11.000	3,700	
ce or		Pres.	:	:	:	:	•	• •		:	•	:	:	:	:	:	:	•	:	:	:	•				•	:	
ature,) 	Temp	:	:	:	:	:		:	:			:	:	:	:	:	•	:	:	•	:	:			•	:	
		TempT W 10		:	:	:			:	:	:	:	:	:	:	:	:		:	:	:	:	:		:	•	:	:
t of	dg ete		:	:		12.4	2	18.3	14.9		14.6	25 25	33	19.6		17.6	15.6	. 9	24.8	15.3	14.	Ŧ. II	11.6	∞. 27	11.8	11.6	12.4	
NITRO- GEN AS	sə	Nitrat						- 1																				
Nr.	sə	Nitrit	.054	.073	87.0	210	010	38	0.58	.036	870.	.01 <u>2</u>	.03	016	.016	920.	.052	.076	90.	-	.36	ક્રું	.14	80	4.	ಣ	સુ	
IC FEN.		sus	:	:	:	:	:		•	•		•	:	:	•	:	:	:	:	:	:	•	•	:	•	:	•	
ORGANIC NITROGEN.		Dis		:	:	:	:		:		•	•	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	
ON	1.	Tota		:	:	:	:		:	•	:	:	:	:	:	:	:	:	:	:	:		:	:	:	:	:	:
	1 Am	p,pud -sng	.188	₽.	#B.	1.55	201.	0 to	.216	.168	.198	715	. 152	.195	272	.176	.133	.072	.113	0.	210.	980.	1 98.	.152	.10 4	.016	104	
TROGEN AS	Albuminoid	Dis-	.392	.56	966	255 250 250 250	100	201.0	.248	.24	. 168	848.	.176	.248	. 192	356	. 246	288	305	.44	.308 803	.256	. 168	. 152	184	308	.152	
NITROGEN AMMONI	Albu	Total	.58	9.	<u> </u>	# E	14.	713	.464	.408	966.	86.	.328	-44	.464	.433	368	.36	.416	.44	સં	.342	.232	30%	. 288	224	.256	•
		FreeAr	3.9	7	ئة د		0.1	7.6	.:	1.04	1.32	4.	.50 1	.336	.336	.536	.528	.504 40c.	.113	311.	.176	.256	.128	.216	.416	.738	89.	:
N ED.		BySus	5.	rů.	χ.		0	9.6	1.3	1.4	1.4	11.2	2.1	€. 4.	, j	-	ಣ	7	3.9	ة: آ	-	9	0.	1.4	9.	}~	1.4	:
OXYGEN CONSUMED.		By Dis						9.														+	.9	6.5	6.	50.00	5.6	:
်ပ္ပိ	۱٠	Total	·	œ.	دة د ق	ο -		17.5									ος ος				7.3	}- }-	6.	7.9	6.6	6.5	·-	:
.ər	ıiı	СРІО	39.	. 68			14.	0	13.	13.	12.	·-	6.	∞. 7	90°	9. 9.	10.	7. 6	5.6	15.2	<u>:</u>	<u></u>	14.	1.1	19.	16.6	18.	•
ON.	pa	Sus	4.	7	0;	56.	01	36.	16.	57.	:S	352.	99	84.	244.	46.	53.	; ;	16.	61.	16.	٠ì	ಣ	37.	67	33.	.9	
RESIDUE ON EVAPORATION		Pis-	412.	374.	354.	292		10 00 00 00 00 00 00 00 00 00 00 00 00 0	25.4	252.	249.	33	188.	160.	186.	230.	245.	260	360.	246.	.992	997	226.	226.	258.	225	225.	•
RES	-1	Total	416.	378.	354.	318 000 000		473.	270	27.9.	274.	444.	248.	244.	1 30.	276.	. 86%	282	276.	307.	282	268	999	263.	287.	260.	231.	:
	Odor				:	Gassy	:	:						:	:						:	•			•		:	
		Color.	8.					yj a			કું	ıů	₩.	4.	7.	-j.	ಣ	4.	ıū	4.	4.	₹.	ಣ	£8:	7	ক্ষ	ಣ	•
APPEARANCE.		Sedi- ment.	V. Little	,,	Little	Cons'd	V. Little	Much	Little	"	*,	Much	Cons'd	Little	Much	Little	Cons'd	Little	*,	9 9	3,9	"	:	9.9	**	"	9,9	:
Ar		Turb'y.	Decided			Much		Silgnt		7.7	9.7	3.9	:	;	9,9	9.9	Decided		Much		Decided		Distinct		Decided		Distinct	•
OF	000	Exami- nation.	2 Jan. 3		., 17	त्र त ः		reb.	16 91	200	Mar. 8		21	33	3 Apl. 4	6. 11	18	::	May 2	0,,,	,, 16	., 93	30	June 6	., 13	20	37	
DATE OF	000	Collec- Examition.	Tan 2		91 ,,	हा ह	90	rep. 61	06 "	270	Mar. 6	13	07	26 33	Apl. 32		17	24	May 1 1		15	35	666 11	Tune 5.1)	61 ,,	26	:
.ov	ŢŢ	Seris	1165		1244	1272		1342	1401	1435	- Labor	_	1531	1566	_	1645	1681	17.35	1762		1826	1862	1899	1933	1969	5001	2039	:

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

:23
Print
-
-
4
00
Y
-1
4
12
-46
3
\circ
7.
41
16.
CE
-
10
_
7
8
ND
AND
A
A
A
OIS A
A
A SIONL
A Sto
LLINOIS A
LINOIS A
LLINOIS A
-ILLINOIS A
R-ILLINOIS A
R-ILLINOIS A
ER-ILLINOIS A
TER-ILLINOIS A
ATER-ILLINOIS A
ATER-ILLINOIS A
TER-ILLINOIS A
WATER-ILLINOIS A
WATER-ILLINOIS A
ATER-ILLINOIS A
OF WATER-ILLINOIS A
WATER-ILLINOIS A
E OF WATER-ILLINOIS A
RCE OF WATER-ILLINOIS A
RCE OF WATER-ILLINOIS A
E OF WATER-ILLINOIS A

University of Chicago.

Report of EDWIN O. JORDAN,

No. of Bac.	per Cubic Centi- meter.	32,000 33,000 113,000 113,000 114,000 115,000
	Presend lo .sdA	
	riemper TIA 10	
мите С. Т.	Temperate Value	
	Heigh:	ee 5 4 4 1 2 4 2 4 2 4 2 4 4 4 4 4 4 4 4 4 4
02	Nitrates	$\frac{x^2 - x^2}{x^2} = \frac{x^2}{x^2} = \frac{x^2}{x$
NITRO- GEN A	sotiutiV	8544 : 825 25 25 25 25 25 25 25 25 25 25 25 25 2
GEN.	papuad -sus	
ORGANIC NITROGEN.	Dis	
	pnd'd s	8112232888889112122221111112222
AS A.	moid Am solv'd	
NITROGEN AS	Albuminoid Pis-sid b'vlos	
NITIN	- 1'10T	2
	FreeAm monia.	0.8.3. 23.8.2.8.2.8.5.8.1.8.1.8.1.8.1.8.1.8.1.8.1.8.1.8.1
GEN JMED.	BySuspd Alatter	
OXYGEN	Total.	0x + 0 = 0 = 0 = 0 = 0 = 0 = 0 = 0 = 0 = 0
ne.	СһЛогіл	1.1.8.1.8.9.9.0.1.1.3.1.3.0.0.2.2.3.0.0.2.4.0.3.0.2.2.3.1.3.1.3.0.0.2.2.3.0.0.2.4.0.3.0.2.2.3.1.3.1.3.1.3.1.3.1.3.1.3.1.3.1.3.1
	popuod •sng	8,000,800,79,80,82,50,50,50,82,83,84,83,84,50 1
RESIDUE ON EVAPORATION.	solved.	1
RESIT	.Into'l' -si(I	
	Odor	
.:	Color.	$\frac{N}{\omega \omega \omega_{ij} \omega_{ij$
APPEARANCE.	Sedi- ment.	V. Little Little V. Little V. Little Much Cons.d Little Much Little Litt
A.P.	Turb'y.	Much Decided Decided Much Decided Much Decided Decided
OF	1900 Exami- nation.	
DATE OF	ė.	
H	Collection.	Apr.
.0%	Rerlat	18

TABLE 133.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, HENRY.

Report of Edwin O. Jordan,	University of Chicago.
LION.	

·+	r., <u>1</u>	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
No. o Bac.	Cubic Centi meter	9,400 1,000 1,000 19,40
	Presen	
	TeqmeT riA To	
	Temper	000400000000+++555-555555555555555555555
	Heigh Mate	
NITRO- GEN AS	Nitrates	$\frac{z-1}{z^2} = \frac{z-1}{z^2} = \frac{1}{z^2} = $
NIT	Nitrites	8.58 : 12.88.58.58.58.58.58.58.58.58.58.58.58.58.
IC EN.	-sng	
ORGANIC NITROGEN.	Dis-	
ŌZ ——	Total.	
20	E Pud -sng	98 98 98 98 98 98 98 98 98 98 98 98 98 9
AMMONIA.	Pionimold Pis- Dis- Div'd	- 1
NITHOGEN AS	Totyl	83-14688644488448344448848484848484848484848
	Free Am-	588 × 8 + 5 2 6 2 5 8 + 4 + 4 4 4 4 4 4 4 5 8 8 2 4 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5 1 5
N ED.	BySuspd Matter.	000-00
OXYGEN CONSUMED.	By Dis-	
Ççç	Total.	<u> </u>
.əı	Chlorin	81 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
ON.	pepuad eng-	
RESIDUE ON EVAPORATION.	Dis-	######################################
RES EVAI	Total.	88888888488488888888888888888888888888
	Odor	
	Color.	ej w pg w w w = + + pj + o o o o + + pj w pg + + pg + w w o o o o
APPEARANCE.	Sedi- ment.	Little Cons'd Little Cons'd Little Cons'd Much Little Cons'd Much Little Cons'd Little
APPE	Turb'y.	Decided V Slight Distinct Nuch
OF	1	+0-22-0282-2822-0522-05330053
DATE OF	1900 1900 Collec-Examition.	<u> </u>
.o.V	1 0	1170 Jan. 1170 J
1		

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-ILLINOIS RIVER, AVERYVILLE.

. *
17
7.
4
-74
0
)m(
p.a
jenneg
C
-
-
_
\circ
_
2.00
7.
person
ine.
-
—
6.3
ايشا
مسه
-
O.
0
rt o
rt o
ort o
ort o
port o
port o
eport o
Seport o
eport o
Seport o
Seport o
Seport o
Seport o
Seport o
Seport o
Seport o
Seport o
Seport o

No. of Bac.	per Cubic Centi- meter.	23.60 24.60 25	
	Presend No. sd A		
	Temper		
	Temperate Nate	- नार्यक्ष च - च - च च च - च च च च च च च च च च च	
	Helgh otaW	#####################################	
NITRO- GEN AS	Nitrates	- + 8 + 4 + 6 + 5 + 5 + 5 + 5 + 6 + 5 + 6 + 5 + 6 + 6	
	Nitrites	<u> </u>	
ORGANIC NITROGEN.	solved.		
ORG. NITR	Total.		
	Sus- prd'd	25.50.00	
SN AS NIA.	Albuminoid Pis- Dis- Solv'd	882888282828255555828365588255825	
NITROGEN AS AMMONIA.	I'10T'	\$ 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
Z	FreeAm- monia.	8888448888888468855 885-8858	
- <u>0</u>	Bysuspd Matter.		
OXYGEN Consumed.	By Dis- solved.		
000	Total.	<u>ααι-αααρ-ααι-αμι-ααι-ι-ααι-ι-υ-ι-υ</u> <u>ααι-αααρ-ααι-αμι-ααι-ι-υ-ι-υ</u>	
16,	Chlorit	<u> </u>	
ON.	peptied Sus-	表表表現中央學習其中的理論中的學家表示中的學習表示中的中	
RESIDUE ON EVAPORATION	Dis-	88888888888888888888888888888888888888	
RES	Total.	88888888888888888888888888888888888888	
	Odor		
	Color.	၍ သံလံသည့် ညှ က် → သို့ ယံ ႏွင့် ဆွဲ ၍ လို့ လို့ ယံ ညှိ ယံ သို့ → — သို့ ညှိ လံ တံ ၍	
APPEARANCE.	Sedi- ment.	Consideration of the considera	
API	Turb'y.	Distinct Slight Distinct Much	
OF	Exami- nation.	Na	
DATE OF	1900 1900 Collec-Exami- tion. nation.	Jan. 2 Jan. 156b. 6 Seb. 13 Nar. 17 Mar. 17 Mar. 17 Mar. 17 Mar. 17 Mar. 18 May. 1 May	
.oN	Iniroz	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, WESLEY CITY.

Report of Edwin O. Johnan. University of Chicago.

No. of Bac.	per Cubic Centi- meter.	10,000 23,000 47,000 31,000 11,000 15,000 15,000 16,000 16,000 110,000 110,000 110,000 110,000 110,000 110,000 110,000 110,000 110,000 110,000 110,000	1.400
	Presen for sof A		:
	request riv to		30°.
ature T. C.	Temper	<u>ត្ត</u>	-
	Heigh Wate	9-3-4-4-4-3-3-3-3-3-3-4-3-4-3-4-3-4-3-4-	
NITRO- GEN AS	Setantiv		- co
NITRO	Nitrites	8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	. Tes
IIC REN.	-sng		:
ORGANIC NITROGEN.	Dis- solved.		:
ON	Total.		:
on l	p,pud	\$38.50	1 01.
TROGEN AS	Pionimonian Pionim	4288874887888874888 2884589	01.
NITROGEN AMMONI	T'10T'	<u> </u>	
	FreeAm-	\$ 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	+
EN (ED.	BySuspd Matter.		
OXYGEN CONSUMED.	By Dis- solved.	antente and and anterted trustical	9.6
ဝဂ္ဂ	Total.	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	6.3
ne.	Chlorin	814. 77. 1. 2. 2. 2. 4. 4. 7. 4. 2. 2. 2. 2. 2. 2. 2. 4. 7. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	17.6
ON ION.	pepued -sng		14.
RESIDUE ON EVAPORATION	Dis-		202
RE Ev.	Total.		2.0x
	Odor		:
	Color.	ည်း သည်သည်: ု နယ _{ုင္} မွာလက်နနည်းမှုန္တန်နနည်းလိုလ်လယ်ယံတို့ သည်သည်း ကောင်းမှုနှင့် မြောင်းမှုနှင့် မြောင်းမှုနှင့်လိုလ်လယ်ယံတို့	G: -
APPEARANCE.	Sedi- ment.	Much Cons'd Much Cons'd Much Cons'd Much Cittle Cons'd Much Cittle Cons'd Much Cittle Cons'd Little Cons'd Cons'd Little Cons'd	Cons a
AP	Turb'y.	Decided Nuch Decided Decided Much Decided Much Decided Distinct	
)F	1900 Exami- nation.		22
DATE OF	c- Ex	\$25.33.36.59.50.24.50.25.65.35.50.35	-
D.	1900 Collec- I	Jan. Feb. Apr. Mar. May. Line June	
Serial No.		1190 1232 1232 1232 1232 1232 1232 1233 1	3043

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATTER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, PEKIN,

Report of Edwin O. Jordan, University of Chicago.

No. of Bac.	per Cubic Centi- meter.	70.000 530.000 55.600 51.00
	Present lo ,sd A	
	TeqmeT TiA lo	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	riogmoT of BW 10	
	Helgh Mate	-01-4-4-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-
RO-	Setrativ	
NITRO- GEN A	Sitrites	26.00 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
IC REN.	papuad -sns	
ORGANIC NITROGEN.	Dis- solved.	
ÖZ	Total.	
	E P, pud	6 2 2 3 2 3 2 3 2 3 3 3 3 3 3 3 3 3 3 3
ITROGEN AS	T'01'T Albuminoid Dis-	\$\frac{1}{2}\frac{1}\frac{1}{2}\f
NITROGEN AS	Totrl	4818 - 481
	FreeAm- monia.	20.00 - 0
N ED.	BySuspd Matter.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
OXYGEN CONSUMED.	By Dis-	ဆု-သည္။ ဆု-သူ။ ဆု၊ ဆု၊ ဆု-သည္
000	Total.	$\frac{1-\widetilde{\omega}}{4}\widetilde{\omega}_{0}+\frac{\widetilde{\omega}}{4$
.9r	Chlorin	(2) 23 23 24 25 25 25 25 25 25 25 25 25 25 25 25 25
ON ION.	papuad -sus	→ E Y - E W F F F F F F F F F F F F F F F F F F
RESIDUE ON EVAPORATION.	Dls.	88 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
RE	Total.	21 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -
	Odor	
	Color.	who was a second
APPEARANCE.	Sedi- ment.	Consider Little Little Much Consider Little Consider Consider Consider Little L
AP	Turb'y.	Much "" "" Decided Much Decided Much Decided
DATE OF	ec-Exami- nation.	2
D	tion.	Tan. Mar. Way
No.	berial	2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

TABLE 137.

STREAMS ENAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

Sounce o

Y
-
7.
- 1
200
-
-5
Ξ
ER,
200
5.7
pales
200
E
NOIS
-
0
1
94
T
=
. 3
-
\blacksquare
- 1
5-3
F
TER-
AT
A
VAT.
A
W.A
W.A
W.A
A

University of Chicago.

Report of EDWIN O. JOHDAN,

No. of Bac.	27,600	63,000	46.600	000,09	50,000	4.700	4.000	9.900	1.100	40,000	11,200	31,000	20,000	19,000	10,500	7,400	002,01		
ce or		:	:		•	•	:			:			:	:	:	:	:		
ature, C.	19 11	qmoT A lo		-15	6	6	₹~	22	=	<u>8</u>		-3	36.	63	66	26.	22 23	31.	:e:
ature D, r.	er er	qm9T sW lo	0.	0.		د ه	9	6	=	11.5	S	- L	24.	- T	27.	.54	25.	.36.	27.
lo i	911 .q2	Bi9H SW	4.06	-1-	7.7	17.	14.9	14.3	12.9	13.	11.2	10.3	9.1	9.	8.4	8.	 	∞	6.5
NITRO- GEN AS	sə	Nitrat							_										ಹ
	!	Nitrite	.046	330	100.	800.	910.	.032	.031	160	33	3	:063	360.	3	91.	2.5.	.13	1.1.
NIC GEN.	_	-sus	:	:	:	:	:	:	:	:	-:	:	:	:	:	:	:	:	:
ORGANIC NITROGEN.	-	-si U -solve		:	:		:	<u>:</u>	:	:	:	:	:	:	:	:	:	:	:
		Total		:	on	<u> </u>	.: 9	:	:	· ·	·:	:	:	:	:		20	9	:
90	id Am	-sng	80.	81.	. 16	. 16	910.	₹ -	80.	<u> </u>	76.	:	7.	.13	7-	<u></u>	<u> </u>	. 13	80.
NITROGEN AS AMMONIA.	Albuminoid	Dis-	1				.176	_			_	:	.248	808	. 192	. 168	. 168	.208	ું.
NITRO	Alb	Tot'l					.192					٠	_	_		_			85.
		FreeAr inom	1.33	3.21	1.8	.464	7.	 	. 168	= :	. 144		408	.408	.341	7.	.288	.344	.36
ED.	r. bd	Bysusi	. i.	6.	1 6	€. ∞	1.5	₹.	?? 	50.	1.9	.5	-1		6:	1.8	9.	70	9.1
OXYGEN CONSUMED.	p	siU va solve	ت.			10	rc. oc	5	ગ		9	9	9	<u> </u>	1-		6	6.	9
03	.1	Total	6.3				7.3		<u>ښ</u>		$\dot{\infty}$	∞			3-	2 7 .5	1-	∞	
.9r	ii	СРГО	.12	21.		6.	4.6	re,					<u></u>	23					17.
ON ION.	pa	sng	3.	13.	59.	131.	34.	59.	54.	5	18.	<u>[</u>	200	103.	63	φ.	æ	.99	£3.
RESIDUE ON EVAPORATION		Solve	352.	306	285	131,	180	197.	218	219.	313	333	262.	127. 17.	368	.368	33.5	244.	. 866
REEVA	٦.	rio'l'	355.	318.	314.	262.	214.	256.	272.	234.	230.	328	: : : : : : :	376.	330.	346.	337.	310.	293.
	Odor					•							:						
		Color.	9.	33.	Muddy	े चं	7	io.	કરે	ಬ	ಣ	က္	7	જ	υį	ಣ	ः	٠ <u>٠</u>	55.
APPEARANCE.		Scdi- ment.	Little	:	:	Much	Little	:	:	:	:	*	:	*	4	Consid	Little	3.9	Cons'd
17:		Turb'y.		_			3,9		9.7	Decided	"		-	Much		;	:	79	93
OF	1000	Exami- nation.			30		Apr. 6		6:1	96 ,,	ಉ		17	:: 31	E 33	June 7		: 31	; ??
DATE OF	1900 Collec- Extion.		=	30	2	8	Apr. 4	=	$\frac{\infty}{2}$	155	≎३	<u>6</u> .	91	: 33	31	June 6		06	27
No.	al	Seri	1231	1314	1353	1555	1612	1651	1691	1730	1769	1801	1838	187.1	190-1	1936	1971	2010	2014

STREAMS ENAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of EDWIN O. JORDAN,	University of Chica
SANITARY WATER ANALYSIS-PARTS PER MILLION.	SOURCE OF WATER-SANGAMON RIVER, CHANDLERVILLE.

No. of Bac.	Cubic	Centi- meter.	006	000.1	7.300	15,700	000	29.300	117.000	70.000	45,(10)	50,000	120.000	30.00	18,500	7.000	5.600	3,(500)	1.900	16,900	2,700	8,000	11.3(8)	12.000	8.300	×.83.5	11,600
Presence or Abs. of Coli.						:		:	:		:				:		:	•	•								:
		TempT	20.	© }	·-	15.5	5.	~~	7		x	-	-	6.	=	33.5	19.	15	••••	55	3.	33	57.	30.	30.		30.
		Temp W 10	0		- ·	.c	0.	ei	0.					_		7,070				_		_			5.5		
10 1	oju Zpi		esi	E.	ين ن:	4	· ·	5.	·-	::	1-	16.	21	x	σċ	.9		***	575	٠÷	35	~	000	7	5.		<u></u>
RO. N AS	sə	Nitrat	2.65	1.9	- CF.	.T.	∞ •	1.8 8	1.35	1.3	£.50	1.15	.8. .8.	2.6	. 8:	- 68	- 2	£.		9.	1.05	 	1.3%	15.	£.	1.5	.cs
NITRO-GEN A	sə	Nitrit	.016	800.	800.	10.	5	<u></u>	£30.	810.	.015	.012	910.	01710.	910:	???	10.	0.	800.	0.	800.	10.	980	33:0.	.048	100.	×20.
OEN.	_	-sus		:	:	:	:	:	:	:	:	:	:	:	:	:			:	:	:	:	:		:	:	:
ORGANIC NITROGEN.		SiU 97fos		:	:	:	:	:	:	:	:		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
0%	1	Total		:	:	:	:	:	:	:	:	:		•			:	:	:	:	:	:	:	:	:	:	:
	1 Am	p,pud -sus	.048	.024	.053	.064	150.	.073	1.13	. 144	.199	.392	£	880.	570.	<u>0</u>	.048	-	10.	:	. 152	. 232	1000	F (2)	91.	<u> </u>	.168
SEN AS	Albuminoid	olv'd Solv'd	. 136	.088	880.	x	XX0.	++1.	.16	. 13x	XX.0.	101.	19.	880.	990	. 136	.0.96	£.	144		. 136	<u>3</u> :	101	.13	136	<u>:</u>	<u>=</u>
NITROGEN AS	Albu	Tot'l	381	.113	==	361	<u>:</u>	316	1.38	37.5	×.	.496	:26:	. 176	. 13x	.176	=	155	- X	:	SXX	.353	878:	.34:1	296	33	888
		FreeAn		. 168												180.	+20:	10.	(F)	:	860.	:01:	.034	33	(F.S.)	:013	.01
ED.		Brsus	ıc	≎}	_:	7.	٥!	??	19.3	G:	* •	0.7	10.3 2.0	<u></u>	 	9.	寸.	10	<u>.5</u>	83 6	* :	55	6.4	4.9	X.	5:	5.1
OXYGEN CONSUMED.		By Dis		2.4														3.5							寸. 寸	-	-:-
Cos	1	Tota	7	2.6	3X	-	2.6	x.	83	+	?? ∞	14.3	<u>x</u>	6.	÷		8.3	10	ۇي	6.9	6.	10 30	oc ici	≈ ∝	3.6	≈ ∞	9.1
,9r	ιμ	СРІО	5.3	6.4	<u>ت</u>	30 00	10.	۳. دن	د ه	,	٥١ <u>.</u>	÷	က္	χ. જ:	∞ ∞	rc ej	9.+	ος 65	4 ∞	.c.	6.4	-	5.6	4.6	8.	က်	6.
o.N.	po	puod sus	0.5	14.	16.	51.	2.	300	704.	œ œ	184.	300.	334.	:01:	33	<u>∞</u>	33	.99	-	- 2	ž	224	x 000	202.	173.	180	328.
RESIDUE ON EVAPORATION.		si(l ovfos	348.	200.	200.	307	350.	.000	<u>S</u>	230.	226.	 	· £	- in	276.	282.	- 55	7.7.	205	266.	.505	244	285	25:3.	211.	257.	238.
RES EVAI	ĵ.	stoT'	350.	331.	:306:	350	3332	346.		31x	410.	338	520	355.	358.	330.	347.	340	3:36.	150.	:70.	168	303	155.	414.	456.	566.
	Odor			:			:			:			:		:	:				:							
		Color.	<u></u>	-:	~:	• ?	 	•	: :	·	9	•			· ·		•		?? ??	· · ·	•	~	٠ <u>.</u>	· 55.	: ::	÷.	٠ <u>.</u>
CE.		5	0	41			_				_	_						_	_	_	_			_			
APPEARANCE.		Sedi- ment	V. Litt	Little	1,1	* 9	,,	19	Much	f'ons'd	9 9	Much	9 9	Little	*,	:	p,suo,)	Little	;	p.suo.)	Little	Consid	9.9	99	* 9	Little	9 9
AF		Turb'y.	Decided	:	*	,,	Slight	Much	* *	:	:	:	:	:		Decided	Much	Decided	:	Much	Decided	Much	3.0	9.9	17		*
OF	1900	Tollec- Examition.	an. 4	11	61	÷	'eb. 3			dar. 2		:		0 2	1 jr. 6		5:		lay 3					une 7	-	- 5	×21
DATE OF	_	- Sc -	3.lan.	10	13	77	-	į		1	1-		77	X ?	1.	-	<i>3</i> .	5	2 1	10	16	~	30	6.11	20		7.
0	1983	Colle tion.	Jan.	:	:	:		1. 6. 1.		Mar.					Apr.				Mai		9 9		,,	June	9.9	99	9 9
No.	[11]	Seri	117.5	1315	1350	1276	1315	1355	· · · · · · · · · · · · · · · · · · ·	7	1170	EX.	154	1573	1613	1655	16.92	17:31	17:0	18.50 18.50	18:30	1875	11911	1945	1981	105	285

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, BEARDSTOWN.

EDWIN O. JORDAN,	University of Chicago.
O.	
Report of	

No. of Bac.	per Cubic Centi- meter.	10.700 17.000 17.000 17.000 17.000 18.700	
	Presenc		
	rempers		•
	riadmaT ataW lo	000000000000000000000000000000000000000	
	HelghH otaW		
RO-	Xitrates	· · · · · · · · · · · · · · · · · · ·	
NITRO-GEN A	Nitrites	9.00.00.00.00.00.00.00.00.00.00.00.00.00	
NIC OGEN.	solved.		
ORGANIC NITROGEN.	Total.		
	p,pud	0.088	
EN AS NIA.	Piori Pistoria di Piori di Pistoria di Pis	25.50	
NITROGEN AS	T'10T	23.25.25.25.25.25.25.25.25.25.25.25.25.25.	
N	FreeAm.	1	
ED.	Bysuspd Bysuspd	13 15 16 17 <td< td=""><td></td></td<>	
OXYGEN CONSUMED.	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
Con	Total.		
.er	Chlorin	%;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;	
ON ION.	pepued sns-	26.5. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.	
RESIDUE ON EVAPORATION	Dis-	8 2 3 3 8 8 3 1 1 2 2 2 1 1 2 3 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	
RE: Eva	Total.	23.33.33.33.33.33.33.33.33.33.33.33.33.3	
	Odor		
	Color.	وي و إ دا ه و ي و يو نه مد تر ـ مد دا سه مد سه مد مد نو مد نو مد دن ده نوي .	
APPEARANCE.	Sedi- ment.	V. Little Liktle Much Cons'd Much Much Much Cons'd Much Much Cons'd Much Much Much Much Much Much Much Much	
AP	Turb'y.	Slight Much "" "" "" "" "" "" "" "" "" "" "" "" ""	
DATE OF	1900 Exami- nation.	88	
DAT	1900 Collection.	Nan. Nan. Nan. Nan. Nan. Nan. Nan. Nan.	
.oV	Serial	12864 1536 1536 1536 1536 1536 1536 1536 1536	

TABLE 140.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-LLINOIS RIVER, KAMPSVILLE.

Report of EDWIN O. JOHDAN,

of C.	per Cubic Centi- meter.	2000 2000 2000 2000 2000 2000 2000 200
No.	D Corre	1-021-62226
	Presen To .sdA	
	toqmoT il/. lo	<u> </u>
	request of Walt	6 8 4 9 8 - 0 % 9 % 6 % 10 % 1
	Heigh United	
NITRO- GEN AS	Solventin	
	Nitrites	2000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
NIC OGEN.	solved.	
ORGANIC NITROGEN.	Total.	
	p,pud -sns	25.5.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.
EN AS NIA.	Albuminoid 1.30T	86888888888888888888888888888888888888
NITROGEN AS	Tot'l	4 # # # # # # # # # # # # # # # # # # #
	FreeAm' monia.	2 3 3 4 6 6 7 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
EN.	Bysuspd Matter.	
ONYGEN CONSUMED.	By Dis-	Q
30	Total.	<u> </u>
.91	Chlorin	© 8 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
ON FION.	popuod -sus	2
RESIDUE ON EVAPORATION	Dis- solved.	
By.	Total.	\$\\ \text{8.2} \\ \text{8.2} \
	Odor	
	Color.	· · · · · · · · · · · · · · · · · · ·
APPEALIANCE.	Sedi- ment.	V. Little Little Much Little Cons'd " " Cons'd " " Much " " " " " " " Much " " " " " " " " " " " " " " " " " " "
AP	Turb's.	Decided Slight Decided Mich Mich Control of
OF	1900 Exami- nation.	Jan. 15 19 19 19 19 19 19 19 19 19 19 19 19 19
DATE	1900 Collect F	Jan. 31.7 17. 25. 17. 17. 17. 17. 17. 17. 17. 17. 17. 17
.o.X	Iniro8	1171 58 1216 5

TABLE 141.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON.

University of Chicago.

Report of Isbwin O. Jordan,

No. of Bac.	per Cubic Centi- meter.	8.8.8.6.00 9.8.8.6.00 9.8.6.00 9.8.6.00 9.6.000 9.6.
	Presenc to .sdA	
	Tempera	<u> </u>
ature D, T	Tempera	00,40,000,40,40,1,40,8,8,8,8,8,8,8,8
	Heigh otsv	<u></u>
NITRO- GEN AS	səts:riiN	
EIN E	Nitrites	84999999999999999999999999999999999999
CEN.	-sns	
ORGANIC NITROGEN.	Dis- solved.	
	Sus- bibid bibid	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
N AS		200
NITROGEN AS AMMONIA.	Albuminoid -sid b'vlos	88888888888888888888888888888888888888
N	.ainom	4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
	Bysuspd Matter	
GEN	bevios	
OXYGEN CONSUMED.	Total.	$\alpha\alpha\alpha\alpha\omega$
.er	Chlorit	8 12 4 4 4 4 4 4 8 8 9 9 1 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9
NO.	pəpuəd -sng	4.6. ± 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8. 8.
RESIDUE ON EVAPORATION.	Dis- solved.	88.58.98.88.59.59.59.88.88.88.88.88.88.88.88.88.89.89.89.89
RES EVAF	Trotal.	4 2 3 3 3 3 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5
	Odor	
**1	Color.	
APPEARANCE	Sedi- ment.	Little Cons'd "" Cons'd "" Little Cons'd ""
ΛP	Turb'y.	Nuch Slight Much Decided Much
OF	1900 Exami- nation.	Nay 36 (1987) 1987 1988 1988 1988 1988 1988 1988 1988
DATE	1900 Jollec- E tion. na	+5742-742-484-588-05886-5888
		2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
No.	Serial	1861 1861 1886 1886 1886 1886 1886 1886

TABLE 142.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON.

University of Chicago.

Report of EDWIN O. JORDAN,

of .:	Ti i	er.	300	. 100	3,900	19,200	100	0003	3,000	0000	0.000	000.	0.00.0	0.000	3,000	5.0(10)	005.0	.500	3.800	3.100	3,500	3.700	9.600	000	3.400	000'1	•
No. Isag	Cubic Centi-	met		21	00		2 {-	33	33	3	£	170	3 -	14		152	7	_		- t						_	
	resend s. of			:	:	:			:			:	:	:	:	:	:			:	:	:					:
	mper Tiv l		6.	e:	2	- 0	: <u>cc</u>	ទាំ	10	25	-	÷	17.	14.55	e i	13.	21.	G		20 20	500	27.	£	31.	863	30	
eture O (re	niperi Wate	9'l' 10	0.	0	₩.	÷ <		0	0.	0		<u>ښ</u>		ī.	ж —		15.	-19	19.	233	19.	152	25.	.96	26.	61	
	leigh State	1	2.9	£.5	10 t	. o) 00 0 00	6.6	30 60	9.3	16.3	14.	12.7	11.8	13.1	10.9	12.7	10.7	5. 5.	 	<u>≂</u>	6.5	e	ات ده	33.	rc 1-	
NITRO- GEN AS	eeja1	IIN	0.	-:	: (uj a	18	0.	.55	.05	· +:	-:	19	. 15	9.	-	3	٠. ن	0.	-:	55	ું. કું.	:S:	0.	.85	eć.	
	bəbn		0:	0:		3000	88	014	800:	<u> </u>	900:	900:	0.	011	018	.018	018	30:	0:	× 00.		016	.CO.	0:	0:	89. -:	:
ORGANIC NITROGEN.	SIS. Jved.	os		:		:		:		:	:	:	:	-:	:		:	:		:	:	:	:		:	:	:
ORC	tal.			:	:	:			:		:	:	:		:	:	:		:	•	:		:		:	:	
an	A Pu	oud oud	.144	कें:	हें। इंटें	1980	180	.392	793:	<u>\$</u>	1.064	89.	373	% ₹ ₹	.856	30.	£	<u></u>		146.	98: -	<u>:</u>	3;	880.	.13	. ±35	
NITROGEN AS AMMONIA.	Albuminoid is-v'd	IOS	.208	.176	.176	82	#	255	300	.156	.216	.221	.216	. 58	:361	22	25.	. 136		.176	.16	9	.136	.16	. 136	<u>.</u>	
NITRO	I'I	оТ	.352	.416	.416	# S	328	624	.56	919.	×	1 06.	<u>∞</u> .	911.	<u>:</u> :	.301	93:	23			£	X.	.336	<u>∞</u>	.256	.56	
	-mkə-					. IS										_			:	_	_				1020	_	
GEN IMED.	ensbq lags lags lag	1375		_	- 0	ю ос э — э е	=	10	9							C	_		-		- i				4 1.6	_	:
OXYGEN CONSUMED.	Jajo Leid	BY	9	9. 7.	9	210	8.3	600	÷	9	0:	-	_	_	9	- 1	-	a.	=	XC :	x	33.	Ŧ,	_	9.	φ_	
'ət	rirofd	G	_	4.67		· +		- - - - - - -		-						25 - 20 -									3.5		
×. C	pəpu -sng	bel E			٠.	- - - - - - - - - - - - - - - - - - -	64.	324.	119.	687.	350.	416.	226.	55	616.	130	£30	X.	200	217	.6.	- 80	 	-6.2	œ.	113.	•
RESIDUE ON EVAPORATION.	sic.				_	000		_	_		=		-														
RESI	Jero	o'I'	:	231.	197.	2000. 0-150.	262.	510.	330	739	 	5-10.	350.	370.	33			323			210	- O.S.	.300)	535	210.		
	Odor		:					:		:	:	:	:			:	:	:	:	:		:			:	•	:
			- :-		: :	6. 50	<u> </u>	155	5		idy	ldy	:	•			:	0	:	:	:	:	:	:	:	: 	
E2	Color.							_			_					_				_	-				es !		:
APPEARANCE	Sedi-		Little	:	99	Little	33	Much	9 9	uš Lp	9 49	1.0	ug ug	40 1	9 9	Consid	Much	Little	Consid		Much	Cons.d	al a	Little	Cons'd	Much	
11.	Turb'y.		Much	:	: :	:	:	*	:	*		**	•	*	;	:	:	4	4		:		:	:	:	;	
F4 (1900 cami-	tion.	n. 5	11	61 .	8 °	X	10								5			2				HILL X		() () () () () () () () () ()	₹; •	:
DATE OF	1900 1900 Collec- Exami-	n. na	+Jan	10	- 2	21 15	(-	=		{-		<u>₹</u>		4		X		S Mis				30	6 6.1	1:3		6.2	
1	Co.13	tio	8 Jan.		: :		r Feb.		;; G					6 Apr.					: :			23	-		0	:	
.0%	[BlT9	S	20	121	200	3	133	1:3	110	1+1	100	153	200	161	165	163	1	1-1	3	X	100	191	1:41	14.	2019	502	

TABLE 143.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM ILLINOIS SHORE, ALTON.

A.N.,	Chicago.
JORDAN	of (
	ity
sport of Edwin O.	University
70	
11011	

No. of Bac.	per Cubic Centi- meter.	1,500	200 200 200 200 200 200 200 200 200 200	21,800	17.000	85,000	59,000	175.000	55,000	50,000	17,000	15,000	18,000	7.300	10,700	4,400	2,000	11,600	6,500	13,500	3,600	7,600
	Presen lo .ed						:				:			•	:				:		:	:
	raqma riv ro	. 4.	o <u>c</u>	0,	50	0			:	:	10.	ر 2 م	158	233.	16.	.65	21.	33.	26.	36.	<u>કુ</u>	65
ature or, C.	ragma' atsV/ 1	0.0							:	.9	 o: o	x c	15.	20.	19.	3.1.	19.	24.	24.	24.	23.	.98
	Heigh Mate	0.	رة الإن الإن الإن الإن الإن الإن الإن الإن الإن الإن	0.0	2 .0	70.0	÷⊆		14.1	11.5	1.0;	1.4 5.5	6.11	10.1	11.4	∞ ∞.	~ ∞	2.5	6.1	ت وغ	 	7:3
30- N A S	itrates	<u>.</u>	-ر ب	.c.	4. ç.i	1-1	ر بن با	7	∞.	, .			=		∞	.55	:£	9.	1.15	ıc		
NITRO- GEN A	setinii	80.	0.013	.01 .00 .00 .00 .00 .00 .00 .00 .00 .00	5.8 8.8	.012	85.	.018	800.	10:	910.	920	960	.022	810.	120.	-03 1	ි ව	900.	.018	.01	.034
JEN.	papua-		: :	:	: :	:	:		:	:	:			:	:	:	:	:	:	:	:	
ORGANIC NITROGEN.	Dis-		: :		: :	:	:	:	:	:	:			:	:	:	:	:	:	:	:	
oz —	rotal.	1:		:		:	: ;		:	:	:			:	:	:	:	:	:	:	:	:
	-sns	80.	# 88 	.072	1.248	19:	5.55	.856	558	.328	.192	. 136	201	.328	:	. 232	ં	.104	.13	80F.	.07.	916.
TROGEN AS	Albuminoid -siq	232	5.80 80 80 80 80	63.	308	184	200 200 200 200 200 200 200 200 200 200	104	91.	.128	128	38.	. 128	. 136	•	. 192	. 192	.176	ં	.16	.192	.16
NITROGEN AS AMMONIA.	[K 1.10]	.312	.528	272	1.456	.83. 183.	656	8	.74	.456		272	352	191		+24	365	82.	.35	.568	.36±	.376
	reeAm- monia.	976	22.80	.92	010: 82:	.44	368	88:	. 256	- 255 - 255	36.5	85	.044	.036		.004	.0 .	900.	.034	870.	3	.034
N ED.	ysuspd	0.0	ν.		6.6	10.7	0 00		{- }-	8.6	က ၈) (20 co	3.1	٠c.	5.8	2.1	63 63	e.;	<i>∞</i> ∞	ાં	ന	
OXYGEN CONSUMED.	y Dis-	00.0	25 00	00.1	7. 7.	9.	2 7	1 .		9	∞	5 is	∞	∞	1C	∞	35	7	6:		7	∞
CON	lotal.	∞.	10.8	∞ °	0 TO	17.3	× ∞ ∞	16.	13.7	14.2	x 0	x ∞ 0 ∞	6.8	8.01	12.3	8.9	S. 53	8.4	~ í	ŵ	7 ! ∞	8.01
.9u	Chlori	15.9		25.5	4.6	6.1	ο Ω 1Ω		7.7	4.6		a. 4	8:	8.0	7.5	œ	∞ ∞	رة زن	-7 55	8.6	œ.	9.9
on.	pəpuə -sns	8.58	38°5	0.00	1334.	753.	. 600 1000 1000	1218.	839.	374.	300.	184	213.	310.	.133.	178.	265.	160	196.	516.	122.	327.
RESIDUE ON EVAPORATION	Dis- olved.			258.					_													
RES! EVAP	Total.	332.	440.	338	- 75t-	943.	55.6	350.	976.	534.	470.	388 380 380 380 380 380 380 380 380 380	405.	541.	629.	430.	505	384.	420.	730.	342.	499.
	Odor			:		:	:		:	:	:			:	:	:		:	:	:		:
		1:		:	· · ·	:	:		<u>:</u>	· ·	:	: :		:	:	_:	:	:	:	:	:	-
	Color.	85.	Mudd	6.5	Pinda	G) 6	3 =	123	.15	Mudd	ين. د	4.00	. 23	ೞ	₹.	٠.	ું.	ಲ	ಀ	wi	ej.	.25
APPEARANCE	Sedi- ment,	Little	Munca 	Little	Much	61	Much Much	7,7	*	*	* :	: 3	Cons'd	3,9	,	7 4	Much	Cons'd	9 0	Much	Little	Much
AP	Turb'y.	Much		; ;	: 3	9.9	. 3	7,9	",	3	3 3	: 3		Much		*,	,,,	"	9 9	**	;	27
OF	1900 Exami- nation.			Feb. 3			Mar, 3	15	22		Apr. 6			May 4		" 17	75 ,,		June 7	14	3	86
DATE OF	1900 1900 Collec- Examition.	Jan. 4 Jan.	2 8	Feb. 2 I	14	ह्य	1 1	;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;;	25	28	Apr. 4 2		25.		6	91 ,,	233	30	une 6J	" 13	02	<u> </u>
No.	Serial			1321 1321				1503	1539							1833	1868		-	1976	3013	2045

TABLE 144.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM ILLINOIS SHORE, ALTON.

Report of EDWIN O. JORDAN.

o. of Bac.	per Cubic Centi- meter.	1 300	8.200 8.200	93,500	32.800	41.000	143,000	135,000	51,000	80,000	110,000	150,000	110,000	35.000	67,000	15,000	38,000	13,500	4.600	1000	5,700	8000	2000	10.000	35.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00	1,000
Coll.	Presence of the Presence of th													:												
	Temperative To	-	- 0	.01	0	10.	0	-	0	6	9			10.	57.	9	33.	£	16.	65	÷ 23	25	.96	96.	183	30.
	remperation of Male	-			0		0.	0	Ċ.	· ·	+	•	.9	6.	эc	10.	5.	.06	19.	21.	16.	. 45	.51	24	33	36.
	idgiəH əts7/	-	. T	0.0	0	9.8	5.	5.5	20 4.	10.	17.	4.4.	6.11	10.1	11.4	10.5	6.11	10.1	1.1	∞ ∞	8.4	£-;	6.1	5. 63.	50	63.
10-	Sitrates	-	1.22			_			-	_	_		_			_	_			-						
NITRO	Nitrites	010	885	010	810.	800.	.007	015	300	0	800.	900.	10.	910.	3	.03	.055	800.	10.	1.20.	.016	10:	004	110	0	970
IC EN.	papuad -sng				:	•		:	:	:	:	:	:	:	:	:	:	:	:	:		:	:	:		
ORGANIC NITROOEN.	Dis- solved.		•		:	•	:	•	:	:	:	:	:	:	•	:		:	:	:	:		:	:		
5Z	Total.				:	:	:	:	:			:	:	:		:	:	:	:	:	:	:	:	:		
	p,pud -sng	18	150	333	.152	.088	.944	1.	.476	504	888.	3.5	358	<u>.</u>	.432	. 136	199.	.33		198.	+ ???	.136	803.	. 192	111	200
SEN AS	Albuminoid Pis-sid Divios	32.1	2000	180	ा	316	.176	122.	199	. 159	=======================================	803	864.	91.	S	. 136	.176	<u> </u>	:	.176	181	91.	. 136	. 16	136	=
NITROGEN A	Tot'l	226	200	504	352	305	1.12	199	368	.056	1.008	1.0%	.456	∞ ??	3.	3733	Œ.	++.		#.	.108	(C)	.341	355	086	.376
A	FreeAm" monia.	12.4	672	424	888.	808	192	.36	.336	.353	273	883	- 5 08:	195	.15	80.	90:	870.		183	10.	900.	.03£	310.	(55)	140.
N ED.	BySuspd Matter.	10	3		65	<u>∞</u>	16.7	9.7	≎5 1.	9.11	10.3	10.1	9.1	35	<u>ထ</u>	··:	9.3	7	1-	1.3	7.1	2.} 	17.	23	33.	3.9
OXYGEN CONSUMED.	By Dis- solved.	1 10		9	T	-	-	:0	9	0	6:	3	+	93	??		33	C.		-	121	03	7	9	9	75 20
ÖÖ	Total.		- 00																16.				9.1			
16,	Chlorin		2 +				47.	4.4	7					4. 63									50.	∞ ??	.9	5
ON ION,	papuad -sng	01	36.	328.	100	63.	958	591.	 S:	1016.	1972.	638	344.	27.0	239	144.	548	2238	338.	174.	333	173	207	397	127.	353.
RESIDUE ON EVAPORATION	Dis-	950	242.	1.08	254.	255.	154.	168.	152	161.	1.38	11.5	131.	160.	- 288	17%	000	1.5	130.	196.	170.	.00%	191	207.	202	166
RE	Total.	836	000	436.	328.	320.	1083	759	237	-28	1100.	130	178.	130		355	698	370.	168	370	19.5		401.	601.	3331	519.
	Odor									:					:											
	Color.	36	े ००	Muddy	, ee:	-	+	ıG	ಆ	<u>∞</u>	27	<u>27</u>	<u>.</u>	સં	7.		ઌ૽	લ	rė	. 15		uj	ಣ	K	ç.	199
APPEARANCE	Sedi- ment.	Littlo	Much		Little	:	Much	:	p,suo,)	Much	*	**	:	*	;	**	*	p,suo,)	**	**	Much	9 9	Consid	Much	Little	Much
IV.	Turb'y.	Much		;	:	:	:	:	**	*	3:		**	*	:	:	:	:	•	:	*.	3.9	9.9	;	4 4	**
3 OF	1900 Exami- nation.	To no	10 11	:	Feb. 3	× :	15		Mar. 5		: 15	::		Apr. 6				May 1		:	- C		June 7	11	57	85 3
DATE OF	roon Collection.			: 67	Feb. 2	!-	11	100	Mar. 2	C. ,,	11 .,	;	× ;	Apr. 4	3	<u>x</u>		May 2		16	?;		June 6.	33	50	:
No.	Iniroz	1129	1214	X	355	1349	1300	1120	1432	1166	1501	1540	157.5	1618	1656×	1696	1735	1771	1800	1834	1860	1906	1911	1977	2013	2046

TABLE 145.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, ALTON.

O. JORDAN.	University of Chicago,
EDWIN	Univ
Jo	
Report of	

No. of Isac.	per Cubic Centi- meter.	11.000 11.0000 11.000 11.000 11.000 11.000 11.000 11.000 11.000 11.000 11.0
	Presenc to .sdA	
	TempaT TiA 10	400000000000000000000000000000000000000
	request etsW lo	0000-0000+ 0000000000000000000000000000
	Meigh:	1910 0 01 10 10 0 0 17 4 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1 0 1
RO-	Nitrates	ؿ ؿڮڗٷۺڮۺؾؖڗۺڎؖ؊ۼؿڿڮۼڔٷڹڎۺڎۼڂڹٷۺڔڿۺڿ
NITRO- GEN A	Setititi	9.000000000000000000000000000000000000
IC JEN.	pəpuəd -sng	
ORGANIC NITROGEN.	Dis-	
OZ	Total,	
202	B -sus	
< 3 3	Albuminoid Pis- Divios	**************************************
NITROGEN .	Tot'l	86.5.4.8.8.4.5.6.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8
	FreeAm' monia.	28
ED.	Bysuspd	$\frac{9-66}{2}\frac{2}\frac$
ONYGEN CONSUMED.	By Dis- solved.	လုံးကရာလုံလုံလုံးကုံကုံကြောင်းသည်တွင်လောလုံလုံလုံ တည်းသုံးကိုလောသည်ကို သို့ သို့ သို့ သို့ သို့ သို့
CoS	Total.	$\frac{\kappa_1 \times \infty \otimes \infty}{\infty} \stackrel{\sim}{\sim} \frac{\kappa_1}{\kappa_1} \stackrel{\sim}{\sim} \frac{\kappa_1 \times \kappa_2}{\kappa_1 \times \kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_1 \times \kappa_2}{\kappa_2 \times \kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_1 \times \kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_2 \times \kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_1 \times \kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_2 \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_2} \stackrel{\sim}{\sim} \frac{\kappa_2 \times \kappa_2}{\kappa_2 \times \kappa_2} $
ıe.	Chlorin	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
ON.	pəpuəd -sng	2.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8
RESIDUE ON EVAPORATION	Dis- solved.	\$\frac{2}{2}\frac{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac{2}{2}\frac
RESEVA	Total.	28.66. 28
	Odor	
	Color.	$\frac{e^{i}}{2} \cot \omega + i + i + i + i + i + i + i + i + i +$
APPEARANCE.	Sedi- ment.	Little Much Cons'd Much "" Cons'd Much "" Cons'd "" Much "" Cons'd "" Much Little Much Little
AP	Turb'y.	Much
OF	1900 Exami- nation.	Jan. 55 35 .
DATE OF	1900 Collec-E	40401- ## 00
	Co.	Jan.
.o.M	Serial	11.84 12.13 12.13 12.13 13.13 13.13 14.43 14.43 14.43 14.43 16.13

TABLE 146.

1900 1900 Collec-Examition. Turb'y.

Serfal No.

DATE OF

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

HDAN,	University of Chicago.	No. of Bac.	per	Centi- meter.	3,500	4.500	30.200	37,500	000.6	000.311	000	500.55	(BO) 061	180,000	260,000	130.000	60.000	20.000	69,000	7.200	00000	5,700	7.500	2,600	5,000	4.100	2,600	20.000	
EDWIN O. JOEDAN	sity of	ce or		Pres Abs. c					:	:	:				:	:	:	:	:	:	:	:			:			:	:
EDWIN	Unive			dmoT Lio	7	0	10.	0.	.0.		= 0	ċ			:	10.	υ.	.0	: :	??	16.	65.		: :3:	26.	.36.	£8.	28.	:
Report of		ature er, C.	19 110	qmoT W lo	0.	0	0.	0.	1.	O	<u>.</u>		÷ ÷		6.	6:	œ	10.	5.	30.	19.		10.		24.	21.	23.	.96.	:
Repo				Hell W		4.03	3.0	0.	e : : :	. i	- د د د د	; ; ; Q	- 1	14.1	5.11	10.1	1.4	10.5	6.11	10.1	-	ος ος	Ţ.	٠. ز ت	6.1	ت. دة	50.	63.	- : : : :
		RO-NAS	ธอ	Nitrat	0.	.055	.55	iG.	5 i c	3 6		o _	6	.c.	.e.		9.	કું	ıc.	ro.		<u>.</u>	27.	18	} -	16	62.0.	<u></u>	:
	TON.	NITRO- GEN A	89	Mitrit	30.	8	.00:	.016	50.0	3.5	5.0	900	800.	800.	100.	10:	920	X.0.	.016	600	300	10.	910.	600.	900.	300.	0.	610.	:
	ε, Λι	HC GEN.		sng-sng	:	:	:	:	:	:	:				:	:	•	:	:	:	:	•	:	:	:		:		:
	нов	ORGANIC NITROGEN.		Solve Solve	:	:	:	:	:	:	:	: :			:	:	:	:	:			-							:
	IRI S	OZ	1	Tota	:		~	~:	:	:	:				:	:	:	:	:	:	:	:	:	~	:	-	:	-:	:
LION	MISSOURI SHORE, ALTON		d Am	p,pud -sng	. 168		<u>88</u>	.16	200	, c	÷2	16	1.08	:90	33	8	969	<u>x</u>	. 752	<u>න</u> 		25	œ.	<u> </u>	. 170	100	<u>8</u> :	185	
NIL.	FROM]	ITROGEN AS	Albuminoid	-sid b'vios	.192	. 193	.176	184		281.	2000	119	136	903	.216	. 155	208	. 136	. 168	. 136		<u>2</u>	86 T	? <u>?</u>	.16%	E.	.136	. 158	
S PEF	NCE F	NITRO	Albu	['10T'	.36	: :::	¥24.	.352		010.1	120.	656	1.216	1.26	.536	.416	106	33	33	.30		.512	. 512 120	X X X X	.311	296	.320	<u>x</u>	
NALYSIS-PARTS PER MILLION	ONE-FOURTH DISTANCE	A		FreeAn	.914	% % %	101.	.656	.376	23 0	8. s	16.	155	376	.216	.192	881.	076	083	210.		.03		0.	330	910.	.035	0.00	:
IS-I	ктн	ZD.	bd bd	Bysus	₹"	00	£. €5	+ :		0.0	× - ×	- 00	0	6.0	8.6	00	€. 1.	23	÷	0C (5.7	- · · ·		63 (C)		1-	£0.	7.1	
LYS	-For	OXYGEN	·p	By Dis	0:		÷.	9	23 -	+ =	7 10		50						G	}			25	G	6.6		6.1	9.9	
INA	ONE	Con		Total	7.6	∞	9.e	တ်)))		φ. α α	0 1	0.1	?? ∞	χ. (C)	(c)	9.3	oc :	©: ∞	G :	6.5	10	6.7	00	6.3	₹-	9.6	1+.	:
TER.	IVER,	.91	ııı	СРГО	15.9	6.3		= -		200	φ- π-π	7 17	00	- · · · ·	3.1	55 50	1.6	9.0	03 03	ر د د	9. e	7		63 60 60	55 55	√	 		:
SANITARY WATER A	IPPI B	N.C.	pa	sng	12.	46.	158.	.69	57.	380.	20 ±	931.	544.	194	265.	220.	527.	× 2	- i	1,00	410.	226.	512.	10.	178	:08:	101.	126.	:
IT.AR	Irsiss	RESIDUE ON EVAPORATION		solved				_			151.									-						_			:
SAN	TER-	RESI	-1	T'018	302.	21:1	350.	307.		.He)	620.	SE	618.	608	381.	380.	662.		914.	315.	552.	.05	6.31	396:	332.	256.	258.	573.	
	SOURCE OF WATER-MISSISSIPPI RIVER,		Odor						:								:	•							:				
	Sourc			Color.	હ	18.	es;	7.		7.1	رة <u>-</u>	. <u>~</u>	63	?	3)	œ.	. 43	183	.e.	4	2-0	ဆ	₽.	.35	ည့	₹.	ઝ	c;	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
		PEARANCE		Sedi- ment.	Little	Much	-0	Little	:	Much	P. suc.	Much	27	**	9.9	99	;	9 9	99	Consid	9 :	33	Much	9 9	Consid	Much	Little	Much	•

1213 ... 1 Jan. 5 1213 ... 10 25 1321 Feb. 2 Feb. 3 1321 Feb. 2 Feb. 3 1452 ... 14 4. 15 1452 ... 25 1

TABLE 147.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS—PARTS PER MILLION. SOURCE OF WATER—MISSISSIPPI RIVER, 100 FEET FROM MISSOURI SHORE, ALTON.

JORDAN,	University of Chicago.
	ty o
EDWIN O.	Universi
rt of I	
Report	

No. of Bac.	per Cubic Centi- meter.	3.400 3.400 3.400 3.400 3.400 3.400 3.400 1.500	
	Presen Abs. of		
	Temper TiA lo	40000000000	
	Temper otsW lo	0000-0000+ 0000000000000000000000000000	
	ugi9H otsW	1920 03 02 02 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
RO- N AS	Nitrates	Sgriftstage - Februargeringstage 400 08	
NITRO- GEN A	Nitrites	90000000000000000000000000000000000000	
TIC GEN.	pəpuəd -sng		_
ORGANIC NITROGEN.	Dis-		_
OZ	Total.		_
76	p,pud -sng	######################################	
ITROGEN AS AMMONIA.	Pioning Pionin	0.00	
NITRO	TotT ToT	S.	
	FreeAm- monia.	24444888888888888888888888888888888888	
ED.	BySuspd	<u>- 4 - 1 - 2 た - 1 - 1 2 2 2 2 2 2 2 1 2 1 で で で 2 - 4 3 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2</u>	
OXYGEN CONSUMED.	By Dis- solved.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
Con	Total.	$r \sim 0 \cdot r \cdot r \cdot r \cdot a \sim r \cdot a \cdot r \cdot a \cdot$	
,9n	Chlorin	ಪ್ರಾಲಂ ೦೦ ಜಲು ಎನ್ನು ಎಲ ಬಲು ಬಲು ಎಲು ಬಲು ಎಲು ಬ ಪ್ರವಹ್ತ ಹೆಚ್ಚು ಪ್ರಮತ್ತು ಪ್ರಭಾಗವಾಗ ಪ್ರಭಾವತ್ತು	
NO.	pəpuəd •sns	38.5.2.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3.3	
RESIDUE ON EVAPORATION	Dis- solved.	28.6.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2	
RESEVA	Total.	28.88.98.66.66.66.66.66.66.66.66.66.66.66.66.66	
	Odor		
63	Color.		
APPEARANCE.	Sedi- ment.	Little Much Cons'd Much Cons'd Much Cons'd Much M	
T.	Turb'y.	Much	
OF	1900 Exami- nation.	Jan. Jan. Jan. Jan. Jan. Jan. Jan. Jan.	
DATE OF	1900 1900 Collec- Examition.	Jan. 10 Jan. 20 Feb. 24 Jan. 20 Jan. 2	
.oV	Serial	1186 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	

TABLE 148.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 400 FEET FROM ILLINOIS SHORE, AT CHAIN OF ROCKS, PUNPING STATION ST. LOUIS WATER WORKS.

Report of Edwin O. Jordan. University of Chicago.

			1888	38	9	23	8	8	38	38	8	8	38	3	9	000	300	
	No. of Bac.	per Cubic Centi- meter.	3,000	76.0	76.0	150.0	80.08	23.0	0.7.	0.12	9,5	10.3	51. 5 51. 6	2.0	16.5	300	15,0	
		Presence or Abs. of Colf.							:			:	:					:
	Temperature of Air, C.		1G	: ; T	- 	<u> </u>	1-	17.	: i g	3	×	÷	7.00		- 1	101	23	•
		Temperature of Water, C.		>	0.	i ai	ıc.	{- i	<u> </u>	5.	17.	٠ ٠ ٠	F) F	189	5	159		
		Heigh Hate	63 00 10	0 1-	သ	<u>∞</u> <u>n</u>	12.	14.5	e ie xi x	17.5	16.3	12:	100	2 2 2	10.7	100	13.9	:
	RO- N AS	Sitrates	86.18.18	=	1.15	19 =	Z	2	_ = %	0	લ્લ	10.1	7.	100	335	0	199	
	NITRO- GEN A	Nitrites	510.	800	810.	9.9 8.8	.013	.017	8 8	0.0	.013	.019	- SA. N	000	300	000	.003	
	IC EN.	papuad gas-			•			:			:	:	:					
-	ORGANIC NITROGEN.	Dis-									:	:	:					
	OX	Total.			:			:	:		:	:	:				:	:
		B p.pud	80.41	836	9.36	000 000 000 000 000 000 000 000 000 00	505	316	20.00	33		.576	: : -	.92	30	.346	.530	
	NITROGEN AS AMMONIA.	Pioniminos propins de la propi	55 ci 25	7	- FEE		<u>86</u>	.136	ž ::	.176		144	132.1	176	0.83	960.	.136	
	NITRO	T'10T	+ + + *	86.	1.16	3; B	£:	355		.196		27.5	101.	539	:06:	= -	.656	
		FreeAm. monia.	969	576	395	35 E	.316	28.7	S 95 9 5 9 5 9 5	.040.		98	100 X	3	330	+10	810.	:
Į	.D.	Bysuspd		01	7.6	201	©};	3.6	0 01	or. ∞	1-	0.3	 	. w	5: 1	6.4	1.5	
1	FGE	paalos	7-1-1-	G			_	G) :	0 0	21	10		7. 00	ı	-	9	10	
١	OXYGEN CONSUMED.	Total.	- − − − × ×	22	{- :	ः ल	??	x. :	c m		œ.	e: -	- 12	-			-	
	.et	Chlorine.				.6			-									
-).X (0.X.	popuod -sus	13.1.0	1305.	1269.	-3. -3. -3. -3. -3. -3. -3. -3. -3. -3.	470.	305.	157.	. S. S.	331	. 116	0.00	712.	2383	706.	1059.	
l	RESIDUE ON EVAPORATION	Dis-	9 9 9 9	216.	9	10.00	Ξ.	155	176.	178.	173		3 2	220.	210.	203	276.	
-	RE: Eva	Total.	201. 504.	1518.	1119.	£ 500	619	450.		486.	5(€):	165 185 185 185 185 185 185 185 185 185 18	126	303	3503	.908	1235.	
	Odor				•						:						•	
		Color.	Si S	•	₹.	28	<u>x</u>	es		55	9	es e	ું જ	999	33	e.	£6.	
	APPEARANCE	Sedi- ment.	Cons.d Nuch	:	:		:	* 1	Cons.d	Much	7 9	66	Much	93	9 9	9 9	*	
	1.	Turb'y.	Much	:	90		9.9	: :	10	,	9	0 1	:	9.9	9 9	:	-7	0 0 0 0 0 0 0 0 0 0 0 0
-	OF	fxami- nation.	:: :a:::											œ	14		. 38	
	DATE OF	- 6 E E	5 Jan.	X	200	三 智	5.	500	12 50	16	7. 0	(~ ~	1 2 2 2	1-	+	-	x	:
	DA	1900 Collec- tion.	lan.											June		:	31	
	No.	Iniros	1981											194×	1985	90.30	2000	:

TABLE 149.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

University of Chicago.

Report of EDWIN O. JORDAN,

No. of	per Cubic Centi- meter.	2. 1. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.
	Presence Abs. of C	
	Tempera of Air,	4
	Tempera of Water	000
	Helght 1918W	ej e. re ej oo re oo re
RO-	Vitrates	8000 4 1 - 0 4 1 0 0 1 0 0 4 4 1 1 1 1 1 1 1 1 1 1
NITRO-	Nitrites	999998888888888888888888888888888888888
NIC GEN.	-sng	
ORGANIC NITROGEN.	Dis-	1 : : : : : : : : : : : : : : : : : : :
	Total,	
on on	A -sng	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
ITROGEN AS	In I	138 138
NITRO	T'10T	85.5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	FreeAm.	25.00.00.00.00.00.00.00.00.00.00.00.00.00
SN IED.	BySuspd Matter.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
ONYGEN CONSUMED.	By Dis- solved.	<u></u>
<u>ి</u>	Total.	& r r r r r r r r r r r r r r r r r r
.91	Срјоги	919144 1919 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
ON TON.	pəpuəd -sng	88 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
RESIDUE ON EVAPORATION.	Dis-	9.50 1.50
EVA	Total.	846.6 1855.0
	Odor	
	Color.	3 0 8 8 4 9 8 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
APPEARANCE.	Sedi- ment.	Little Cons'd Much
AP	Turb'y.	Much
E	1900 Exami- nation.	n r r r r r r r r r r r r r r r r r r r
DATE OF	c- Ex	11 :: 28 Feb. 28 Feb. 29 Feb.
DA	1900 Collec- tion.	Tan. Mar. May. May. May.
No.	Serial	1198 1298 1298 1298 1298 1298 1298 1298

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT. SANITARY WATER ANALYSIS—PARTS PER MILLION.

SOURCE OF WATER — MISSISSIPPI RIVER, AT CHAIN OF BOCKS, INLET TOWER, ST. LOUIS WATER WORKS.

EDWIN O. JORDAN,	University of Chicago.
O ſ	
Report	

No. of Bac. per Cubic Centi-meter.	2. 2. 2. 6. 600 2. 2. 2. 6. 600 3. 2. 2. 6. 600 3. 2. 2. 2. 600 3. 2. 2. 2. 2. 600 3. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2.
Presence or holi.	
emperature of Air, C.	<u> </u>
Comperature Of Water, C.	00110146665565656565
Height of Water.	
NITTRES GEN OF THE SOLUTION OF	
	20000000000000000000000000000000000000
Sus Sandad	
Potal. Nitraogen. Nitraogen. Onganical Sus-	
Yu -sns	
AMMONIA. Albuminoid	1
NITRO INDI	888 588 588 588 888 888 888 888 888 888
reedm- monia.	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0
YSuspd B	
Total. Young Total. Young Total. Young Total. Young Total.	
Potal. So	ででに下でる では では で で で で で で で で で で で で で で で で で い で で い で い こ
Chlorine.	<u> </u>
Sus- Sus- Sus-	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Total. EvaPonation Colved. Col	1
Total. E	23.50 23.50 23.50 25.50
Odor	
Color.	Mudddy
APPEARANCE Sedi- ment.	Cons'd Much Much Cons'd Much Cons'd C
APP Turb'y.	Much
tor 1900 Exami- nation.	0512 0 2 0 2 0 2 0 2 2 2 2 2 2 2 2 2 2 2 2
DATE OF O 190 llec- Exa on. nati	28 28 28 28 28 28 28 28 28 28 28 28 28 2
DATE OF 1900 1900 Collec- Examition.	Mar. Mar. Mar. Mar. Mar. Mar. Mar. Mar.
Serial No.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

TABLE 151.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIFFI RIVER, 400 FEET FROM MISSOURI SHORE, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

JORDAN,	University of Chicago.
EDWIN O.	University
Jo	
Report of	

No. of Bac.	per Cubic Centi- meter.	25.000 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300 11.300
	Presenc	
	RTSqm9T TiA To	
	rempera	001-0-4767-5767-388888888
	Height 91aV/	8 9 1 9 1 9 1 9 1 9 1 9 1 9 1 9 1 9 1 9
RO-	Nitrates	6. 14.14.00 10 10 10 10 10 10 10 10 10 10 10 10 1
NITRO- GEN A	Nitrites	20000000000000000000000000000000000000
HC BEN.	-sng	
ORGANIC NITROGEN.	Dis- solved.	
ōΖ	Total.	
	Bud, pud	44 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 +
EN AS	Pioniminos -siq	0.096 0.096
NITROGEN AS AMMONIA.	Tot'1	824317444 517 88 82 82 82 82 82 82 82 82 82 82 82 82
A	FreeAm monia.	1.01.02.03.03.03.03.04.04.05.03.03.03.03.03.03.03.03.03.03.03.03.03.
N ED.	Bysuspd Matter.	4 61616 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
OXYGEN CONSUMED.	By Dis-	000000000000000000000000000000000000
Con	Total.	441-61-60-61-68-60-61-68-40-61-60-61-61-61-61-61-61-61-61-61-61-61-61-61-
,9t	Chlorin	で 0 2 4 4 8 0 0 2 2 2 8 8 8 8 8 8 8 8 8 8 8 8 8 8
NO.	pəpuəd -sns	36. 166. 166. 178. 178. 178. 178. 178. 178. 178. 178
RESIDUE ON EVAPORATION	Dis- solved.	88.88.88.89.89.89.89.89.89.89.89.89.89.8
RES EVAI	Total.	1236. 1027. 1102. 1102. 1102. 1102. 1103.
	Odor	
	<u> </u>	
	Color.	
APPEARANCE.	Sedi- ment.	Little Cons'd Much
Ār	Turb'y.	Much
OF	1900 Exami- nation.	Jan. 6 " 126 " 27 " 27 " 24 " 24 " 27 " 26 " 27 " 27 " 27 " 27 " 27 " 27 " 27 " 27
DATE OF	1900 Collec- E tion. na	10 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
	Col Col	Jan. Feb. Mar. Apr. " " " " Jung
.oV	Serial	1195 13300 13300 13300 1427 1427 1427 1530 1530 1742 1659 1659 1743 1741 1816 1816 1816 1816 1816 1816 1816 18

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

NOURCE OF WATER-MISSOURI RIVER, FORT BELLEFONTAINE, WEST ALTON, MO. SANITARY WATER ANALYSIS-PARTS PER MILLION.

University of Chicago.

80
7.
P 4
V
20
-
0
-
0
\cup
0.00
7.
penti
-
, min.
2000
F+2
Jordana
Carrel .
Ç
-
5
-
~
-
Q.
Page 1
-

No. of 13ac.	per Cubic Centi: meter.	13.26.0 17.0	25.000
	Presenc lo .sd A		• • •
	Temper	wie a we Fire # Fire Fire Fire Fire Fire Fire Fire Fire	36.
	rempers		8
	Heigh Uate	- w + r o s s s s s r c c c c c c c c c c c c c c	12.9
NITRO- GEN AS	Nitrates	٥ ٢٤ تا ١٥ تا تو الآ الأ يو عن الا ما الإ الله الله الإ عادة الديو لـ م ها الا إ	§ :
NITR	Nitrites	865 - 95 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	8: :
GEN.	pepued -sng		
ORGANIC NITROGEN.	DIS- solved,		
	pnd'd "		
S S	Sus-sug	2	- :
NITROGEN AS	Pionimoid Pivios Divios	28	3) :
NITH	T'10T	25	26.1
	Freedm. monia.	0.054 0.054	
EN ded.	Bysuspd		10.1
ONYGEN CONSUMED.	By Dis-	- 10 to - 0 + 0 to	G :
	Total.	<u></u>	
,9r	Chlorin	#####################################	x :
ON ION.	-sng	68 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	.5018.
RESIDUE ON EVAPORATION	Dis-	######################################	× 550
REEVA	Total.	44.88 62.09.88.19.19.19.19.19.19.19.19.19.19.19.19.19.	\$7.16.
	Odor		
1	1	- 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/ 6/	: :
6-3	Color.		
APPEARANCE.	Sedi- ment.	Little Nuch Cons'd Little Little Cons'd Much Little Cons'd In the cons'd	
Ar.	Turb'y.	Nuch	
Es,	1900 Examination.	60 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	8:
DATE OF	c. Ex	Jan. 1987 × 1987 × 1987 × 1987 × 1987 × 1987 × 1988 × 1988 × 1987 × 198	
DA	1900 Collection.	Heb. San.	
.o.V	Serial	11.59 J. 11.59 J. 11.50 J. 11.	3000

TABLE 153.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 VARDS FROM ILLINOIS SHORE, JEFFERSON BARRACKS. MO.

JORDAN,	University of Chicago
EDWIN O.	University
Report of	

No. of Bac.	per	Centi- meter.	008	4,300	6,600	40,000	8,400	102,000	39,000	90.000	135,000	190,000	210,000	120,000	29,000	46,000	7.000	32.000
ce or Coli.		Pres.	:	:	•	:	•					:		:	•	:	:	:
		Temp To	:		:		:	:		:	:			•		:	:	:
		TempT W lo	0.	0	က်	4.	0	· 0	0	0.	.0	21	େ:	ос°	<u></u> -	œ	5	16.
tot i.re	d ₂	ieH W	1:	35 70	9.€	0.0	.9	4.7	6.4	6.3	14.	25.5	18.5	13.	14.8	15.8	16.	17.4
NITRO- GEN AS	sə	Vitrat	ಬ್															
NE	sə	ninin	.014	986	.008	.012	.012	.00%	<u>.</u>	.016	.016	.006	800.	.014	.01	8	8	.03
C EN.		puəd sns	:	:	:		:			:	:	:	:	:		:	:	•
ORGANIC NITROGEN.	.b	SiU Solve	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
OR		Tota	, :	:	:	:	:	:	:		:	:	:	:	:	:	:	:
	Am.	p,pud -sus	.112	.136	368		.064	1.032	.693	.112	968.	186	.613	.164	888	.456	. 248 842.	191.
ITHOGEN AS AMMONIA.	Albuminoid	p'vios	.192	1364	.176	.176	184	. 168	.208	802.	7 90.	.136	.138		.136	. 168	144	.16
NITROGAM	Albu	Total	304	005.	544	.496	.248		6;	£.	96.	1.19	₩.	:504	484	.62 1	.392	.624
		FreeAr	.056	.355	8	. 296	9.	.64	++.	.384	. 19	.254	98.	288	. 184	. 164	.048	.058
ED.		Bysus	1.1	0.	ં	4.6	T: -	13.7	5. 00	2.6	14.	13.	11.8	<u>رن</u>	4.1	9.6	3.6	9.4
ONTGEN CONSUMED.	.ba	By Dis	6.1	C.5	9	寸	5	∞	<u>G</u>	9	9	∞		5	ci.	4	લ્યું	es .
Cos	_	Total	3- 0;												9.3			
re.	ıļı	СРЈО	18.6	_		_					_	_				_	_	
ON.	pe	epuəd sns	35.	36.	56.	270.	39.	1330.	697.	120.	1168.	1129.	801	534.	348.	138.	349.	523.
RESIDUE ON EVAPORATION.		osid solve	330.	 0.51	236.	192	285	306.	176.	194.	160.	163.	114.	158.	146.	166.	159.	164.
REA	-1	Total	362.	286.	- - - - - - - - - - - - - - - - - - -	164	321.	1536.	873.	314.	1328.	1292.	915.	695.	191	604	508	687.
	Odor																	
		Color.	. 18	77.	ec.	Muddy	e;	: :8:	ē.	.45	रु	٠: د:	??	ಚ	ಣ	달.	77.	35.
APPEARANCE		Sedi- ment.	Little	Cons'd	Little	Much	Little	Much	:	Cons'd	Much	:	:	3 3	;	7.7	9.9	,,
AP		Turb'y.	Much		",	:	,,	",	9,	,,,	*	*	**	9.9	40	**	9.7	77
OF	1000	Exami-	Jan. 5				Feb. 5										07,	27
DATE OF	1000	Jacon Jollec- F	7	** 11	<u>∞</u>	500		6.	000	00	œ	16	35	66		= ?	06 ,,	96
7.0.	<u> </u>	Seris															1705	1744

TABLE 154.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER. EAST OF MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of EDWIN O. JORDAN,

University of Chicago.

No. of Bac.	Cubic	Centi- meter.	2,900	1.000	18.000	16, 100	1.60G	56,000	£0,000	70,000	112.000	180.000	170,000	115,000	23.000	000°66	9,000	29.000
re or Coll.		ser4 sed A.	:	:		:	:	:		:	:	:	:	:	:	- :	:	:
		dmoT an		:	:			:	:								:	:
		TempT V/ To	0.	0	···	-	C	C	<u>.</u>	c.	٠			x.	} ~	x	27	. 9
10 1		Ileli W		55	9:10	g. c	÷.		 G.	ç. ç		6.53	(C) (C)	~	∞. <u>+</u>	<u>s</u>	16.	
NITRO- GEN AS	6 3	Mitrat	65	35	ા ે	£,	Z)	ક્	GI. I		£.		x;	x.		σ:	91	c.
NIT		Nitrit	210.	900.	30.	800	50.	3	.012	910.	910.	5.5	.01.	.01	.015	.032		99
GEN.		sns		:	:	:	:	:	:	:	:		:	:	:	:		:
ORGANIC NITROGEN.		Dis		:	:	:	:	:		:	:	:	:	:	:	:		:
OZ	1	Total		:	:	:	:	:	:	:	:	:	:	:	:	:	:	:
200	d Am	p,pud	ઃ	.156	8	.335	. 196	ž.	199	886	1.016	1.168	955.	.036	18	=	300	<u>×</u>
NITROGEN AS AMMONIA.	Albuminoid	-sid b'vios	.16	:224	9	 		.192	.176	£	190	. 152	<u>z</u>	∞ 	-	. 152	35.	. 168
NITRO	Alb	T'10'T	.36	88.	+	91.	50.	1.072	支	156	80.		7	.416	333%	.593	± 1.	9.
		FreeAr	.618	:33	.018	138	.595	52	116	3233	280	6	16. 16.	<u></u>	.176	e j	.018	9
SN (ED.		Bysusi	1.5		<u> </u>	4.6	- :	22	=	£3.	Ξ	2		-	~	- ه	3.5	
OXYGEN CONSUMED.	p	By Dis	5.4		<u>-</u> -		0		: Q		 	9:5	 	2	0.0	5.3	9:3	5.5
000		Total'	6.9		∞							× 1			<u>с</u> :	5.6	C	5.5
.91	rļı	Сијо	19.1	6.	G:		0.0								∞. •••		1.6	
ON ION.	pa	sng sng	œ	104	.96:	219.	9	32.5	656	100	<u>8</u>	1.0×	230	+33	286.	107	470.	619.
RESIDUE ON EVAPORATION		el(I	336.	236.	230.	500	296	20%	188	<u>.</u>	119.	136.	130.	143.	150	110.	170.	
REEVA		Total	344.	330.	326.	171	336.	1380.	====================================	289	1930.	16:34	%: 000 100 100 100 100 100 100 100 100 10	574.	436.	517.	640.	779.
	Odor					:	:			:		:				:		:
		Color.	5	£.	\$6 \$6	eć	ej.	ક્ષ	ņ	£.	<u>e</u> .	ti.	o į	er.	re.	300		£.
APPEARANCE		Sedi- ment.	Little	Cons'd	Little	Much	Little	Much	.,	Cons.d	Much	9.7	• •	:	:	:	:	:
V		Turb'y.	Much		**	;	:	:	:	:	:	:		:			:	:
OF	000	1900 Exami- nation.	1	:	:	4 -0	Feb.	9 9	40 3	Mar.	:	10 0	18 19	10	Anr.		21	7 9
DATE OF		Collec- tion.	1 .	:	2	:S	Feb. 3	0.	:	Mar. 3	œ	. 16	?}	हों हैं।	/ 11 J		: 20	96
No.	11	Berlin	1177	()(27)	1255		1330 F					1516					200	7.15

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, JEFFERSON BARRACKS, MO.

University of Chicago,

Report of EDWIN O. JORDAN,

17																		
No. of Bac.	per	Centi- meter.	4,200	3.500	32,900	40.000	16,200	39,000	75.000	(17,000)	100,000	260.000	230.000	130,000	19,000	104,000	11,000	31.000
		Prese																
		dm9T A lo										•						
r, C.	911 911	qmoT sW lo	0.	0	600	4	0.	0	0	0	0	c)	ં લે	∞	i-	00	2	16.
		BiəH B'//	-	3.5	9.6	5.9	9	4.7	6.4	6.9	14.	23.55	18.5	13.	14.8	15.8	16.	17.4
'RO-	89	Nitrat	ŀ															in.
NITRO- GEN A	sə	Nitrit	.01	900	000	80	990	800:	.012	.012	.014	.016	210.	012	.017	03	0.18	.018
	1 -	-sng							:	:	:			:	:			
ORGANIC NITROGEN.	q.	Dis		:				:	:		:	:	:	:	:			
OR		Total		:				:	:		:	:	:	:	:			:
	Am.	p,pud -sns	.092	. 144	321	332	890.	.648	.648	.248	976	1.212	.652	.116	184	.552	344	.504
NITROGEN AS AMMONIA.	Albuminoid	Dis-	.138	.176	. 136	128	.152	.136	. 184	808	.064	. 168	. 168	.34	.152	.176	144	.168
NITROGEN AMMONIA	Albu	T'30T	.26	33	.46	.46	8	184	.832	.456	1.04	1.38	£	.456	.336	7.58	.488	.672
		FreeAi	.052	.376	880	108	.544	304	.352	.286	.16	333	.252	.236	808	966	.036	.05
ED.	r. pd	BySus	e;	-:	2.6	3.5	<u>ઃ</u>	11.4	9.5	25.50	15.						@ 1	12.1
ONYGEN CONSUMED.		By Dis					₩.										6.7	6.1
် ဗိ	_	Tota					6.8					19.8		12.6				18.2
.ər	ıįı	СРЈО	21.4	10.	<u>:</u>	- 2 . 2	17.	11.5	5.6	6.6	ت. زه	υς ∞	3:3	6.5	2.5	33	5.4	5.5
ON.	pe	puəd Sus	£	110.	162.	318.	122.	856.	543.	175.	1910.	2252.	821	498.	333.	555	710.	824.
RESIDUE ON EVAPORATION.		sojve Dia	339.	252.	256.	251.	280.	246.	186.	300	116.	142.	113.	126.	156.	134	157	156.
REVA	1.	Tota	403.	362.	418.	569.	405.	1102.	739	375.	2026.	2394.	934.	624.	489.	689	36	980.
	Odor			:					•				•			•		
		Color.	.13	7.	ಸು	<u>دن</u>	જ	33.	ಣ	.35	. 15	<u>دن</u>	ા	<u>ښ</u>	ಣ	.38	4.	용.
APPEARANCE		Sedi- ment.	Little	Cons'd	•	Much	Little	Much	9.	Cons'd	Much	;	99	99	99	**	7,9	99
AP		Turb'y.	Much		:	:	***	1	;	\$:	*	9,	7.9	7,	9.	99	7 9
OF	1000	Exami- nation.	Jan. 5														30	22
DATE OF			4 38		18	25	3	6	3	3 M	000	16	55	66	5 4	5	02	36
DA	1000	Collec- tion.		:	:	9.9	Feb.	7,9	9.	Mar.	;	79	"	9 9	Apr.	9.	7.9	7
No.	[B	Seri	1178	1231	1256	1292	1331	1367	1416	1451	1479	1517	1552	1590	1629	1663	1707	1746

TABLE 156

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION,

Socree of Water-Mississippi River. West of Midstream, Jefferson Barracks. Mo.

Report of EDWIN O. JORDAN.

University of Chicago.

No. of Bac. Per Cubic	Centi- meter.	1 000	200	10 200	74XXX	000.0	8,200 40,000	43,000	45.000	24,000	140,000 900,000	990,000	200,000	000,061	000,000	000,000	00°88
sence or .ilo.) to										:	:	:	:	:	:	:	
perature Air, C.				•					:		:	:		:	:		
perature Jaier, U.			i d		· +:	: <	; c	; c	; c	; -	်င	i÷	, ,	ć i	- 0	· ÷	16.
ight of Tater.		-	24	0	210		. 1.	2 2	7 0	? -	9 2	100	101	1100	1 14 1 14 1 14	16.0	
Seri seri	Nitra	1									7	-					.33
	Nitr	010	000	(3)	008	00.	8	250	300	3.5	110	250	210	200	0.00	210	0.0.
NIC Sec. Sec. Sec. Sec. Sec. Sec. Sec. Sec.	ng						0	•	:	:	•	•	•	:		:	: :
1 4 % 1 6 %	ros Nos						•			:	:	:	:				
ox la	toT							•		•	•	•	:	:	•	:	
V	sns	96	.164	248	18	136	300	803	1.11	1 506	1.410	() L	3	2 2	7.17	100	35
	sid vlos	8	960	300	0.1	550	50	168	1.38	()()	1.00	168	36	2	378	500	28.
NITRO AMA	'30T	85	.26	.376	296	386	97	576	9,70	38.	10	C	17		- 1	250	.656
	Free.																.048
rspd By Teach	Bysu	4.	-	3.6	4	2.1	-	5.6	61	5	33	21	1	- }-	1	0. 10	8 11.9
ONYGEN CONSUMED.	Bk D	55	10	3.										0.7			
al. Se	toT	5.6	6.7	X:	œ	6.7	30	27	1	38	19.5	18.6	51	<u>01</u>	27	9	16.7
lorine.	СР	1.	16.	06	15.9	19.	53	10.6	19.5	10	-	6	4.6	oc ⊙	8.23	-	×.
ded box	ns ned	57.	- 128		383	=======================================	600	5.	160	3576.	2310.	993	118	681	965	1322.	1360.
RESIDUE ON EVAPORATION	Nos D	318	280	308	267.	288	236.	2008	318	1.80.	152	15.	134	218.	190	333	201
Eval Jan	10'1'	375.	38.	37.5	555	100	35	676.	178	3706.	2162.	1008	552.	800	1161.	1541.	1574.
Odor																	:
10101		.15	125	ıc	શ	છું	8	33	18	15	ବହ	cj	00	27	řě.	wh.	8
APPEARANCE.	ment.	Little	Cons.d		Much	Little	Much	9 4	Cons'd	Much	4.9	**	. ,		:	:	
Turb's		Much		•	a up	:	9 0	:	9.0	, ,	7 9	4 4	9.0	9 9	9 9		;
1900 1900 Exami:	nation.	Jan. 5	<u>?</u>													02	33
DATE OF		- miles	=	$\frac{\infty}{}$	12	60	0	Ĝ.	33	ЭÇ.	16	37	230	٠.0	2	06	38
oN fatr		1179 Jan.														- X3-	

TABLE 157

Serial No.

STREAMS EXAMINATION—CHICAGO SANITARY DISTRICT.

SANITARY WATER ANALYSIS-PARTS PER MILLION

Source of Water — Mississippi River, 100 Vards from Missouri Shore, Jefferson Barracks, Mo.

University of Chicago.

Report of EDWIN O. JORDAN,

286.000 175.000 175.000 175.000 175.000 175.000 175.000 27,500 27,500 28,100 38,100 38,100 38,000 No. of Bac. per Cubic Centimeter. Presence or Abs. of Colf. Temperature Of Air, C. Temperature of Nater, C. Height of Water. المنافق عند المنابعة GEN AS NITRO-SHIRITA Nitrites pəpuəd -sng ORGANIC NITROGEN. solved. Dis-Total. p,pud -sns Albuminoid Am NITROGEN AS p'sid b'vlos AMMONIA. 848844469944468888 [,10], monia. OXYGEN CONSUMED. By Dis- $\begin{array}{c} rv \otimes r \cdot \otimes rv \otimes \overline{-r} \cdot \overset{\mathcal{G}}{\otimes} \overrightarrow{\circ} \overset{\circ}{\circ} \overrightarrow{\circ} \overset{+}{\rightarrow} \overline{-r} \overset{\circ}{\circ} \overset{\circ}{\circ} \overset{\circ}{\circ} \\ r\dot{\circ} \overset{\circ}{\circ} r \cdot \overrightarrow{r} \cdot \overset{\circ}{\otimes} \overset{\circ}{\circ} \dot{\circ} \dot{\circ} & \overset{\circ}{\circ} \overset{\circ}{$ Total. Chlorine. beugeg RESIDUE ON EVAPORATION. Dis. 23.50 128. 1786. 1786. 1756. Total. Odor Color H er er er er er er gj er H en er er er gj + gj APPEARANCE Sedi-ment. Little Cons'd Much Cons'd Much Cons'd Much Turb'y. Much 1900 1900 Collec-Examirtion. 3 Feb. 9 · · Apr. : : : .. Mar. DATE OF 31 m m m 231 82 m m m m 23 1180 Jan. 1233 ... 1258 ... 1294 ... 1333 Feb. 1369 ... 1418 ... 1453 Mar. 1519 ... 1582 ... 1582 ... 1665 ... 1709 ...

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-ST. LOUIS, MISSOURI, TAP WATER.

Report of Edwin O. Jordan, University of Chicago.

No. of 13ac.	per Cubic Centi- meter,	600 600 600 600 600 600 600 600 600 600	2,400
	Presence Abs. of		
	Temperation To		
	Temperator Nate		
10 1	Heigh ots7//		
RO. N AS	Nitrates	-(a + b + b + b + b + b + b + b + b + b +	25
NITRO GEN	Sitrites	990000999999999999999999999999999999999	00 o :
NIC GEN.	-sug		
ORGANIC NITROGEN	Dis-		
	b'baq	x = x = x = x = x = x = x = x = x = x =	÷ : : :
02	Sus-sus	988	
TROGEN A	Pionimoio pionimoio pionimoio pionimoio pionimoi		21.
NITROGEN	E I'10T	### ### ### ### ### ### ### ### #### ####	91
	FreeAm- monia.	60000000000000000000000000000000000000	
EN IED.	Bysuspd Matter.	10000 4 4 1 1 0 0 0 1 1 1 1 1 1 1 1 1 1	- :
OXYGEN	By Dis-	$\frac{\omega_1+\omega_2}{\omega_1+\omega_2} = \frac{\omega_1+\omega_2}{\omega_1+\omega_2} = \frac{\omega_1+\omega_1+\omega_2}{\omega_1+\omega_1+\omega_2} = \frac{\omega_1+\omega_1+\omega_1+\omega_2}{\omega_1+\omega_1+\omega_2} = \frac{\omega_1+\omega_1+\omega_1+\omega_2}{\omega_1+\omega_1+\omega_2} = \frac{\omega_1+\omega_1+\omega_1+\omega_1+\omega_2}{\omega_1+\omega_1+\omega_1+\omega_2} = \omega_1+\omega_1+\omega_1+\omega_1+\omega_1+\omega_1+\omega_1+\omega_1+\omega_1+\omega_1+$	_ ic :
၂၀၅	Total.	44444444444444444444444444444444444444	χ - : : : : : : : : : : : : : : : : : : :
*9t	Срјоги	$\frac{\text{deg}(2)}{\text{deg}(2)} \frac{\text{deg}(2)}{\text{deg}(2)} \text{$	
O.N.	papuad -sug	である。 である。 では、 では、 では、 では、 では、 では、 では、 では、	97. 158.
RESIDUE ON EVAPORATION	Dis-	%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%	916.
RESE	Total.	######################################	374.
	Odor	0:::::::::::::::::::::::::::::::::::::	
	1 .		-
1	Color	NI NI	יים פיים יים
APPEARANCE.	Sedi- ment.	V. Little Cons'd Little	V. Little Little
A	Turb'y.	Much d	
OF	1900 Exami- nation.		88
DATE OF	c)		63 A :
	1900 Collection.	Jan. Feb. Reb. Apr. Apr. Apr.	9 9 9
No.	Serlal	# 25	9099

TABLE 159.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, BRIDGEPORT.

Report of Adolph Genemann, D. B. Bisber, Municipal Laboratory, Chicago, III.

No. of Bae.	per Cubic Centi- meter	metet:	105,000	81,350	362,250	200,000	515,000	642,000	2,650,000	186,000	160,000	392.000	845,000			2,160,000			8,425,000	
	resenda jo sq		+	+	+	+	+		+	+	+	+	+	+	+	+	+	+		
	eniper viv lo	JL			27.	ਲ ਨ	24.	21.	19.	15.	19.	10.	10.	20.	10.	=	****	10.	0	
ature D., T.	emper ots <i>II</i>	o L	:	6.	25.5	10.	19.	19.	13.	6.	13.	13.	i-	10.	0.	œ	6.5	5.5	9.	
	Heigh Hate			1-	5.4	7.	5.5		5.	řÇ.	ت ن	4.7	5.6		5.5	5.6	ī.c	5.	5.	
GEN AS	sətsiti	Z.	0.	97.	.30					:		:	:	:	:	-			08.	
	settriti			7 9.				003		٠				9	- -	Ė	003		c	:
GEN.	Sus-				51.73	_	01	0	:	3	3	1 2.46	_	35		_	35	90	31	
ORGANIC NITROGEN.	Dis-	8	31.73	≎5		_	-	_		0.1.6	3.5.5	1.84	ल १	1.7	3.2.4	0.20	1.70	2.3	20.0	:
OZ	leio'.	L			3.08														4.7	:
7	ed Amsterday		1.46	€.	89.	99.	1.0	68.	98.	1.01	.86	1.26	.92	08.		17.	1.15	1.83	1.90	:
NITROGEN AS	Albuminoid -sic	os I	6.	07.	1.20	07.	<u>08</u> .	1.16	1.04	1.23	1.39	1.26	1.30	1.13	1.46	1.26	1.64	1.50	1.30	
NITHO KWA.	[1.10]	\mathbf{T}	2.40	1.55	1.88	1.30	1.80	60. G	1.84	3. 2 <u>4</u>		9.50 50.50		1.93	2.68	3.00	2.79	3.39	3.20	:
	reeAm- nonia.	H	12.60																	
ED.	ysuspd latter	B	18.	14.6	₹ ₹	<u> </u>	30.	18.5	<u>.</u>			19.6		18.	21.6	14.4	18.4	30.	30.4	:
OXYGEN	y Dis-	В	23.		4	G)		1-		9	c)		4	₹.	∞	9.	77	77	9.	
Cos	lotal.	L		9	6.14	33		G?		00	7	9	C.5	4	4		∞	TJ.	0.	:
.əu	hlori)	.63						5					10	10		5			
ON.	ended Sus-	d	.99	70.	.69	11.	76.	51.	65.	42.	63	:58:	<u>.</u>	.89	.89	54.	.69	151.	157.	
RESIDUE ON EVAPORATION	Dis-	os	416.	534.	555	389	414.	489.	468.	482.	572.	441.	459.	330.	456.	373.	498.	556.	438.	
REVA	'lato'	L	482.	604	591.	436.	765	540.	533.	524.	635	499.	532.	458.	524.	427	567.	707	595.	
	Odor		Gassy	3lGas'y	St'gGas	Gassy	*	:	9.9	St'gGas	3	23	Gassy	St'gGas	Gassy	;	7,9	**	33	
	Color.		Muddy			_	-	**			:	:	:	* * * * * * * * * * * * * * * * * * * *	:		:		:	
APPEARANCE.	Sedi-				1I. N			11.	S.	VS.	VS.		. S.	VS.	S.	Š	VVS.	VH.	II.	•
AP	Turb'y.		VVIII.	VVII	VVIII.	VVII.	VVH.	VV.11.	77.11.	VV111.	VVIII.	VVII.	VVII.	VVII.	VVIII.	VVII.	VVIII.	7.7.11.	.11.7.7	•
OF	1899 Exami-		July 3		F6 .,			11 ., 11			192					Nov. 6		06 ,,	37	:
DATE OF	1899 Collec- F	- ,	July 31	17	31	31	Vug.	:	sept.	9 0	: ::	Det. 2			30		13	06	27	
.o.X	Serial		31	333	공	55	36	33.	40 %	1	43	43 (:	46	1-	87	65	000	1451	•

APPEARANCE-II., Heavy. VII., Very Heavy. VVII., Very Heavy. S., Slight VS., Very Slight. VVS., Very, Very Slight. M., Medium.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS AND MICHIGAN CANAL, LOCKFORT, ILL.

Report of Adolph Gehrmann, D. B. Bisbee. Municipal Laboratory, Chicago, III.

No. of Bac.	Cubic Centi- meter.	102.250 630.600 12.250 600 12.250 600 17.500 1.260.600 1.260.600 1.670.600 1.070.600 1.070.600 1.070.600 1.070.600 1.070.600 1.070.600 1.070.600
	Presen To .sd A	+++++ ++++++
	ioqmoT ii k. To	สียยุยยยยยยยยยยยยยยยยยยยยยยยยยยยยยยยยยย
	requer of Wat	REPRESENTED REPRESENTED
	1161gp 1877	
NITRO- GEN AS	estantiv	
1	Nitrites	0090 008080888 5 88800
OEN.	Sus	8
ORGANIC NITROGEN.	-sl(I beylos	88888885 88888885 888888885 88888888
ON	Total.	**************************************
11 1.	p,pud -sng	85%34836833383288366888
NITROGEN AS AMMONIA.	P. VIOS -SIG -SIG -SIG -SIG -SIG -SIG -SIG -SI	85888888888888888888888888888888888888
NITRO	[.10.I.	85854848884858485888886858
	FreeAm- monta.	8886437777888888888888888888888888888888
ED.	BySuspd	$\frac{48\pi r + 6\pi r + 68\pi r r r r s + 7r r r s + 88874884}{160144666666666666666666666666666666666$
ONTGEN CONSUMED.	By Dis-	55722777722222222222222222222222222222
03	Total.	<u> </u>
ou.	irolil')	PP4888898989888888888888888888888888888
0.N 10.N.	papuad -sus	88888888888888888888888888888888888888
RESIDUE ON EVAPORATION	paxios Dis-	======================================
RES	Total	10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
	Odor	Stransky Gassy Gas
	Color.	Mudddy San
APPEABANCE.	Sedi- ment.	
A.P.	Turb's.	
s oF	1899 1899 'ollec- Exami- tion. nation.	# 2 L 2 2 L L 2 2 L L X 2 2 2 2 2 2 2 2 2
DATE OF	1899 Tollection.	# = = = # = = = # = = = = = = = = = = =
.o.Z	Iniros	======================================

APPEARANCE—H. Heavy VII., Very Heavy, VVII., Very, Very Heavy, S., Slight, VS., Very Slight, VVS., Very, Slight, M., Medium,

TABLE 161.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS -- PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, LIGCKPORT.

Report of Adolph Genemann, D. B. Bisbee, Municipal Laboratory, Chicago, III.

No. of Bae.	Der Cubic Centi: meter:	100,000	150,000	280,000	260.000	520,000	287,500	62,000	100.000	- 4 - 4 0 7 0 4	72.500	45,500	19 750		35,000	1.65°E	33.350	28,650	24.550	24,750	17.000	
	Presenc to set		:+		-+-	+	+	1	 - 	-+	+	-+	1	:	+	+	+	+		- :	+	
	Tempera of Air	86.5	000	6 6 6	200	27.	•	ਰ ਵੇਜ਼ ਵੇ	. 0	26.		92	- 	:	÷.	33	<u>8</u>	30.		<u>∞</u>	<u>∞</u>	:
	Tempers	33.		26.	100	33.	:			: ଗ	18.	15.5	<u>-</u>		Ç.	00:		6	00	1	∞ ∞	:
	Heigh ots'//	-		_	_	-	:	÷;					:	:		_:	+-	_:	_			:
RO- N AS	Nitrates	.45 52	2,8	69	0.	02.	0.	07.	2	06.	- 40	.30	:	:	0	၁	0.	0.	30	0.	.55	:
NITRO-GEN A	Setitiv	.013				٠							٠	:	.00.	:	0.	10.	.013		.005	:
IC FEN.	pəpuəd sns	.03					٠											ď			30.	
ORGANIC NITROGEN.	Dis- solved.	8:0								-											39.	:
52 	Total.	. S. S.	80.	8	.46	88. —	æ;	7.8	0 0€ 0 0€	1.1	16.	: 23		:	7.	<u> </u>	.58	1.8	7.	.62	.64	:
on .	my pud -sus	10.	38	80	80.	90.	91.	3.	03	3	.0.	.016	3	30.	3.5	60.			80.	20.	3.	:
TROGEN AS	Picor Dis-	.55. 07.		84.	84.	.52	.456	77.	0 0 0 0	.53	.43	.40	<u> </u>	#.	4.		.36		.36	86.	801	:
NITROGEN AMMONI	I'10'I	.56	37	.56	.56	.58	.616	5.0	5.00	.55	.50	.416	++.	94.	<u>×</u>	=	36	.56	#.	.30	£.	:
	Ргеедие. тонів.	80.	30	80.	80.	80.	80.	3.8	000	3	.048		80.	3	+£0.		80.	<u>. 1</u>		10 .	.0	
ED.	BySuspd Statter	्र द	00	-				: -	্ কা							:		_				:
ONYGEN	By Dis-	13.6			27	11.0]: []:	2: -	13.2	2	11.6	4.6	ος ος	9.4	G	0.00	∞ ∞	<u>∞</u>	7.	<u>}</u> -	6.3	:
000	Total.	13.8	5.	13.8																		:
re.	Chlorin	6.3	, m	4.3	5.1	3	က် () -	11.1	9.0	00	о. С	; {~	₹~ : :0:	9.6	.⊥ .⊥	6.7	} }	12.3	14.8	16.3	17.3	:
ON ION.	popuod -sus	13.	, 20 20 20 20 20 20 20 20 20 20 20 20 20 2	14.	∞ ∞	- ;	:3	×i⊆			ئىن.	10.	:		· ·	ó	19.	4	-	10.	-	
RESIDUE ON EVAPORATION	Dis-	383.	190	309.	337.	335	308	30 1 .	25.55	309.	287	317.	343.	372	851.	365	415.	.398	165	449.	117.	:
REG	T'otal.	396.	368	323.	345.	339.	308	2000	33.	308	294	337	33%	37.1	308.	373	434	30%	169	459.	161.	
	Odor	None	:	7,7	9 8	:	: :	: :	Gassy	None	. ,	*	* .	;			9 9	;	9.9	;	:	:
	Color.			:: ::	寸.	.e	<u></u>	:			:	:			:	:	:	•	•	•	:	:
APPEARANCE.	Sedi- ment.	VS.		'n	VVS.	1.7.5.	S. S.	27.7	· · · · · · · · · · · · · · · · · · ·	VVS.	VVS.	V.V.S.		VVS.			VVS.	VVS.	VVS.	VVS.	VVS.	
AV.	Turb'y.	VVS.	N. N.	V.S.	VVS.	VS.		21.2		VVS.	VVS.	VVS.	VVS.	VVS.	2.7		VVS.	VVS.	VVS.	VVS.	VVS.	•
DATE OF	1899 1899 Collec- Examition.	July		**	:	Aug.	: :	: :		11	.:	35	2 Oct.	5.		:	30 Nov.	,, 9	13			:
		glul'g	:	::	: 1	19 Aug.	: : []	: :	176 Sept.	8	: 08	: 25	84 Oct.	: 98	20 9	;	26	Nov. 10	,, 96	,. 86F	., 10	:
11 27	Serial								-		+	7	*	andra .	uga v	-	7	Traps.	+	-	1001	:

M.. Medium. S. Slight. VS., Very Slight. VVS., Very, Very Slight. APPEARANCE-H., Heavy. VII., Very Heavy. VVH., Very, Very Heavy.

TABLE 162.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER. NORTH OF JACKSON ST., JOLIET.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, 111.

No. of Bac.	Cubic Centi: meter.	55.000 385.000 385.000 280.000 197.500 385.000 385.000 385.000 385.000 385.000 385.000 385.000 385.000 385.000 385.000 385.000 385.000 386.000
	Presenc To .sd A	+++++++++++++++++++++++++++++++++++++++
	TeqmeT	<u>भैवव्यक्षित्वर्षक्ष्यम्</u> यव्यव्यक्षित्वर्
	TeqmeT of a Value	श्चित्र संश्चित्र श्चित्र च स्था श्चित्र च ने ल ळ च
	dgioII otaW	
RO- N AS	sətertiN	
NITRO- GEN A	setritiv	88.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0
IC FEN.	papuad -sus	80868 : 55-8 : 426 : : 58886 558
ORGANIC NITROGEN.	Total. Dis- Solved.	8. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4.
	a p,pud	842846888=888 188598458
N AS		88888888888888888888888888888888888888
NITROGEN AS	Albuminoid b'vios	88488848888488884888888888888888888888
NIN	.sinom	4888588555555
-	Matter. Freedm-	
GEN	Bysaspd	00
OXYGEN	Total.	40000000000000000000000000000000000000
	Chlorin	<u>8888888888888888888888888888888888888</u>
	1 .	
S ON TION.	pepued 	
RESIDUE ON EVAPORATION	Dis-	
Ev Ev	Total.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	Odor	Gassy Gassy Gassy Gassy
	Color.	Muddy 5 5 Muddy 3 3 3 3
APEARANCE	Sedi- ment.	日本語を表別を表別の名がある。
A V.	Turb'y.	
O.P.	1899 Exami- nation.	2017-2017-2017-2017-2017-2017-2017-2017-
DATE	1899 Collec- F	
70.	Inirad	88888888888888888888888888888888888888

M., Medium. VVS., Very, Very Slight. S., Slight. VS., Very Slight. VII., Very Heavy. VVII., Very, Very Heavy. APPEARANCE-II., Heavy.

TABLE 163.

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-DESPLAINES RIVER, SOUTH OF TOWN, JOLIET.

Report of Adolph Gehrmann, D. R. Bisbee, Municipal Laboratory, Chicago, III.

No. of Bac.	per	Centi- meter.			960,000									
ce or Coll.	nə	Prese	+	+	+	+	1	-		1	+	+	1	+
		dm9T A to	2.i.	19.	19.	53	<u>∞</u>	18.	-1	18.	<u>∞</u>	17.	22	11.5
ature er, C.	er ete	gm9T 37/ lo	33.55	31.	25.	ક્ક	33	26.	33	33	233	રેરે	17.	18.
		ieH W	:	•	:	:	:	:	:	:	:	:	•	
-03 N A S	sə	Mitrat	0:	82:	.26	.05	.40	88	.40	200	.0±	:	:	.18
NITRO GEN	sə	Nitrit	:		50.	:	0:	.018	.007	.023	.065	.035	0,	<u>8</u> .
NIC OGEN.		sug			1.73		3.40				-	3.24		1.60
ORGANIC NITROGEN		Dis.	05	33	1.28	1.48	1.48	1.48	1.60	1.48	88.	.56	1.52	1.80
OR	_	Total	58	38	3.00									3.40
	Am	p,pud -sns			.87									
EN AS	Albuminoid	solv'd	1.20	1.40	.63	1.00	<u>8</u> .	90.1	1.06	1.14	1.00	09.	1.06	1.16
NITROGEN AS AMMONIA.	Albur	Lioti	2.13		1.50	3.00	2.24	9.60				2.00		
Z		TreeAr inom	7	08	7.00	_		_	0f	09	99	8	40	9
- <u>.</u>	ı.i	BySus	5.8	8.1	13.	3.5		.9	5.4	7.	· 63	50	5.4	8.8
OXYGEN	·pa	evios	_ CS	7	3	00	es.	7	7	C.	03	T	03	35
OX	-	Total		03	.2 16		63	T.	00	35	7	+	9	
] '	[010[]	30		. 39									
16.	LĮI	СРЈО	117	105	85.	16	108	138	111	117	119	131	115	=======================================
ON ION.		sus Sus-	78.	49.	141.	130.	196.	174.	æ æ	217.	165.	94.	46.	54.
RESIDUE ON EVAPORATION		eiu Solved	483.	551.	441.	536.	493.	557.	513.	483.	514.	507.	157.	427.
RE Eva	.1	[B10']'	561.	900	585	656.	689	731.	601.	700.	679.	601.	503	481.
	Odor		SlGas'y	"	;	Gassy	3	**	7.9	",	**	**	3,0	:
		Color.	Muddy	:	.45	ું.	. 15	က	.85		:	:	:	
APPEARANCE		Sedi- ment.	H.	M.	H.	H.	H.	H.	M.	H.	VII.	VH.	ś	VVS.
API		Turb'y.	VV1II.	V.H.	V.11.	1.1.11.	VH.	V.V.H.	V.H.	VH.	VH.	Š	VIII.	VVII.
3 OF	1800	Collec- Examition.	3 July 3	101	17	÷~	31	Aug. 7	77		:	Sept. 5	11	18
DATE OF	1800	Collec- tion.	July 3	01	: 17	: 31	31	Aug. 7	11	₹ :	86	Sept. 48	-	.: 18
				_	_	_	_	_	_	_	_	_	_	_

M., Medium. VS., Very Slight. VVS., Very, Very Slight. S., Slight. VH., Very Heavy. VVH., Very, Very Heavy. APPEARANCE—H., Heavy.

TABLE 164.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Adolph Genemann, D. B. Bisbee, Municipal Laboratory. Chicago. Ill.

SOURCE OF WATER-KANKAKEE RIVER, WILMINGTON, ILL.

4000].c	er.	9,350	:	1969	3.000		:	1.750	1.X(E)		3.000	5.000	11 000	1.350	:	1.9(0)	19,500	5.000	:	0.000	30.000	5.250
No. of Bac.	Cab Cab	met		:	₹7 T	•	:	:	?	-		**	41.	_	_		***	3:			400	×.	•••
(,01f.			!	:	+-			:		+				;		:	+-	+	1	:	7	+	4
ature , U.	aədu uber		81	18	<u> </u>	9 9	<u>∞</u>	17.	55	900		16.	16.	16.	10	ž	19.	<u>x</u>	17.	<u> </u>	; -	īĞ.	0 -
ature n, C.			FG :	£	31 7	· ·	?}	3	99	왕.	23 33	19.	50.	11.	оċ	<u>.</u>	19.	-	10.	.c.	ië.	œ	iG.
	dai9 otran		. 833	<u>.</u>	<u>ء</u> ن	; rq	-	-							œ.		-	-		ë: I	<u>.c.</u>	<u>∞</u>	1.6
RO- N AS	หอ1 <i>ธ</i>	MIL													۰					0.			
NITRO-GEN A	soff	nin			690.				-											0.			
GEN.	geq -st	18	3 . 15																				
ORGANIC NITROGEN.	-si	n			.58									$\overline{}$									
	a b	pud	·		20:	_	_	-			_	-		harmag			-	_	-				
r AS	[24]	Alos							_	_					•	_		-			-		
NITROGEN AS	1bum	10T			01.				-	_				_	-	_							
N	-mkg .sina.	out		-	.07		-	_	_		_	-	_	:		-		.0:0		016			
	ter:) Isi			∞. ∞. s	3 31		00	_	{ ~	33	@ ?	-	•					- 1				
OXYGEN CONSUMED.	D97		11.9	∞ ∞	9 11	2 00	2	oc. ₹~	oc.)C	-3	-}	Ğ: 9	8.9		5.6	9.1	4.6	1G		07	en oc	65 65 65
65 —	.IEI	0,7,	11.6	=	₩. 0.3	2 =	12.6	11.6	10.9	13.4	9.01	9.6	10.	ж ж	6.9	6.7	9	ω. Ω.	 	5.6	0 1-	oc oc	0.01
16.	loriı	СР	2.6	35 00	e) :	9 99	65	3.6	50	species - -	OC	oc.		5.1	9.4.	4.6	7	4.6	÷	-	oc.	??	3.0
ON ION.	pəp -sı	nəđ 18	16.	-	ල	. 66	67.	70.	Ţ.	67.	7.	35.	9	36.	œ.	33		36.	??	-		·	12
RESIDUE ON EVAPORATION	is-	(I	.380	27.5		9 19	25	949.	253.	35.5	254.	200	53.30 	219.	245	ox.	257	1307	÷1 ≎1 ≎1	200	321.	313.	307
EVA	.fBl.	од	296	316.	363	35.1	310.	319.	324	319.	335	207	500	150	763	130 ·	300	300	150	27.0	323.	334.	35.5
	Odor		None	9.9	-A - 19	:	3 3	:	9.9	;	:	:	:	:		0 79	;	•	:	:	•	:	:
	Color		ೞ			9 19	÷ 00	100		•		:		:		:			:		:		:
APEARANCE	Sedi-	ment.	V.S.	M.	Ä,	1.1.1	1.7	1.7	1.7.5.	1.5.	N.	5.7	1.5	5.1.2	ジンン	7.8.	1.5.	1.5.	1.7.5	1.1.5.	1.7.5	1.1.8.	5.7
11.	Turb'y		1.5.7	Ĵ.	: X:	, II.	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	V.11.	-	VII.	point invest	- Period		1	5.7	M.	M.	M.	M.	1.7	7.7.7	5.7.7	1.1
OF	1890 Evami.	ation.	July 3	9	:	5 55	112. 7	=	100	2007			<u>×</u> :							ior. 6			17
DATE OF	1899 1890	on. n	_0.5	0		57 67	1-	upolic beauti	50	¥.1	it. 4 Sept	=	y	19	67	C.	16	3	08:	0	155	()(-	7.2
	2 0		O July	;	27.0	:	5 1111	., .,	**	:	/		;		r,	-		:		10% ×	-	:	30
.0%	luin	18	00	天	UC U	, X	K	X	X	X.	1	J.	5.	3	4.	0.	2	₫.	Ç	Ö.	C:	10	127.8

M. Medium.

S., Slight VS., Very Slight, VVS., Very, Very Slight,

VIE. Very Heavy. VVII., Very, Very Heavy.

APPEARANCE-II. Heavy.

TABLE 165.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, MORRIS, ILL.

Report of Adolph Gehrmann, D. B. Bisber. Municipal Laboratory, Chicago, Ill.

of .c.	bic otti- ler.	19.650	000.00	0.000	(S)(S)(S)	0.000	36.500	33,750	000.0	000.0	7.000	0,000	91.500	39,250	6.250	65.000	067.71	000.73	33.250	16.000	2.500	
No. Ba	per Cubic Centi- meter.		7.	-	=	7	• * *	:3 :3	<i>5</i> .	ॉट -	Ğ.	3	1		14.	7.	6.	5	ক	*1	_	
	Presenc	11	+	+	+	+	1.		+		+	-	+	+	+-	+	-1	+		-}-	+	
	Temper	15.0	-81	2) 2)	<u>00</u>		÷ ;	6.	į -	+	16.	=	10.	2	16.	91	?	+	{ -	-	Ť	
	Temperate Vate	8.5	24.	27.		33	- - - - - - -		36.	33.	30	13.	17.	10.	30	15.	10.	16.	_ ∞	6.	6.	-
	Heigh Wate	34.03 29.53	<u>1</u>	6.3	Ģ.	6.3	ιυ : ∞ ·	5.	4.1	4.1	ت. ص	ت. زه	5.6	5.4	5.6	5.6	5.1	5.6	6.3	6.3	8 3	
-01 I AS	Nitrates	85		.67	67.	02.	S;	.40	<u>. 15</u>		♂	0.	98.	80.	. 15	0.	9	17	1.64	19.1	11.1	
NITRO- GEN A	Nitrites	9. E.E.	.11	80.	10:	.01	+ 00.	.017	1.08	.075	91.	.37	114	<u>8</u>	.052	.013	800.	.03	.065	60.	60.	
EC.	pəpuəd -sns-	32.															_	$\overline{}$				
ORGANIC NITROGEN.	Dis-	1.16	88.	.76	1.08	1.20	86.		1.34	1.08	2.10	1.24	1.02	1.02	.80	1.22	1.50	1.16	1.38	1.18	正	
OR	Total.	1.56	1.16	80.1	1.62		.68	. 68	1.74	1.74	2.60	08.1	1.62	1.76	1.02	2.08	2.58	2.24	2.04	1.54	2.30	١
	E p,pud	.13																				
EN AS	Albuminoid	£.5	0f.	99.	٠ <u>.</u>	.67	<u>6</u>	99	∞.	.67	77	:69:	69.	99.	.56	02.	88.	99.	99.	.58	.76	
ITROGEN AS AMMONIA.	Tol.1	86.86	96.	89.	8; 8;	1.03		<u> </u>	1.10	8.	1.40	33	66.	1.00	99.	1.44	1.50	1.10	90:	88.	1.26	
Z	FreeAm- monia.	8.71																				
SD.	Bysuspd - Matter	8.5					no 0													3.6	6.4	
OXYGEN	By Dis- solved	€: 1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.		00			 													<u></u>	∞ ∞	
OX	Total.	15. 1 16.4	च	00	7	35	٠ ر	21	00	ा	9		00	?!	_		T	_		000	<u>~</u>	
.9ı	Chlorin	57.					_	_						_					_			
z ż	pəpuəd Sus-	9.					9 9														:	
RESIDUE ON EVAPORATION	Dis- solved.	380. 450.	327.	439.	417.	428.	395.	 	452.	437.	429.	408.	446.	136.	331.	390.	386.	378	380.	382.	417.	
RES	Total.	389.	380.	448.	428.	423.	401.	413.	463.	451.	145.	417.	455.	459.	343.	403.	397	405.	401.	409.	410.	
	Odor	None	"	9 9	-d -e	*	3 3	:	None	9.9	9 4	;	7.7	3.9	97	*	10 10	7,	Slight	,	•	
	Jolor.	ç; 4	-		왕.	ಲ	က့	:	:			•	•	•	•	•			-			
CE.					-		-	:	•	•	:	:	-	:	:	:	-		-		:	
APPEARANCE	Sedi. ment.	V.S.	M.	VS.	VVS.	VVS.	VVS		VS.	M.	VVS.	VS.	Ś	ś	ś	VS.	M.	V.S.	M.	M.	M.	
41.	Turb'y.	VS.																			MH.	
OF	1899 Exami- nation.	uly 3						۰													86	1
DATE OF	1899 Jollee- E tion. n			33																		1
No.	Serial	106 July	108	109	110	111 Ang	113	113	., †11	115 Ser	116	117	118	119 Oc	120	121	193	123 No	124	3	353	-

M., Medium. S., Slight. VS., Very Slight. VVS., Very, Very Slight. VH., Very Heavy. VVH., Very, Very Heavy. APPEARANCE—II., Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-FOX RIVER, OTTAWA, ILL.

Report of Adolph Gehrmann, D. B. Bisber, Municipal Laboratory, Chicago, Ill.

No. of Bac.	per	Cuore Centi- meter.	12.350	13.500		11.250	13,000	13,000	906					0000	19, 950) (m 1 m 1	20.650	00.01	10,000	000.15 11 500	005.01	18,500	
		Prese Abs. c		+	:	+								+	-		-			- -	-		
		qmə'l' A lo	35.		33	56. 56 5. 56 56 56 56 56 56 56 56 56 56 56 56 56 5	1 22	2	33	37.	31.	500	3	6	0		96				<u>.</u> S		
		qm9T sV/ lo	88	∞	:36:	83 P	1	.98	3,	31.	.98	18		18.51	16.		: (-	10	13		15	1-	_
		라하 당기/ 	00.	20° I		က်စ	÷ -1	3.6		e5.	3.6	-	-	-		:		10			. 2	3.00	
30- N A S	80	Mitrat	88:	£	1.07	<u> </u>	2 G	0F.	.30	08:	02.	02.	0.	01.		_	•	9	10	-	9	23	
NITRO- GEN A	80	Nitrit	l																			3	
ic EN.		opuod -sng	80	97	8	99	06.	10	.93	=======================================	25	10.	.e.	10.	8	;		2	10	8	2		
ORGANIC NITROGEN.		siU solve		_		3 %																38	
ÖZ	1	R10T	.52	30	1.56	2 3 - -	5 3c	.68	.76	.68	. 7.3	3 .	39.	.58	3		10	9	6:1	15	101	<u></u>	
on.	d Am	-sns	.01	3	14.	:e0	0.0	01.	ō.	.05			<u></u> 30.	90.	0.					0.1			
NITROGEN'AS	Albuminold	-si(I b'vios	.31	999	₽;	99		.36	01:	=	٠ <u>.</u>	88.	.34	.30	8		33	35	3.5	100	200	30	
NITRO	Albu	Toty	88.	89	<u></u>	<u> </u>	· +	. 46	=	94.	£.	œ.	38.	38.	.31		33	~		O _G	00	× .	
		Ргеел. пош	<u>21.</u>	3.8	80.0	5,8	9.6	80.	.01	0.	2.	910.	20.						0.03			:8	
ZD.		enska Bysus	-		9) (1)	:10	0.	-	1.33	**	-:	?}	¢ć.	-:		600				3	-	
OXYGEN	p	By Dis				~ _ _																1-1	
Con	-	Flota I				ρ: ο Ο α				<u>.</u>	- 0		7.6									6.1	I
16,	ijΙ	Срјо	37°1	0,0	D .	i 4	6.7	6.4	5.6	6.3	7.5	5.6	6.1	0.8	œ.		6.7	7.	7	1 -	6.5	8.9	
ON.	pa	penaq sus	6.	3	3	13.	-	***	: ::	<u>x</u>	20		10.				7	•		4		6.	
RESIDUE ON EVAPORATION	E	Solve	300.	200	2003. 0.000.	2 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	289		X X	- X	300		280.	298	335		319.	337.	291	298.	306	305.	
RES	1.	rio'l'	306.	500°	416	250	303	305.	303	300	55.3	25 26	290.	31.	313.		326.	33.5	291	305	306.	308.	
	Odor		None	7 7	, ,		9 9	9.9	9 7	4	,		9	9	9 9		None	-0	9 9	9 9	9 9	**	
		Color.	- 3	9,	e e	<u> </u>	-E	. 5				:		:									
APPEARANCE.	;	Sedi- ment.	V.S.	.00.1	= 0	7.7.8	VVS.	7.1.S.	VVS.	VVS.	s.	.3.	S.	V.V.S.	1.1.8.	V.S.	VVS.	VVS.	1.7.5.	7.7.7	1.7.8	1.7.8.	
Ψ		Turb'y.	VVS	6.0	0.5	i .i.	17.8.	·s.	None	3.	5.7	1.3.			5.1.2	V.S.	VVS.	1.7.8.	1.1.5.	1.1.1	1.1.2	1.7.8	
OF	1899	Exami- nation.	11/7 3	10	0 10	ug		15				<u>~</u>	61 13		ct. 3	9 5	2 12			10v. 7	11	*	-
DATE OF			S.July	1		7	- }			X	- Sept		α	20	200	00	6			1	13 ,	1-	1
DA	1899	Collection.	July		9.0	: ::	Aug.	:	:	:	sept.	:	:	:	Oct.	**	7,7	3		Nov.		33	
.07.	[13]	198	134	120	14:	2	111	146	*		1.	Z.	艾	200	XOX	CX	01 02	814				1304	

M., Medium. VVS., Very, Very Slight, VS., Very Stight. S., Slight. VVIII., Very, Very Heavy. VII., Very Heavy. APPEARANCE-H., Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

SOURCE OF WATER-ILLINOIS RIVER, OTTAWA, ILL.

No. of Bac.	per Cubic Centi: meter	88.000 138,500 138,500 13,000 100,000 100,000 19,13
	Presen lo .sdA	1 +++++++++++++++++++++++++++++++++++++
ature, O.	ragmaT riA to	· · · · · · · · · · · · · · · · · · ·
ature 17, C.	Temper of Wate	<i>१११</i> १९ १९ १९ १९ १९ १० १८ १८ १८ १८ १८ १८ १८ १८ १८ १८ १८ १८ १८
	Heigh Wate	• • • • • • • • • • • • • • • • • • •
RO- N AS	Setrativ	
NITRO- GEN A	Nitrites	KK + 3 & YK & 8 8 8 4 4 8
NIC GEN.	Sus-	0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.
ORGANIC NITROGEN.	Dis-	S
	Total.	28552980842559
A8	A sus	
NITROGEN AS	Dis-	\(\frac{1}{2} \) \(\frac{1} \) \(\frac{1} \) \(\frac{1}{2} \) \(\frac{1}{2} \
NIT		3398883: 8888888848884888
	FreeAm-	www.q-q-w-q-w-q-w-d-d-y-w
EN MED.	BySuspd Matter.	\(\phi \) \(\phi \
OXYGEN	By Dis-	_ <u>බණ්ණක් 4 ක්ෂණක් කික් මේ මේ ක්</u> මුණුමුට්ට්ටුමුට්ට්ටුමුට් මුමුණුමුකුට්
	Total.	888.888.888.888.888.888.888.888.888.88
	Chlorin	
ON FION.	-sus	0,400040 1400 1200 1000 1000 1000 1000 100
RESIDUE ON EVAPORATION	Dis- solved.	29.5. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.
RE EV	Total.	88.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8.8
	Odor	None
	Color.	gi - 1, g
APPEARANCE,	Sedi- ment.	
IV	Turb'y.	
DATE OF	1899 Exami- nation.	3 July 3
DAT	1899 Collec- tion.	July Sept. Coct. Nov.
.o.Z	Serial	######################################

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VH., Very Heavy. VVH., Very, Very Heavy. APPEARANCE—H., Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-BIG VERMILION RIVER, LA SALLE, ILL.

Report of Abolph Genraann,
D. B. Bisber,
Municipal Laboratory, Chicago, Ill.

No. of Bac.	Cubia	Centi:	11.250	15,350		96.250	000.00		20,000	1.600	28.333	31.000		30,750	39,000	25.750	0.000	20,000	21.000	21,500	99,750	32,350		16,650	
ce or Coll.		Pres. o	1						ļ.,	+	+	÷		٦	+	t	:	+						+	-
		qməll 7. 10					:																		-
		Temp T			:		:								:										
10 1		Heli I'W			11.6							5.				0.0		9.7							
NITRO- GEN AS	R9	Mitrat	_	-	30.00		_																		
	_	Nitrit			100																				:
IC FEN.		sus	8																				-		:
ORGANIC NITROGEN.		Dis-	.33																						:
ON.	1.	вюТ	.38	.68	00.1	0f-	5.6.	689.	69	.76	17.	.70	32	08.	<u>√</u> .	31.00	5.	. 1.	$\frac{\infty}{\top}$.56	:36:	3.		.30	
	1 Am	p,pud -sug	80.																						•
NITROGEN AS AMMONIA.	Albuminoid	Pis-			5.5																				
NITRO	Albu	['10T]	.24	88	11.	7주:	\$? .	33	305	:ic.	=	04.		88	.30	7!	200	61.	જ	£.	.16	<u>= :</u>	.16	<u>=</u> :	
		Ггеелі іпоят			5.50																			3.	
ED.		BySus	_:	0.4	1.4	ŧĊ.	3.		\$?	30	33 33	33.	35 35	1.9	G.	es es	-	;;	- 1		7.	7.	9.	:	
OXYGEN CONSUMED.		solve solve	හ දෙ	- j	∞ 1~		oc ∞	- -		÷.	- 1	50	- j	.C.	-	50 53	33.	e:	6.5			33			:
Cox	-	Total			6.6							∞ 				5.4	<u>; -</u>		9.4	4.1	G	⊕1 ⊝C	3.1	53	:
to:	ri.t	СРБО	÷.	10.	18.5		.0.	5.	533.51			68.5				13.0		138	110.5		121.	131.5	±	105.	
ON.	pi	snS	33	 	39	19.	÷0	65	.0 .	- j	.j.	76.	333	.88	130	÷0.	30.	11.53	. <u>i e</u>	33	16.	34.		.01	
RESIDUE ON EVAPORATION	1	sid earlos	684.	314.	. S. S.	513	708.	917.	508.	561.	(35).	973	1101	1191	1713.	1619.	1702.	1591	1567.	1611.	1645.	1751.	1336.	1581.	
RE		Potal	706.	352	101	535	-1 10c	283	63.X	638	802	1049.	11:31	1530	1785	1659.	1831.	1631.	1618.	1675.	1661	1788	1327	1591.	
	Odor		None			:	:	;	:	:	:	:	:	ig e	:	:	:	. 9	:	:	:	:	9 •	:	
		('olor.	0.	- G.	. (c).	(F)	0.	.05	.65																-
APPEARANCE.		Sedi- ment.	j.	1.7	N.	N.	j.	7.7.7	M.	M.	N.	= -	N.	N.	M.	s.	ジン	M.	ŝ	1.7.5.	1.7.5.	1.1.5.	1.1.5.	VVS.	
11.		Turb'y.	7.7	3.1.7	·j.	1.7	j.	N.	N.	N.	M.	N.	N.	j.	N.	M.	M.	M.	M.	5.11	5.1.1	5.1.1	5.7	ジンン	
2 OF	1900	Exami- nation.	July	:	61	97	AHE. 2		165	??	**	Sept.	1	:			.9.	:	:	10%	:	:	:	£	:
DATE OF	1500	Collec- tion.	July 4		J	·	VIII. 1	3	:	***	06	Sept. 5.				Det. 3				16. 31	1- 20%	11	:	· ·	
l oz	[11	rios	169	57	X X	522:	37.5	5	大人.	364	3,64	397	400	103	11.63	100	07	10	X	4.51	XIX	- X	15%	1387	

M., Medium. VVS., Very, Very Slight. VS. Very Slight. S., Slight. VVIII. Very, Very Heavy. VII., Very Heavy. APPEARANCE-II. Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.
SOURCE OF WATER-ILINOIS AND MICHIGAN CANAL, LA SALLE, ILL.

Report of Adolph Gehrmann.
D. B. Bisbee.
Municipal Laboratory, Chicago, III.

No. of Bac.	Cubic Centi- meter.	69.000 60.000 60.000 60.000 60.000	
	Presen forselv	+!++++ ++ ++++++++++++++++++++++++++++	
	rik to		
	Temper of Male		
	figi9H etaW	co.T.c.c.c.c.c.c.c.c.c.c.c.c.c.c.c.c.c.c	
NITRO- GEN AS	Sitrates	8239080	
NITRO	Nitrites	68.888.98.58.58.58.58.58.58.58.58.58.58.58.58.58	
GEN.	pepued -sng	- 188	
ORGANIC NITROGEN.	-si(l solved.	24255242455545555555555555555555555555	
- 7.	pnd'd "	0888848484868454448888888888888888888888	
sa	Sus-	<u> </u>	_
TROGEN AS	Albuminoid -sid b'vios	44,58,86,46,45,86,86,88,86,88,88	
NITROGEN AMMONI	[[1,10,L	868888888888888888888484	
	FreeAm-	<u> </u>	
ED.	BySuspd		
OXYGEN CONSUMED.	By Dis-	ထလ္မမႈ-၄	
၁	Total.	047-808 0 0 0 0 0 0 1 1 0 0 0 0 1 1 0 0 0 0	-
.ər	Chlorin	3 2 2 3 2 2 2 3 2 3 3 3 3 3 3 3 3 3 3	
ON ION.	pəpuəd -suS	<u> </u>	
RESIDUE ON EVAPORATION	Dis-	######################################	
REEVA	Total.		
	Odor	None : : : : : : : : : : : : : : : : : : :	
***	('olor.	15-16-1919-	
APPEARANCE	Sedi- ment.		
AP	Turb'y.		
DATE OF	1899 Exami- nation.	1	
DAT	1899 Collection.	July. Aug. Sept. Sept. Nov.	
.oV	Serial	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	

M., Medium, VVS., Very, Very Slight. VS., Very Slight. S., Slight. VVIII., Very. Very fleavy. VII., Very Heavy. APPEARANCE-II., Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-ILLINOIS RIVER, LA SALLE, ILL.

		III.
ż.		Chicago,
H GEHRNANN.	BISBEE.	Laboratory,
ADOLPH GER	D. B.	unicipal
Report of		Mun

No. of 13ac.	Per	Centi: meter.	000.4	40.000	13.650	14.0(x)	18.650	68.000	11.350	21.500	(000)	21.000	77.5(0)	86.000	41.750		66,000							59.950	
10 99 (10)		Prese	-	-	+		+	.	+				t	+	-	1		+	-+		+	- 1	+		-
		tineT'								•															
971176 ;) , 79	119 911	qmoT sW lo		:		:	:	:	•	•						:						-			:
lo t	916 ZD:		9.1	6.6	11.6	9.11	9.5	£. €.	6.		9.4	ñ. 6		1.6			6.				9.11			? OI	:
0.0-	89	Mitrat	3.05	3.95	3.06	1.70	30.5	3.5	名 ??	€.10	9.50	1.95	1.G	1.37	27.15	9.	- :	1.11	1.51	?!	.76	1.31	2.01	1.9	:
NITRO- GEN A	sə	Mitti	:8:	555	11.	.40	£.	57.	02.	12	£.	1.10	.65	£	30	ã.		38:	.36	97	11	.16	38:	96	:
IC SEN.		pund- sns	.16	₹?	₹ ?:	:96:	.70	33	07.	.25	.26	97.	90.	1	×.	10.	?!	023	90.	3	.:30	:	10		
ORGANIC NITROGEN.	1 ~	sid			98:		-		-								-					_			:
ÖZ	1.1	RIOT	1.34	1.16	는 왕. 당.	1.64	30.53	1.16	S: 1	1.18	<u>×</u> : =	1.06	8.1	1.01	3.	30.1	1.26	원. -	× .	3:	.93	2	1.0.	3	:
pro	d Am	p,pud -8ng	08.	+-	.26	?}	0ge .	.O.	.136	95.	91.	유 주	.10	≅.	910.	90.	30.	01.	3	80.	:	3	80	60	:
NITROGEN AS	Albuminoid	Pis-	.40	<u>x</u>	38	34	9.	=	2.	24.	77	<u>=</u>	01.	=	=	90	<u>∞</u>	23	9	9	01.	::	23	3C	:
NITRO	Albu	1.10.L	09.	63	19.	56	0,7	81.	.536	.73	09.	33	500	<u>«</u>	156	56	.50	300	===	oc.	9.	.36	110	01.	•
		Freev	1.10	8. 8.	3.00	80.		<u>~</u>	9.	06.1	1.70	20.8€	4.10	4.62	6.33							-		3.60	:
N. ED.	ı. bd	Brsus	2.6	ن د	∞ .∞	4.6	3.6	∞ ?ì	<u>~</u>		9.1	₹.2	-	\$1	9.	9.	??	23	9.1	-	7	œ.	9	?!	
OXYGEN CONSUMED.		By Dis	1.1	10.8	8.6	9.4	9.4	9	c:	8.6		10.	9.1	.0:		9.1	ж. С:	φ. ∞	7.6	7.1	· ·	7.6	30	oc.	
000	1	R10T	-	∞. ??	13.6	14.		11.4	10.8	=	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	13.1	∞.01		ος σ:	10.	=	$\frac{\infty}{2}$	C:	œ.	×	oc	φ α	0	:
.9t	ПİЛ	орро	#	63.5	30.	35	<u>-15</u>	17.5	50.00	(F)	-J.C.	7.0	£3.5	15.5	10.00	Œ	76 5	680	59.	53	55.	31.	. C.	339	
2N 10N.		ppuad sus	35	£3	00 10	. 25.	92 93	49.	, , ,	27.	12	56.	<u>∞</u>	+0.	.65	∞.	??	∝.	-:-	-	10.	=		οċ	
RESIDUE ON EVAPORATION		oovfos	:389.	410.	313.	356.	10.5	381	385	395.	102.	115	5	395	380	416.	138	.886	369	:50.	:	326.	251.	37.5	:
RES	.1	rio'l'	5	+333	371.	+533.	470.	430.	=======================================	155	120	501.	++3.	133	<u>x</u>	161.	150.	11.	38.	561.	385	340.	351.	<u>x</u>	:
	Odor		None	*	:	:	-0 -0	*	• •	* *		# #	**	:		1	:	*	*	4 .	* 9	*	:	1	
		Color.	8	98:		. 15	.35	. 255	25.		:	:													:
APPEARANCE		Sedi. ment.	S.	5.7	<u> </u>	N.	==	N.	S.	÷	·ŗ.	=	. N.	M.	Š	7.5.	ĿĊ.	Ż.	ý.	Ý.	17.5	・バンン	1.7.7	5.7	
1.		Turb'y.	7.8																		1.7.7	グンン	5.1.7	1.1.1	
DATE OF		Exami- nation.	t July 5																		α :	15	900	· ·	
DAT		Collec- tion.	July	:		:	1119.	:	:	:	:	Sept.	:	4 4	9 9	Jet.	:		:	:	Nov.	4 10	. ,	· %	:
.oV	18	Serl	170	173	376	ÎX.	33433	实											4119	155	1379	1386	13%5	13%	:

APPEABANCE-H., Heavy, VH., Very Heavy, VVII., Very, Very Heavy, S., Slight, VS., Very Slight, VVS., Very, Very Slight, M., Medium,

TABLE 171.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, HENRY, ILL.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

No. of Bac.	per Cubic Centi- meter.	26,750 12,500 12,500 12,500 333,750 12,500 290,000 50,000 50,000 13,550 11,550 11,500	
	Presenc	+++++++++++++++++++++++++++++++++++++++	
ature , C.	riedmeT rin lo	8	dium.
	Tempers	gi = 26.55.55.55.55.55.55.55.55.55.55.55.55.55	M., Medium
10 1	Heigh eta <i>W</i>		
RO-	Nitrates		VVS., Very, Very Slight
NITRO- GEN A	Nitrites	88888844888885 Signost	Very
nc abn.	papuad -sng	2	ery,
ORGANIC NITROGEN.	-siQ solved.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	S., 1
02	Total.	82.28.21.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2	1
	Bud'a	#858855884-67885858	ght.
TROGEN AS	Tor'i	<u> </u>	ry Slig
NITROGEN AS AMMONIA.	I'10T	885564658688888448848	VS., Very Slight
	FreeAm- monia.	<u></u>	
ED.	Bysuspd Matter.	<u>+</u>	Slight
OXYGEN	By Dis-	စ်စွေထွတ်တာ - စစ်ဝိဝိဇ္ဇေဇဇလတ က - စာ ∞ : စစ် ကို စစ်သည် သို့ ထိုလို စစ် ကို ထိုလိုက်	U.
Cop	Total.	85000000000000000000000000000000000000	
.en	Chlorit	850 950 950 950 950 950 950 950 950 950 9	Heavy
ON ION.	pepued Sus-	**************************************	Very
RESIDUE ON EVAPORATION	Dis- solved.	3866.6.6.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.	Very.
RE	Total.	28888888888888888888888888888888888888	VVH Very. Ver
	Odor	None	
	Color.	ej ej ai ai ai gg ai	VH., Very Heavy.
APPEARANCE	Sedi- ment.		VH.,
ΑV	Turb'y.		APPEARANCE—II., Heavy.
DATE OF	1899 1899 Collec- Examition.	28	RANCE-
DAT	1899 Collection.	July Aug.	APPEA
No.	Serial	88 60 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	

TABLE 172.

Serial No.

Report of Abolph Gehrmann, D. B. Bisner. Municipal Laboratory, Chicago, III. STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, AVERYVILLE, ILL.

No. of Isac.	per Cubic Centi- meter.	11.000 10.0000 10.000 10.000 10.000 10.000 10.000 10.000 10.000 10.00	9000 9000 9000 9000 9000 9000 9000 900	
	Presenc forse	+ + + + + + + + + + + + + + + + + +	++	
ornin O, C.	apduoj apduoj		≘ - :	dium
ornii .U ,u	Pempera of Wate		10.	M Medium
	Helgh otaW	त्रात्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त्रवात्त् 	81 87 81 87 81 87	
30- N AS	setantiv	84588222888885658558585 845888888888888886886	रू. <u>।</u> इ. । :	Slight
NITRO-GEN A	soffativ	ยู่ยู่ส <u>หน่า</u> ย่ากล่ายแล่ง : 8,7588	S 15 :	
HEN.	popuod -sus			ery.
ORGANIC NITROGEN.	-sid solved.		•	VVS. Very, Very
	Total.		•	=
œ.	a Sus-sus			ider.
TROGEN A	Piori Illouminoid	* 455%%%%% = %	## ## ##	ry Sti
NITROGEN AS AMMONIA.	['10']		.36	VS., Very Slight.
•	FreeAm- monta.	1 <u>1686438888885</u>	3.00	1
ED.	Bysuspd		= ∞ :	Slight.
ONFGEN	By Dis-		∞.5 5.5 5.5 5.5 5.5 7.5 7.5 7.5 7.5 7.5 7	<i>S</i>
93	Total.	33000000000000000000000000000000000000	∞ t- :	1.
.e	Chlorin	<u> </u>	3.50	Heav
ON FON.	penqeq gus-	T	70, 65	Very
RESIDUE ON EVAPORATION	-sid solved.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	368	Very.
Eva	Total.	在存货投资股票等的表现是 2.	365	VVII., Very, Very Hea
	Odor	None	· · · · · · · · · · · · · · · · · · ·	
Pa 1	Color.	ω', ''ÿ − − '; '; '; '; '; '		VII., Very Heavy.
APPEARANCE.	Sedi- ment.		VS.	VII.,
A.P.	Turb'y.	ॐळ <u>्</u> र्रेॐचंबंबंबवंबंबंबंबवंबवंबवंबवं	VS.	APPEARANCE-H., Heavy.
Es.	1899 Exami- nation.	5 3 3 · · ·	R R	1
DATE OF	2- Ex	Nov. : : : : : : : : : : : : : : : : : : :	- x	LEAN
DA	1899 Collec- tion.	July Nug Sept 88 88 88 8 8 8 8 8 8 8 8 8 8 8 8		APPEA
		I ENERGE STAN SHIP SHOP SHOP SHOP SHOP SHOP SHOP SHOP SHO	X 82 ·	

TABLE 173.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-ILLINOIS RIVER, WESLEY CITY, ILL.

Report of Adolphi Gehrmann.
D. B. Bisbee.
Municipal Laboratory, Chicago, III.

No. of Bac.	per Cubic Centi: meter.	1,916.000	5,510,000		500.000	1 000 000	1,020,000	240.000	1,550,000		5,000,000	77,500			
	Presend to .sdA	+1	+		++	- :	- : - :		-11	-	+		:	:	
	remperative Temperative Temper	36.22				888	; ∞	- FG	£ 61	15.	15.	16.	16.	<u>.</u>	
ature D. T.	regmer etavi lo	33:		35.		88 8		13.	2110	16.	10.	6.	10.	 •	
10 1	Helgh Hate	6.	9		6.7	: -		<u>.</u> ;		-	1.5		c) -	e. I	
-0.	Nitrates	2.93 88.23 88.23	8		8.3	88 9	06.	1.40					6) - 6) - 6) 1		
NITRO- GEN	setitti N	82.53	883	. 33	£ 60.	8	10	08.	Z &	96	1.44	99.	हैं।	60.	
IC EN.	bended Sus-	.13	80	•	89. 7										
ORGANIC NITROGEN.	Dis-	1.08	.76		8.8	-	-	,	- -	-	-		_		
ÖZ ——	Total.	1.36	8.	1:00:	1.58	1.55	1.42	1.30	1.74	1.84	3.30	1.54	8.30		
	p,pud -sng	8.8	80.	.12	.45	.55	.30	.36	£ 50	.40	1.03	.34	99:	7.	
AMMONIA.	Piori Piori Dis- Dis- Dis-	14. 44.	.36	.36	£;∓	<u> </u>	7.	.36	3.5	.48	·84	.55	∞ ;	17.	
NITRO	T'10T	.64	7 6	48	86.09		37.	£ .	£ ∞	88.	1.86	98.	1.44	- -	
	FreeAm- monia.	.03	42	1.10	99.	₩. ₩.	88	.16	2. %	1.24	1.30	2.80	3.60	5.74	
ED.	Bysuspd Matter.	1.6	4. :		9.89			က တ်					ट इ		
OXYGEN	By Dis-	11.4	∞ :	6.	10.	10.2									
Coo	Total.	55 55 55 55 55 55	∞ 4.		12.6				15.2						
'əu	Chlorin	8.2	31.5	31.5	4 4 .	73. 73.	56.5	63.	66 65 65 65 65	67.	67.	61.	55 11 11	6.16	
ON ION.	pepued gns-	37.	ફ્	12	## ##	87.	 31.	25.	8 8	ક્ક	46.	18.	11.	.68≈	
RESIDUE ON EVAPORATION	Dis- solved.	356. 381.	305.	347	377.	352.	359.	391.	306. 306.	404.	426.	395.	390.	991	
RE	Total.	400.	357		420. 381.	439.	390.	418	410.	429.	473.	413.	401.	200.	
	Odor	None	None	None	3 3	; ;	3	9 9	: 3	13	9,	9,	: ;		
	Color.	शंक्ष						•				:	:	•	
APPEARANCE.	Sedi- ment.	S.	ŵ	M.	ZZ	ΞÞ	1.S.		ń vi	M.	M.	vi;		i	
A.P.	Turb'y.	VS.	V.S.	:	VVS.							%	;;	.11.	
E OF	1899 Examination.	July 5	24 July 26		8 8 3 3 8 8 8 8			Oct. 4			Nor. 2	6	16	**	
DATE OF	1899 Collec- I	July 3	Ang. 1			Sept. 5	96	Oct. 3	: :	26	31	Nov.	: :	2	
No.	Serial	350 July 330 ::	344		331		245	25.0	1030	1032				oenr	

S., Slight. VS., Very Slight. VVS., Very, Very Slight. M., Medium. *No Sample.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-ILLINOIS RIVER, PEKIN, ILL.

Keport of Abolphi (FHRMANN)	D. B. BISBEE,	Municipal Laboratory, Chicago, 111.

No. of Bae.	Cubio	Centi- meter.		686.500	370.000	1.653,500	2,880,000	1,157.000	16.000	49,000	45,000	187,500	1,020,000	462,000		580,000	3,490,000	617,500	9,255,000	305,000	60,000		
ce or Colf.		Prese		+	+	- <u></u>	+	-1	-	+	+	-	+	-		+	+	+	+	+	1		
		dm9T'	30.	33	32.	34	34.	33	33.	40.	40.	#1.	27.	12	∞.	34.	555	21.	50	+	15.	16.	36
ature D. T.	19 910	qm9T sW lo	24.	86	26.	30.	55	21.	55.	.98	. 68	88	30.	33	<u>5</u>	<u>5</u>	150	<u></u>	16.	10.	9	10.	0
				-	e3	ī.	_	-	ıc.	iù			-	1.6			,			7.0	_	93	1.5
0. 7 AS	69	Mitrat		1.74	1.97	1.66	1.03	7.	1.50	1.39	64.	.54	5.7.0	.52	25.	1.00 000	92.	2.36	1.70	3.74	1.90	2.58	1.27
NITRO- GEN A	89	Mitrit		.36	٠ 0	₹?:	89.	18.	.30		.11	.16	.07	.085	.B.	88	19.	19.	1.13	.36	0×.	37	
IC EEN.		puəd -sns																					÷.
ORGANIC NITROGEN.	.b	siQ 9vfos		-									-						quint(8.
02	1 .1	B10T	_			_	_	_		_		_					_	-	_	_	1.46		
100	d Am	-sus p,pud	80.	96.	.16	.10	.30	.16	88.	. 26	35.	÷:	91.	30.	€.	.22.	.33	.33	38.	4.	.20	.16	.18
GEN AS	MMONI.		.52	14.	.48	.30	07.	04.	04.	#	.42	æ.	040	£.	4.	88.	08	8	5.54	01.	.36	£.	.59
NITRO	Ed Fluor		99.	57.	1 9.	9+.	90.	.56	84.	.70	8.	£.	.56	8.	.63	3.	.70	98.	1.01	∞.	.56	81.	.70
	FreeAm- monia.						(med												_		3.00	C/5	1.00
ED.	RySuspd B Natter Hysuspd		1.6	€3 ∞	e3 e3	7.	3.1	∞.	6.1	e3	<u>∞</u>	6.2	٠ :	4.	€.	65	٥;	3.6	<u>~</u>	€5 ∞	1.6	25 25	ej.
OXYGEN CONSUMED.	pi	sid val	Ξ	23	5	<u>∞</u>	o,	∞	œ	9	9	<u>م</u>	œ.	∞	∞	∞	œ	с: _	x	œ.	7.6	8.9	à
000	Total.								10.4			15.6									6.6	6	10.
Chlorine.		35.	36.5	40.5	31.	55	83	35.5	43.5	54.		55	33	51.5	55.	56.5		67.	93	99	59.5	50.5	
ON ION.	pa	sng	18.	35.	46.	स	22		38	<u>:</u>	<u>∞</u>	33	33	£ .		19.	21.	=	;;	33	જ	30.	21.
RESIDUE ON EVAPORATION	1	-slU oovlos	365.	391.	353.	200.	319.	315.	370.	37.5	366.	360.	365.	355	356.	37.5	337	300	10.	<u>:</u>	395.	370.	370.
RES		[n10T]	383.	416.	300	333	346.	36.50	3000	.381	38.	136.	33.63	127	373	391	358.	101	437.	427.	115.	390	391.
	Odor		None	9 9	9.9	•	•	;	. ,	9.9	9 9	3.9		:	9 9	9 9	9 9	9 9	*	9 9	:	;	**
		Color.	्रा.	??	-	-:	4	185	ಣ				:										·····
PEARANCE.	Sed! C ment.		ŝ	Š	ŝ	N.	Ý.	M.	M.	· · · · · · · · · · · · · · · · · · ·	Ŀ.	M.	.N.	. I.	ジン	M.	ŕ.	·j.		·j.	Ŀ.	s.	ý.
A V	Turb'3		VS.																				
DATE OF	1000	Exami- nation.	6 July 8																		ō .,		
DAT	1000	Collec- tion.	July 6		61 .,			o. ∶				Sept. 5)ct. 3					10V. T		
No.	1 0				137	500	236	133	大きい	48.60		.,				-					7. 1551		

M.. Medium. VVS., Very, Very Slight. S., Siight, VS., Very Slight. VVIII., Very, Very Heavy. VII., Very Heavy. APPEARANCE-II., Heavy.

TABLE 175.

Report of Adolph Gehrmann.
D. B. Bisbee.
Municipal Laboratory, Chicago, Ill. STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO. SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-ILLINOIS RIVER, HAVANA, ILL.

Parenteance Parenteance																								
Parameter Para	No. of Bac.	Cubio	Centi- meter.	11,500	19,065	74,500	23,650	3,000 1,000 1,000	21.950	9.250	34,500	30,333	67.000	350,000		35.500	145,000	052,19	17,000	33,750	11.000	845,000	:	106,000
Paper Pape				+	+	+	+	1	+	+	+	+		+		-}-	+	+	+		+	+		+
Park Park				कं	83	30.	88 88		300		31.	: :38:	33.	:		1.	<u>13</u>	21.	G	19.	∞	5.	=	13.
Section Control Cont				.92	જુ	27.	30.	:	36	33	36.	- 36 36	66			16.	14.5	15.	17.	12.	6	<u>}</u> -	10.	=======================================
Section				4.7	-:	6.4	∞.	:	4.1	က (၃	ಣೆ	e;	7:02	:	3.6	က	ಣ	9.8	دن زه	<u>က</u>	3.5	9.4	٠٠ ٠٠	4.4
September Parameter Para	-0.	รอ	Nitrat	• !	1.45	1.74	1.12	69	1.04	86.	02:	. 16	.17		٠ <u>٠</u>	67.	80.	99.	09.	.60	2.16	09.	1.98	
Pape of Papearance Papearan	NITH	sə	rinin	• 1	જ	86	. 28	:3	.165	. 33 26 27	3.6	56.	.036	•	 	=	57		09.	200	1.04	1.30	?}	<u>:</u>
PATE OF PAPEARANCE. PAPE	NIC GEN.		sns														_				•			-:
PATE OF PAPEARANCE. PAPE	ORGAN	-	sid					_									_							
Formula Part of Pa			p,pud				_		_		_	-		_			_	_		-	-			_
Boate of the control of the contro	K AS		p, Alos			_	_							_	_	_								
Boate of the control of the contro	ROGEL	(lbumi																						
Pate of the color Date of the color Date of the color 1899 Lurb's Sedi- Color	TIN															_								
Pate of the color Date										9.	6.	7	ा	00		33	<u>}</u>	7	7	33	-	00	_	
Pate of Pate of Pate P	GEN	pd	BySus	es es	6 1	0 8	00	-	23	<u>ල</u>	55	- 9	6 2	ಞ	To To	<u></u>	4	\$5		9	_	₹ 1	03	9
Pate of Pate of Pate P	OXY	Total Cox		63	C)	9	7	∞	-	50	4		00	<u>@</u>	7	03	9.	9	7	∞			03	9
B99		1				٠0	10	10		10	_		10		10	_		0			321	10		
1899 1899	.9r	rir	СРГО	ನ	ਨ 	3,	33	ಸ _	 ਲ	<u></u>		7	7	7	<u>ښ</u>	<u>بر</u>	<u></u>	- 6.	<u>ෂ</u>	⊛ 	9	<u>,,,</u>	70	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~
1899 1899	ON MON.	pa	sng -sng	14.	333		31.	भ	27.	100	70.	330	53	33	107	7.0	.99	63	, 35	35	£	6	16.	- FG
1899 1899	SIDUE	1	SOIVEC SOIVEC	387.	374.	379.	317.	322	330.	347.	349.	384.	345.	343.	303.	350.	359.	397.	348.	391.	393.	395.	369.	308
PATE OF APPEARANCE. 1899	RE		Total	401.	107	384.	348.	364.	384	424.	419.	493.	398.	435.	410.	429.	₹.	160	423.	446.	435.	404	385	333.
1899 1899		Odor		None	3.9	"	3	:	3	13	18 e	18	9.9	`. •	9.9	4 .	:	:	"	•	• ,	*	9.4	7 3
1899 1899 1899 1899 1899 1899 1899 1800.			Color.	e;	-	-:	19	ಣ	ં						•					•				
1899 1899 1899 1899 1899 1899 1899 1800.	PEARANCE.			M.	M.	VVS.	Š	M.	M.	M.	M.	Š	M.	M.	M.	ś	M.	ś	M.	M.	Š.	ś	ý.	જ
B99 1899 1899 Collec- Example Link 12 Collec- Example Link 13 Collec- Example Link 13 Collec- Example Link 13 Collec- Example Link Lin	Turb'3		M.	VVS.	VVS.	VS.	M.	M.	M.	M.	M.	M.	M.									S.	si.	
DAT 1899 Collection. July 5 1999 Aug. 28 Aug. 28 28 11 12 13 13 15 18 18 18 18 18 18 18 18 18 18 18 18 18	OF	1000	Exami- nation.	fuly 6	" 13	% 30	27			17 1,	1,54	** 31	ئد	14	" 31	86.)ct. 5	12	,, 19				16	
	DATE		uly 50	12	19	97	lug. 2	6	16	" 23	. 30	ept. 6	13	02 ,,	16	let. 4	** 11	.:	. 25	iov. 1	00	151	÷;	
	No.	1 0				959	260	7.9		263	264				268	692	\sim		37.5	273	274	27.5	101	

M., Medium. VVS., Very, Very Slight. VS., Very Slight. s., Slight. VVII., Very, Very Heavy. VII.. Very Heavy. APPEARANCE—H.. Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-SANGAMON RIVER, CHANDLERVILLE, ILL.

OOLPH GEHRMANN,	B. BISBEE.	original Laboratory (thicago
ADOLPH.	D. 13.	Municipal
Report of		
p-tre(

vo. of Bac.	per 'nbio	Centi- meter.	16.500	24,000	10.500	44.350	34,750	11,750	5,465	62,000	38.38	7,750	14,250	13,250	18.350	11,750		13,750
1100	JC	Preserved Abs. o	+	- 1	+	- +	- (-		+		+			+	1		
		dinaT A lo	28.	300	33.	.68	65	27.		. 18	16.		55	5	x	Ξ	16.	<u>-</u>
		qm9T 37/ 10	27.	57	.96	33	.36.	25	37	19.	{-	6.	15	18.5	10.	10	12	
		11911 177			25	3.6	rċ	:	Nor	3 3	;	9.9	* *	9 7	- • •			25
o-o	89	Mitrat		2.06	338	16:	86:	۰				•					99	· ·
NITRO- GEN A	89	Mirrit		.036	.023	98	.0.S	900.		800	11.	.004	.005		.008		015	0.
IC EN.		puəd -sng		56.	.18	.26	.33	:	.36	.36	81.	* 1.	80	200	.56	.36	37	=
ORGANIC NITROGEN.	d.	SIU						٠										04.
ôZ	.1	T.ota	.60	E	.69	.76	.52	:	93.	.61	1.26	00	555	.56	1.7	.58	99.	10.
00	d Am	p,pud -sns	. 18	61.	. 13	08.	.17	01.	91.	. 16	15.	80.	3.	5.	60	10.	.06	.07
NITROGEN AS AMMONIA.	Albuminoid	Dis-	.14		-	-	91.	.50	91.	. 16	#	02.	Ξ.	**	=	1.	.30	.15
NITRO	Albi	T'toT	33	.36	.30	.37	. 33	99.	33	.32	89.	89	. 53	92.	0	<u>∞</u>	.36	65.
	FreeAm- monia.		.02	80.	80	14	10.		3	.016	8.	30.		.016		.018	10	27
SN IED.	Bysuspd		4.9	ಬ			20			2.9	63. 50	<u>∞</u> .	4.03	2.1	00	{-		- 31
OXYGEN CONSUMED.	By Dis- solved.		33	÷	20	ಣ	3.6	ь.	er;		$\dot{\infty}$	35	φį		e3	200	35	22
000	. {	nioT	∞ **											8.4				r0 60
.91	111	очо	5.1				30							5.6				
ON ION.	bended S. S. S. S. S. S. S. S. S. S. S. S. S.		157.	136.	64.	1666	33.	332.	93.	95.	79.	30.	.03	45.	.39	26.	33	5.
RESIDUE ON EVAPORATION	T	std oavlos	280.	309	308	255	:Xei:	319.	989	:277:	350.	200.		.X.0X.	394.	201	255	27.0
RESEVA	1	вюТ		- FOE	373	431.	388	651	250	373.	1:30	330	351.	353.	333	317	310.	330.
	Odor		None	• •	:	* 9	:	:	;	;	:	:	*	:	:	* 9	;	:
		Color.									:			:		:	:	:
APPEARANCE	Sedi. C		Ξ.	policy backet	. N.	=	11.	.N.	N.	M.	Ť.	ý.	j.	ý.	ý.	j.	ĿŚ.	M.
17:	Turb's		H.	7.7.8.	N.	M.	.N.	.M.	N.	==	Ξ.		j.	N.	ý.	j	ij.	N.
ATE OF	DATE OF 1899 IRON. Ination.		5 July 6	12 : [3	3.7119. 3	01 . 6	16 17	31 Sept. 1	:	50 · 66	:	ct.	:	35 : 35	01.	:	15 . 16	: 3
	No. DATE 1899 Collect Lion.		To July	:	Aug.			:	ニニナ	:	1.	Oct.	:	:	10%	:		:

M.. Medium. VVS., Very, Very Slight. VS., Very Slight. S. Slight. VVIII., Very, Very Heavy, VII.. Very Heavy. APPEARANCE-II. Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-ILLINOIS RIVER, BEARDSTOWN, ILL.

Report of Adolph Gehrmann, D. B. Bisber. Municipal Laboratory, Chicago, III.

No. of	per Cubic Centi- meter.	30,350 3,350
	Presenc	+1++1+1+1++++++++++++++++++++++++++++++
	rempera riv. ro	<u> </u>
ture . C.	Tempera	######################################
	tdgleH i91877	F-0 0 0 0 F-0 0 0 0 0 0 0 0 0 0 0 0 0 0
NITRO- GEN AS	Setrativ	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
EN E	Nitrites	
SIC GEN.	Pended Sus-	000000000000000000000000000000000000000
ORGANIC NITROGEN.	-sid .baylos	888.2.2.38888. :25.3 :3.4.88.3.25.6 :86 :
	Total.	- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1
m	P -sns	5683847591785799
NITROGEN AS AMMONIA.	bionimin b'vios	<u> </u>
NITRO KICL	I'10'I'	<u> </u>
	FreeAm- monia.	848388884548484848
ED.	BySuspd	### 0
OXYGEN CONSUMED.	By Dis- solved.	
000	Total.	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$
.9I	Chlorin	72 28 88 8 8 2 8 8 8 2 8 8 2 4 4 4 4 7 7 7 2 4 5 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
ON TON.	papuad -sns	<u> </u>
RESIDUE ON EVAPORATION	Dis- solved.	82.8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
RE	Total.	338 339 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
	Odor	None
	Color.	
APPEARANCE	Sedi- ment.	
IV.	Turb'y.	HANNER REPORT OF THE PROPERTY OF STANFORD
DATE OF	1899 Exami- nation.	July 8 28 28 Sept. 18 18 28 Oct. 6 28 Nov. 3 17 Dec. 11
DAT	1899 Collection.	July Aug. 280 1
No.	Serial	34 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4

M. Medium. VVS. Very, Very Slight. VS., Very Slight. S., Slight. VVII.. Very, Very Heavy. APPEARANCE—H., Heavy. VH., Very Heavy

TABLE 178.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO,

SOURCE OF WATER-ILLINOIS RIVER, KAMPSVILLE, ILL.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

No. of Bac.	per Cubic Centi: meter.	2,400 6,150 11,500 12,500 10,5
	Presend lo .sd A	111+4+1++++++++++++++++++++++++++++++++
	Tempers of Air,	
	Teniper 9187/10	हिन्न हिन्न हो हो है है है है है है है है है है है है है
	1dg[91] 91877	第
NITRO- GEN AS	sətmitiN	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
N.r.	Satistics	
IC REN.	Sus-	58 58584 SET85 S
ORGANIC NITROGEN.	Dis-	<u>\$5 84858 8568868</u>
ōΩ ——	Total.	85 588325 56 88285 55
200	B P.pud	F3362289
NITROGEN AS	Albuminoid -sld -sld	स्ट अध्यक्षिक्ष स्टब्स
NITEO	[1'10T	<u> </u>
	FreeAm- monia.	<u> </u>
EN IED.	Bysuspd	6 1 1 : 24 6 1 1 1 1 2 1 1 1 1 2 1 1 1 1 1 1 1 1 1
OXYGEN	By Dis-	FF
ဝဌ	Total.	
.91	Ublorli	57 54 88 88 54 48 10 </td
ON FION.	papuad -sus	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
RESIDUE ON EVAPORATION	-sid -bevios	
Ev.	lsto'l'	30.4 30.4 30.4 30.4 30.4 30.4 30.4 30.4
	Odor	None None None
	Color.	£ - : - : - : - : - : - : - : - : - : -
APPEAUANCE.	Sedi. ment.	
Turb'y.		N N N N N N N N N N N N N N N N N N N
DATE OF	1899 1899 Collect Exami- tion. nation.	12 July 13 16 July 13 16 July 13 16 July 13 16 Sept. 7 17 17 17 17 17 17 17 17 17 17 17 17 1
DAT	1899 Collec- tion.	E Signature of the state of the
.06	Serial	द्वश्चर्यश्चर्यः सहहहहहरूके

M.. Medium. VVS. Very, Very Slight. VS., Very Slight. S., Slight. VII., Very Heavy. VVIII., Very, Very Heavy. APPEARANCE-II. Heavy.

TABLE 179.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ILLINOIS RIVER, GRAFTON, ILL.

Report of Adolph Gehrmann, D. B. Bisber, Municipal Laboratory, Chicago, Ill.

•																							
No. of Bac.	per	Cubic Centi- meter.	11,350	13,700	20,500	15,350	32,750	62,500	25,000	11,750	19,250	44,000	23,000	46,650	17,350	29,000	17,750	47,500	38,000	5.000		40.250	17,500
		Prese Abs. c		+	- 1	1		+	+	+	+	-	+	+	-	+	.	+	+	+		+	+
		qm9T A lo	.38.	%	36.	37.	37.	.58.	930	31.	31.	34.	25.	19.	30.	.02		.54.	10.	17.	14.	1.	E
ature r, C.	- 818	Temp W 10	.68	53		34.	33	25.	.98	31.	30.	33.	33	30.	19.	16.	17.	19.	20.	6	10.	13.	8
		giəH 3VI	_	13.2									4.4	4.8	4.1		63	63	33.4	5.4	6.2	6.	5.9
NITRO- GEN AS	sə	Nitrati	:	1.18		_	0.5	12.5	_				:	.75	:			.57		_	_:	2.43	
NITR	68	Nitrito		.13		.12						•		.057	۰							8	•
GEN.	_	-sng	<u> </u>	.18	•	:															_		91.
ORGANIC NITROGEN		siU solve	09:			-																_	99:
OZ	1	Total	97.	•	•	:															35	_	.72
82	id Am	p,pud -sng		_	_		_			_		_		_				_	_				.30
NITROGEN AS	Albuminoid	-siU b'vlos	 																				
NITEC	Alb	L'10T	.43	œ.	#.	.35	.38	88.	.37	. 335	.31	.41	 88	.37		56.	.34	.33	.36	.38	88.	.72	.72
		гееля inom	.02	.15	. 12	.13	30.	.12	80.	.0 .0	.05	.18	8.	:S	.40		દુકું	. 16	.30	.56	.15	80.	96
EN IED.	. . .		9.	+.	0.1	-	:	3	2.4	00.	-	_	1.1	_	-:	-	щ.	1.6			6.	5.3	1.7
OXYGEN		By Dis		-3	7.4	7.5	-	5.5	5. 5. S.	9.2	6.8	6.7	6.4	6.3	6.2		6.	6.				5.7	
03	Total. 80		8.4	7.5	7. 7.	7.6	2	∞ 3.	9.6	8.4	∞ ∞	7.9		7.00	5.5	6.7	6.3	7.6	8.6	9.9	12.8	11.	<u>∞</u>
Chlorine.		10.5	10.4	1.4.0	24.5	33. 53.	15.5	ж то	17.5	15.	21.	21.	24.5	33 12	30.	31.	56.	33.	39.	43.	31.5	27.5	
ON ION.	pa	epuəd sng	20	7	16.	18.	22.	178.	117.	33.	33.	- - - - -	27.	65	49.	£6.	30.	- 1 2	94.	27.	48.	8	43.
RESIDUE ON EVAPORATION	J.	Dis	306.	300	313.	329.	345.	303.	183.	276.	275	318	308.	301.	256	234	316.	.962	258	315.	303.	301.	301.
EVA	1.	R10T	313.	307	329.	347.	367.	381	300	311.	310.	336.	336.	356.	305.	280.	346.	341.	352.	342.	350.	386.	314.
	Odor		None	-d	9 9	7.7	9.9	3	9.9	9.9	9 9	3 9	9.9	9 9	9 9	• 9	9.9	3	9 9	9.9	9.9	9 1	7.7
ម៉		Color.	.05	6		:S:	ಬ	33.	:	:	:	:	:	:	:		:	:	:	:	:		
APPEARANCE		Sedi- ment.	VVS.	VVS.	VVS.	Z.S.	7.8.	N.	N.	M.	L'S.	S	7.8.	ś	VVS.	L'S.	VVS.	VS.	si,	ZS.	ZS.	VS.	Š
AP	Turb's		VS.	VVS.	N.	VVS.	M.	VII.	V.H.	M.	H.	H.	M.	H.	-	1.11.	M.	M.		M.	VII.	II.	MH.
DATE OF	1800	Exami- nation.	5 July 7	9 :	? ?	9.0	\use	,	9.9	7 7	9 -	sept.	9.9	3 9	3 3	ot.	7.0	,,	100.	;	9 9	9.9	ec.
DAT	1800	Collec- tion.	July	9 :	9 9	99	Aug.	:	91 ,,	31	08 ;	Sept. 6	13	06 ,,	767	Oct. 4	; 11	: 33	Nov. 1	œ :	15	€6 3.	9.9
.o.V	g	ires	514	516	218	521	555	524	551	553	555	557	559	561	563	565	567	569	571	573	576	578	580

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VVH., Very, Very Heavy. VH., Very Heavy. APPEARANCE—H., Heavy.

TABLE 150.

Serial No.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO SANITARY WATER ANALYSIS-PARTS PER MILLION

SOURCE OF WATER-MISSISSIPPI RIVER, GRAFTON, ILL

D. B. BISBEE. Municipal Laboratory, Chicago, III

Report of Abouph Gehrmann,

28,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 11,500 9,000 per Cubic Centi-meter. Presence or Abs. of Coll. Temperature of Air, C. Temperature of Water, C. 8638835588888888 13.4.6 13.2.7.9 17.6.7.9 17.6.6 17.6.6 ा चे दिस्स च च व्य おこれであるないたれずらさいたますら Height of Water. = 8885866 65. 0000 GEN AS SHIERIES NITRO-.00. 90 8 Nitrites pəpuəd -sng ORGANIC NITROGEN. 2 4 2 5 *** solved -SI(I Total p,pud -sng Albuminoid Am NITROGEN AS solv'd b'vlos AMMONIA 0.4%%%41044%%7446% Lon 016 22828 82222238 :18 छ म सक् -201010 Matter BySuspd OXYGEN CONSUMED. By Dis-Total Chlorine, pəpuəd -sng **医黎思士氏学型好好的尼士器单行生物的**的最大 RESIDUE ON EVAPORATION. -sid sofved 8282346888883383538538 Total, Odor Color 1- 69 00 00 00 01 APPEARANCE. Sedi-ment. <u>自己对对对自然对对对对对对反应反对对对反应类</u> Turb'y. Collec- Exami-nation 515 July 5 July 6 July DATE OF 1899 tion.

Medium VVS., Very, Very Slight. VS., Very Slight. Silght. · ſ. VVII., Very, Very Heavy. VII., Very Heavy. APPEARANCE - II., Heavy.

TABLE 181.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM ILLINOIS SHORE, ALTON, ILL.

Report of Adolph Genrmann, D. B. Bisbee, Municipal Laboratory, Chicago, 111.

	No. of Bac.	per Cubic Centi- meter.	30,000 37,330 38,200 17,630 8,500 8,500 8,500 8,500 8,500 17,630 17,630 17,630 18,730 17,630 17,
		Presenc	++++11+++++++++++++++++++++++++++++++++
		Tempers	****************
	eture r, C.	Tempers	***************************************
		Height Stell	ででいいた+ここの + ここの 1 1 1 1 1 1 1 1 1
	NITRO- GEN AS	Nitrates	
		Nitrites	4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	NIC GEN.	Sus-	4888888888888848884888888
	ORGANIC NITROGEN.	-sill solved.	00 +0 00 00 00 00 00 00 00 00 00 00 00
	02	Total.	0.0.4.0.8.0.8.0.0.8.0.0.8.0.0.4.8.0.0.0.0
	AS	a sus	
	NITROGEN A	Dioniminoid Pis-sid	<u> </u>
	NITRO		<u> </u>
		FreeAm- monia.	8.8.8.9.9.9.8
	EN IED.	BySuspd	
	OXYGEN CONSUMED.	By Dis-	#####################################
	၁၀	Total.	1000000000000000000000000000000000000
ı	.en	Chlorin	
	on rion.	Sus- Sus-	444498885888888888888888888888888888888
	RESIDUE ON EVAPORATION	Dis- solved.	200 200 200 200 200 200 200 200 200 200
	Ev.	Total.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
		Odor	None
	r i	Color.	- 19 19 19 19 19 19 19 19 19 19 19 19 19
	APPEARANCE. Sedi- ment.		
	Turb'3		
	EOF	1899 1899 Collec-Examition.	Aug
	DATE OF	1899 Collection.	July 5 199 199 199 199 199 199 199 199 199 1
-	No.	Serial	: 2015 25 25 25 25 25 25 25 25 25 25 25 25 25

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VVII.. Very, Very Heavy. VII.. Very Heavy. APPEARANCE—H., Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM ILLINOIS SHORE, ALTON, ILL.

Report of Adolph Gehrmann.

B. B. Bisber,
Mudicipal Laboratory, Chicago, III.

No. of Isac.	per Cubic Centi- meter.	26.350	8.00.8	15,250	7,750	24.635	19,666	56.650	44,000	36.000	29.500	36,750	15.000	23,000	17.350	11,000	17,000	6,750		21.000
	Present to .sdv	++	+	+		1	-	!	1	+	+		;	+	+	+	+	+		+
	Temper	1000	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	34.	33	× 5	\$ £	35	65 70	?}	20.	19.	16.		19.	19.	10.	12.	2	7
	Temper of Wate	36.	38	⊗	양	26.	8 8	38	30.	24.	્ટ સ	<u>∞</u>	<u></u>	16.	- 20	<u>∞</u> :	27	6:	9	<u></u>
10 1	Helgh Wate	10 m			1-1 25							တ္								
RO.	Vitrates						2 ,0													
NITRO	sethitiv		.003	.013	왕() ()	(G).	000	.012	<u></u>		.0	0.	10.	.016	.008		.015	.00.	0.0	9
c EN.	bended sus-	£8.	: 53	·	₹.	56.	2 1	38	01.	.46	1.0.		3	.16	16.	.30	.3.	1.	10	238
ORGANIC NITROGEN.	Dls-	92	16	TT.	37	04.	8 8	36	.31	+ :	.36	=	 00	∞ ∵	.40	<u>a</u>	55.	Ŧ.	20	.68
ON	Total.	8.	7.6	33	<u></u>	8. 8	3 33	1.5	-	8:	8:	X.	1.01	5	16:	∞ - 1 - 1	.66	∞ ∞	16	8:
	a p.pud	70.	0	<u>31</u> .	. 23	10.1		200	10	÷:		£	91.	5.	<u>ं</u>	=	:	=	96	8
NITROGEN AS	Albuminoid -sid b'vlos	.33	8	87	5.3	89.8	3 5	£	हें: हें:	. 23	. 27	÷	÷;	हों:	25.	₹?	33	200	200	8
NITRO	Tot'l	.40	.40	.40	.57	<u>.</u>	÷.%	.36	.30	77.	=	94.	7	.39	<u>∞</u> .	300	.32	.39	10	<u>10</u>
	FreeAm-	ਰ:	.03	.03	S.	8.5	5	.03		:	:	910.	89.	3	0.	820.	1 0.	80.	0.18	30.
SN (ED.	BySuspd	63					÷ 22										300	33	_	
OXYGEN	By Dis- solved.	14.6					6.9									œ				12.6
000	Total.	17.6	15.2				10.0										10.8			
,91	Срјоги	1.9	3	0. 0.	6.4	का द व्य	5 .c 2 .c	∞. ∞.	5.1	5.5	ص اد	, ic.	5.	6.5	တ တ	တ တ		∞ ∞		
ON TON.	popuod -sus	88	00.	33	= 7	(); (); (); (); (); (); (); (); (); ();	73.	53.	65		00	33	3	0.	76.		3. 3.	989	71.	99
RESIDUE ON EVAPORATION	-sld beyled.	160.	130	190	191.	200	18.5	213.	197.	133	30.5	159.	170	136	197.	£	17.	130	39	176.
RE	T'otal.	258.	268	305	238	1533	10 S	266.	262.	278	.290	300	25.50	356	25.5	257	569	250 250 250 250 250 250 250 250 250 250	- T-55	213.
	Odor	None	None		1 :	7 9	:	9	9	10 I		; ;	-		9 1	10 A	-	9 9	9.1	9,
	Color.	∞.	<u></u>	ಛ	si j	<u>e</u> .			:		:	:								
APPEARANCE.	Sedi- ment.	H.	H.	Ξ	<u>_</u> ;	<u>:</u> :	: ×:	N.	7.	i:	Z:		<u>.</u> :	į:	Z.	. N.	7.	.N.	Ť.	j.
11.	Turb'y.	VII.					- X	ý.	7.		7.7		M.	Y.	×		.N.	M.	.1.	.M.
OF	1899 Examination.	20.	÷	:	YIIK.	: :	: 55	:	ept.	= 2	. :		JCI. 5				10V. 2		91 16	666
DATE OF	1899 Collee- F		. 19	3		क य	2 83 :	30	Sept. 6.S	200	हु ! : :	**	٠'	_ ;	× 1	S.	NOV. 1.NON	00	15	€ 1:
.oZ	Serial	7.05	715	17.0		1000	012	715	155	135	33-								1(55	1059

M., Medium. VVS. Very, Very Slight. VS., Very Slight. S., Slight. VVIII, Very, Very Heavy. VII., Very Heavy. APPEARANCE-H., Heavy.

TABLE 183.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, ALTON, ILL.

Report of Adolph Gehrmann, D. B. Bisber. Municipal Laboratory, Chicago, Ill.

No. of Bac.	per Cubic Centi: meter.	34,750 19,000 65,500 6,250 83,000 19,666 12,000 55,000 11,750 14,750
	Presenc	+ + + + + + + + + + + + + +
	risqual aiv to	# 8 # # 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	request estail to	数数数数数数数数数数数数±20000000000000000000000000
	HelgləH etaV/	ででいるところのでもというとことでも にあら 。
NITRO- GEN AS	Zitrates	88894804080000
NIN	Nitrites	90000000000000000000000000000000000000
IC JEN.	pəpuəd -sns	86488866884866586486
ORGANIC NITROGEN.	Dis-	変字式 35 4 8 8 8 8 8 8 5 5 7 5 4 4 6 5 5 5 8 8 8 8 8 5 5 7 5 6 5 5 6 5 5 5 6 5 6 5 6 5 6 5 6
oz —	Total.	86.5.5.5.5.5.5.8.8.8.8.8.8.8.8.4.8.8.5.5.5.8.8.8.
	B p.pud	
GEN AS	Pionimula I Torr	 <u>ន</u> ន់ដន់នង់នង់ដន់ដន់ដង់ដង់នង់ដង់ដង់នៅន
NITROGEN A	I'10'T	<u> ब्रह्म कुर्य के चर्च क्ष्म के क्ष्म के क</u> ्ष्म के के क्ष्म के कि क्ष्म के कि क्ष्म के कि क्ष्म के कि कि कि कि
	FreeAm- monia.	2000 00 00 00 00 00 00 00 00 00 00 00 00
ED.	Bysuspd Matter.	# @ 16 4 61 60 60 61 61 60 60 61 61 61 60 4
OXYGEN CONSUMED.	By Dis-	4 x x x y y y y y y y y y y y y y y y y
CON	Total.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
.9n	Chlorin	
ON.	pepued -snS	28.88.88.88.88.88.88.88.88.88.88.88.88.8
RESIDUE ON EVAPORATION	Dis-	25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
RES EVAI	T'otal.	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
	Odor	None
	Color.	∞', y 21 ω ω τ ; G
APPEARANCE	Sedi- ment.	
AP	Turb'y.	
S OF	1899 Examination.	12 July 19 19 19 19 19 19 19 19 19 19 19 19 19
DATE OF	1899 Collection.	July 5 1919 1928 Aug. 26 1939 1939 1940 1950 1
Serial No.		:: 8 2 3 3 3 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VVII., Very, Very Heavy. VII., Very Heavy. APPEARANCE-H., Heavy

TABLE 184.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, ONE-FOURTH DISTANCE FROM MISSOURI SHORE. ALTON, ILL.

Report of Abolph Gehrmann.
D. B. Bisber.
Municipal Laboratory, Chicago, 111.

No. of Bac.	per	Centi	19.00	35.500	33.50	24.000	3.250	28.350	15.333	11.000	35.500	20.000	10.500	18,500	10,000	14.500	19,500	17,500	0000 00000	130	10,500		19,500
	Presence or Abs. of Coll.			- 1	1	+	1				£			+	. 1.	+			+	_1	_ 1-		
		Temp T	27.	₹. ₹.	34.	34.	333	æ.	30.	£.	3	.055	??	(3)	19	16.	1-	÷.	19.	5	9	~	=======================================
		gmoT'	.36.	99	\$ 50 \$ 50 \$ 50 \$ 50 \$ 50 \$ 50 \$ 50 \$ 50	30.	55	36.	:36:	× 1	35	30.	.33	.05	-8	Ib.	16.	2	<u>x</u>	21	c.	10.	<u> </u>
10 1	010 Ц2		15.		5	6.			{-	7	20	23	2.2	7	2.2	25	ુ:	21	_	71	77	ಾಣ	
NITRO- GEN AS	89	Jariin		_		08. 6											0.					000	8
1. S		Minit				.005									_								0.
HC DEN.	_	sng -sng																					. 10
ORGANIC NITROGEN.		elU evios				.36																	
ON.	1.1	T'ota	<u> </u>	-											_							_	39.
	d Am	sus-	15.	99.	FG:	Ġ1.	+1.	II.	154	91.	. E.	. 11	??	(0)	+?·	. 19	.11	.21	<u></u>	27	15	80	.13
NITROGEN AS AMMONIA.	Albuminoid	-sitt b'vios	88	\$. 80	36	.36	36.	??	÷;	\$? ·	5	≎?	<u>و</u> د.	? ?	??	19	97.	7.	왕	??	5	35	35
NITEO	Albu	[,10,[,	55.	æ.	<u>S</u> :	=	OF:	:8: :8:	.384	0+.	.32		<u> </u>	. ‡6	<u>∞</u>	7	.35	<u>::</u>	330	98:-	04.	9	=
		Угеел. Нош			·	.01		·			:		:	:	:	:	:	:	:	·	:	•	50.
EN IED.		Byshus	30	×. ±		:	35	ಯ	÷	-	જો	ાં	<u> </u>	30	33	~·:	33	91	??				
ONYGEN CONSUMED.		By Dis		-	_	∞ ∞ ∞	\dot{x}	{=	1-	1-	-	6.	φ.	1-	œ.		c:		œ.			-	
000	.1	вюТ	30.	25.6		:	= ::	=	11.6	Ξ	9.1	0.00	10.6	11 6	13.6			T: =					
.en	111	Chlo				₹. 67																	
0.N 10.N.	pa 	puad sus	210.	162.	159.	150.	÷.	108	164.	-52-	59.	50.	101	=======================================	3; 3;	-36.	76.	33		€	G	101	: :3
RESIDUE ON EVAPORATION		sld earlos	154.	164	2	170.	<u>8</u>	178	137.	<u>8</u>	<u>3</u> 6.	<u>&</u>	163.	162	145.	136.	162.	177.	- 689 - 689	162	163.	139.	139.
Ri- Ev.	Total.		361.	626.	3339.	3530	3	286.	301.	0:27	200	2 1	263	15	133	33	33x	269.	310	25	-	233	303.
1	Odor		None	0 10		:	:	:	•	•	-a -a			-a-	•	:	:		*	:	:	:	:
53		Color.		?}	-	er;	eć.	<u></u>		:	:	:		:	:	:	:	:	:	:		:	:
A PPEABANCE.		sedi- ment.	11.	7.11.	=	=	7.	7.	11.		N.	7.	N.	N.	7.	. N.	Μ.	N.	. N.	N.	7	Ť.	j.
d V.		Turb'y.	VII.		V.11.	II.	-	. N.	V.11.	N.	s.	.v.		Z.	7.	.N.	M.	N.	M.	M.	.1.	.N.	M.
DATE OF	1899	collec- Exami- tion, nation,	Tuly 5 July 7	:	:	36	\u2. 2.\u2. 3	: G	. 16 . 17	3	30		:	: 0?	:	# Ort.	;	; <u>«</u>	33	. 1 Nov.	i àc		: 66
Zo.	[B]	204	707 Ju		-1-		-14		1:31							TIS OFF				70% SSL		1654	1.657

M., Medium. VVS., Very. Very Slight. VS., Very Slight. S., Slight. VVIII. Very, Very Heavy. VII., Very Heavy. APPEARANCE-II., Heavy.

TABLE 185.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Augleh Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 FEET FROM MISSOURI SHORE, ALTON, ILL.

No. of Bac. Per Cubic Centi- meter.		20.500	18,000	62,000	33,000	19,000	13.000	16.000	16,000	32,600	23.500	50.250	12.350	13.350	11,000	16.000	8.350	19.500	9.000		23.000		
Abs. of Coli.			- -	- 1	ſ	1	1	- 1	1									1		:		:	
10 90	uə	Pres		_	_	_									_				_			+	-:
		qm9T A 10	37.7	85	31	34	33	80	33	80	81	22	8	19	91	-1	ର	19	10	10	13	=	:
atnre er, C.	19 316	qm9T 3V/ 10	26.	230	88	30.	36	36.	88	86	30.	24.	30	<u>8</u>	15.	16.	18.	18.	13.	6	10.	13.	
Height of Water.			15.3	15.8	12.2	6	رة دون	· ·	4.3	3.1	ი: გ:	3.6	477	30.00	હર જ	23 23	2.1	1.9	æ. ≈	3.9	5.4	4.7	
30- N AS	ธอ	Mitrat		8.	1.60	.70	.40	.30	٥.	0.	.16	<u>ين</u>	. 10	0.	:	0.		0.	0.	0.	.10	98.	:
NITRO-GEN A	Nitrites			900.	.003	:	:	.003		:	.00.		.007	.005		800.	900.		10.		:	0.	:
IC PEN.		puad	40	83	98.	.30	04.	.40	.42	.30	91.	***	.36	₹8.	.48	.44	.26	08.	.38	.18	.13	.10	:
Organic Nitrogen.		Dis-		.54														_					:
62 	1	Tota	6.	1.36	8.	87.	.78	8.	89.	.68	.50	. 100	∞.	£.	æ. —	<u>.</u>	.74	1.	89. —	- 68	99.	89.	:
16	i Am	p,pud -sng	.31	-18	.14	.12	.31	.16	91.	Ξ	<u>::</u>	.16	91.	02:	.16	. 12	€.	.16	01.	.13	80.	. 13	:
SEN AS	Albuminoid	Dis-	.21	.28	∞ ?:	.32	£.	.24	.24	.24	음 :	.20	8	.24	98.	.25	33.	.24	97.	કે.	660	83.	
NITROGEN AS AMMONIA.	Albu	['10']	55	92.	.43	.44	.55	.40	.40	.35	.34	.36	.41	##.	.43	.37	110	0+.	.36	386	.40	#:	:
		FreeAi inom	50.	80.	30.	31.	10.	.04		:	:	:	:	:	:	:	:	:	:	8	8	30.	:
Bysuspd B x			₩. ₩.	13.	6.	-		3.7		ું	20.30	30	30 63	3.5			2.1		ن	25.	C)	.8	:
OXYGEN CONSUMED.	By Dis-		13.2	11.	10.	8.6			(- (-					35									•
000	-	Total	21.6	34	16.		10.6				0.5		11.						10.8				:
,91	ij	СРЈО		2.0																			
ON.	pa	sng	266.	407.	131,	121.	.09	149.	62.	.09	÷	33	80.	64.	56.	77.	.99	82.	.08	39.	76.	73.	:
RESIDUE ON EVAPORATION		Solved	141.	161.	180.	169.	167.	163.	166.	189.	187.	157.	167.	155.	157.	162.	167.	157.	157.	144.	143.	139.	:
RESEVA	.1	fs10T	407.	568.	301.	290.	227	312.	338	249.	235	240.	247.	219.	213.	239.	233.	239.	237.	183	.618	202.	:
	Odor		None	*	9.	***	;	. 9	;	:	:	:	:	;	:	9	:	;	.,	:	;	:	
		or.	-	ા	_	65	60	15		:	:				:	:		:	:	:			•
5.5		Color.)	***	•	4.9		•	:	:	:	:	:	:	:	:	:	:	:	:	:	:	
APPEARANCE		Sedi- ment.	H.	V.H.	H.	11.	M.	M.	M.	M.	M.	M.	M.	M.	M.	N.	M.	M.	M.	M.	ŝ	Š	
AF		Turb'y.	V.H.	VIII	VIII.	H.	MH.	M.	M.	Š	M.	M.	M.	M.	M.	N.	M.	M.	M.	M.	M.	M.	
)F	1000	Exami- nation.	١.	., 13			. •																:
DATE OF		c- Ex	5 Ju	.; ?;	. 61	96	2.11	6	33	30	6 Se	13	33	37	400	111	38	· ·	1 20	00	15		:
7C	1000	Collec- tion.		;		9 9		77		99	Sept.				Oct.		:	;	Nov.			3	
.o.Z	[13	Seria	7.08	713	718	7.53	7.28	733	743	7.18	749	758	763	768	773	178	783	200	7.93	298	055	990	:

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VVIII. Very, Very Heavy. VII., Very Heavy. APPEARANCE II. Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

Report of Adolph Gehrmann, D. B. Bisber, S. Municipal Laboratory, Chicago, Ill. SOURCE OF WATER-MISSISSIPPI RIVER, 400 YARDS FROM ILLINOIS SHORE, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

No. of Bac.	per Cubic Centi- meter.	21.350	19,000	9,350	7,750	Orange of	17.700	17.000		47,350	28,780 081,880	16.000	6,250	60,000		13,650	8,500	19,000	100 750	14,250
	Presence or Coll.		- 1	1	+-	-	+	- 1	:	+		n-q-	÷	+	. :	1	-+	- 1	+	
	requeT riv to	15	55	36.	8 8	. 6.	222	35	350	24.	20.	<u>x</u>	हर		<u>∞</u>	6:	oş	-		17.
	meqmeT etaV/ lo	6	36.	.56	8:3	970	101	27.	29.	33	.61	.91	15.	17.	17.		6.	6	Ξ	œ
	Helgh Wate	633	9.08	16.6	11 E	13.9	5.0	10.	9.9	9.9	9.9	5.3		دن ن	33	20	3.9	6.5	6.5	7. 76
30- N AS	Nitrates				04.0				:	:		:	:		٠	:				
NITRO	Solititiv	.003		.00 .	0.	200	600	100	:		8	950.		•			.013	<u></u>	10.	.019
IC GEN.	bended sus-	1.36	-	-	_	_														
ORGANIC NITROGEN.	Dis.	32.00																۰		
02	Total.	888																		
00	B P,pud		æ. 	<u></u>); č	i in	Ř	95	şį.	÷;	G)	Çį	===	÷ ;	-	₹.	-	. 18	37	0
ITROGEN AS	bioniminoid -sid -sid	85.55 05.55	.17	. 19	8:5	1 2	.31	∞.	.17	€	çi	25	08:	.e.	57	-	કૃ	.30	89	\$3
NITROGEN AMMONIA	T'10T	<u>8</u> 5.	1.05	11	8: 8	32	19:	÷:	€ <u>₹</u>	÷.	œ.	9.	49	-	3.	333	.34	∞.	.50	.36
	FreeAm- monia.	ಕ್ರ	01.	ව <u>ි</u>	8, 3	5	30.		당		33		•		:	910.	.01	.033	:03	83 83
ED.	Bysuspd Matter.	8.88 8.88	25.5	5.3	20.57 x 57	16.7	12.4	3.0	6.3	00	9.1.	= 1		<u>.</u>	€5 ∞	. i.	5.4		<u>.</u> ;	<u>∞</u> .
OXYGEN CONSUMED.	By Dis-	∞ ∞ ⊙ ∞	6.9	6.4	ان د نه	3 60	3	- Paris	9	}~	-) (-:	ن ن ن	-	œ	9	·-}	:		<u>es</u> es
000	Thotal.	31.6	32.4	9	9 15		8.3	=	₹. 62	x. 0	χ. Ξ	$\frac{\infty}{2}$	2	27	0.	10.1	10 1	::		11.0
.91	Chlorin	3.6	÷	6.4	ाट । इत्रुं ३८	200	α. ic	6.3		x.	- 1	ان ان	- :		φ χ	7.6	9.8	6. 1	10.5	<u>∞</u>
on Ion.	bended Sus	1687.	1896.	1346.	1851.	1159.	G555	617.	367	<u></u>	9:	115.	9 8	. 72	=	- 98	136.	:	157.	51.
RESIDUE ON EVAPORATION	-bis-	135.	187.	191	200	174	179.	-514	214	200	197.	164	.161.	3	201		186.		38	173.
RES	Total.	1822. 2184.	9083	1540.	2006 2006 2006	1333	811.	33	× ×		3.21	3009	261.	600	??	300	33.5	:	319	358
	Odor	None ::	:		: :	:		:	*	• •		: :	: ;		*	. ,		:	:	:
	Color.	3. je	(5)	Ē.	_ '8								:	:						
APPEARANCE.	Sedi- ment.	VVH.	.H.1.1			Y.E.	VVIII	. H.		. I.	Ä,	N.			-	N.	7.		/.II.	M.S.
1.	Turb'y.	VVIII.	11.7.7			VVH.	VVIII.									VVIII.		HAA	11.1.1	MH.
OF	1899 Exami- nation.	July 7	÷ :	or i	= : :::::::::::::::::::::::::::::::::::	:	: :	ept. 1		000		65)cl. 5		:		Nor. 3			Dec. 1
DATE OF	9 ec. F	6.1		10	10.5		24	1		-		,	0 0	· *		98	C1 .		53	30 0
1	1899 Collec- tion.	July	:		A 10 2.	•	:		Sept.			,	Oct.				NOV.		97	19
No.	657	655	566	551	663	6.65.1	53	500	196	XXX	599	010	179	672	67:3	67.1	675	123	1133	

S., Slight. VS., Very Slight. VVS., Very, Very Slight. M., Medium. APPEARANCE-H., Heavy. VH., Very Heavy. VVII., Very, Very Heavy.

TABLE 187.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, AT CHAIN OF ROCKS, PUMPING STATION ST. LOUIS WATER WORKS.

Report of Adolph Gehrmann, D. B. BISBEE, Municipal Laboratory, Chicago, III.

No. of Bac.	per Cubic Centi- meter.	45,000 53,650	35,000	9.	2,309,500	28,000	26,666	8,000	40,650	24.000 25.000 20.0000 20.000 20.000 20.000 20.000 20.000 20.000 20.000 20.000 20.000 20.000	19,300	14,500	25.000		30,350	34,750	18,750	35,000	68,550
	Presence of Co		+	+	-	++	-	+	+	<u>!</u> -	- 	+	+		+	+	+	-+	+
	Tennper ria lo	37.	57	36.	3 3		32.	32.		57 S	200	21.	55	18	19.	જ	14.	10.	17.
	Temper of Wate	26.	26.	39.	29.	9.60	27.	27.	23	3,5	16.	15.	17.	17	17.	6	6	=	ထင်
	dgiəH ə1sV/	22.6	20.6	16.6	<u> </u>	5 CO	9.7	7.	9.9	0.0 0.0	0 00	4.5	20.	33.3	<u>ئ</u>	3.0	6.2	6.2	5.5
RO- N AS	Vitrates	.8:	.30	89.	9.9	9.8	.49	.40	:		2		02.			0.	:	92.	.30
NITRO- GEN A	Nitrites	.00	:	.011		38					.017			•		900:	•	:	.004
C EN.	pəpuəd -sng	8.8	1.98	1.66	$\frac{1.96}{46}$	1.45 1.45	1.42	.64	∞ ∞ 0	70°	55.0	.62	2	.34	.54	.54	•	.62	#:
ORGANIC NITROGEN.	Dis- solved.	84.48	.30														۰		.33
OZ	Total.	32.48	2.28	1.8	2.16		1.74	1.08	1.10	1.84 0.84	9 00	8.	8.	20.	.74	£.	:	88.	.76
	Bud, pud sns.	.90	18:	69.	08.	7.0	.766	.43	. 34 			98.	33	.25	.14	.20	.14	. 23	.19
EN AS	Albuminoid -sid -vios	88	91.	.16	.20	16	.164	.16	7.	<u>. 5</u>	<u> </u>	.14	15	.16	.14	- 10	- 19	.13	. 13
NITROGEN AS	T'10T	1.20	.97		8.8	8.8	.93	.58	24. 20.	.40 64	.4.	.40	.37	.41	.28	8	.33	.36	.32
	FreeAm- monia.	2.8	.10	.16	8,8	5.5	.02	: 1	 20:	60	~	:	:	•	:	•	30:	30.	.03
9	BySuspd	27.6 43.8	9.3	φ. •	9.0	0.00	1.3	0.1		4,00						5.4	:	6.1	8
GEN	pealos		6	4	٥. ٥	10	1-	30.	-			6.	-	9.	٠. ب	<u></u>		دن	
OXYGEN	By Dis-	တတ	9	ા	ທີ່		_	9	· oc	غا∝ تەر	0 03				4	<u>െ</u>	:		8
	Total.	18.13 18.13		8													_	133	10
.9n	Chlorin	ţ			ر ب	ا ا ا	5	÷.	(ł		0	12	<u>2</u>	17	14	13	13.2	10.6	13.
ON NON.	pepued -sng	2260. 2492.	2374.	1974.	2252	1714.	1133.	608	573	5127	148	446	385	321	339	386	:	374	310
RESIDUE ON EVAPORATION	Dis- solved.	196. 172.	188.	189.	210. 189	155.	182.	219.	3000	924	230.	256.	276.	274.	274.	265.		249.	256.
RE	Total.	2456. 2664.	2562.	2163.	2462.	1869.	1315.	1028	833	737	378	702.	661.	595.	613.	649.		623.	.994
the same	Odor	None	* *	· .	9 9	;	7 9	us 19		: :	3 9	9 9	***	**	*,	9.9	**	9 9	* *
	Color.	યં.ફ	.05	.0.	₁ 2	3		:	:	:					•	:			
APPEARANCE.	Sedi- ment.	VVH.	VVH.	VVH.	VVH.	VVH.	VV11.	VVII.	. H		VII.	V.H.	11.	VH.	VH.	VH.	N.	VH.	11.
AP	Turb'y.	VVH. VVH.	VVII.	V.V.H.	VVH.	V.V.H.	VVH.	VVH.	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	H	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVII.
SOF	1899 1899 Collec- Examition.	July 7	160	3	Aug. 4	.: 18		31 Sept. 1		C C		Oct.	99	3 7	*,	Nov.	•	9.9	Dec. 1
DATE OF	1899 Collection.		02 .,	(C)	Aug. 3	.,	:		Sept. 7		80		12		98	Nov. 2	13	93	1 0g ,,
No.	Serial	632	634	635	636	638	633							647				1128	1132

M., Medium. VVS., Very, Very Slight. VS.. Very Slight. S., Slight. VVH.. Very, Very Heavy. VH.. Very Heavy. APPEARANCE—H., Heavy.

TABLE 188.

Serial No

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO

ANALYSIS - PARTS PER MILLION SANITARY WATER SOURCE OF WATER-MISSISSIPPI RIVER, AT CHAIN OF ROCKS, INLET TOWER, ST. LOGIS WATER WORKS.

D. B. BISBEE, Municipal Laboratory, Chicago, Ill

Report of ADOLPH GEHRMANN,

27,323 50,000 27,500 21,500 21,000 21,000 26,730 10,350 30,000 11,650 13,750 13,750 per Cubic Centi-meter. No. of Bac. Presence or Abs. of Coll. Temperature ('. of Air, ('. 1910226226226226236232 Temperature of Water, C. 8 1 9 6 6 8 1 7 1 6 1 6 2 6 1 1 1 1 1 6 6 1 2 Helght of Water. 28277788 01, ន្តខ Selutiin NITRO-GEN .003 8888 888888 017 Nitrites 238837888 ORGANIC NITROGEN. pepued sns solved 88 -sid :88 Total. कां कां कां कां कां कां कां कां कां कां papud sus Albuminoid Am NITROGEN AS Dis-88999977388838889899938 84486686666666666868888888888 ['10'] FreeAm. 33 OXYGEN CONSUMED. Mrsuspd Bratter 10 ** By Dis-0) -Total 000 တတ္သင္ Chlorine. benrqeq 452 350 RESIDUE ON EVAPORATION. -sns paajos -SIU 23.115. 2.25.25. 2.25. 2.25. 2.25.25. 2.25. 2.25. 2.25.25. 2.25.25. 2.25.25. 2.25.25. 2.25.25. 2.25.25 Total 88 88 Odor Color -888-8 APPEARANCE Sedi-ment Turb'y. 1899 Exami-nation. 일었고 그 첫 원 -weggenesse=== DATE 1899 Collec-616 Sept. .1119. Oct.

Medium

. IV

Very Slight.

VVS., Very.

Slight.

Very ?

Slight.

Ś

VVH., Very, Very Heavy.

Very Heavy.

VH.,

Heavy.

APPEARANCE-H.,

TABLE 189

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Source of Water-Mississippi River, 400 Feet from Missouri Shore, at Chain of Rocks, Municipal Laboratory, Chicago, Ill.

No. of Bac.	per Cubic Centi- meter.	## ## ## ## ## ## ## ## ## ## ## ## ##
	Presenc	1-++++
	rempers,	
r, C.	Tempers	82288882222886555556665
	Helgh etaW	88883141111 5000049110000000000000000000000000000000
RO-	Nitrates	
NITRO- GEN A	Nitrites	
ORGANIC NITROGEN.	Sus- bended	\$644295351588875951
ORG	Total.	%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
	Bug-sng	888888888888888888
NITROGEN AS AMMONIA.	Piori piori piori piori piori piori	<u> </u>
NITRO	I'10'T	<u>85806865848448889888888</u>
	FreeAm- monia.	4.85-1-15445 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
EN (ED.	BySuspd Matter.	80088888888888888888888888888888888888
OXYGEN	By Dis- solved.	<u>ಸ್ಥರ್ಣ ಇಟ್ಟು ಎ.ಎ.ಎ.ಎ.ಟ್ರಾಟ್ ಟ್ರಾಟ್ ಟ್ರಾಟ್ ಸ್ಟ್ರಾಟ್ ಸ್ಟ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ ಪ್ರಾಟ್ </u>
03	Total.	80000000000000000000000000000000000000
'əu	Chlorin	<u>*************************************</u>
ON NON.	papuad -sng	28119 29173 29173 29173 29173 29173 2917 2917 2917 2917 2917 2917 2917 2917
RESIDUE ON EVAPORATION	Dis- solved.	20000000000000000000000000000000000000
RE EV	Total.	323 3323 3323 3323 3323 124 1107 1107 1109 1009 692 692 692 692 692 693 693 693 693 693 693 693 693 693 693
	Odor	None
	Color.	-288-8
APPEARANCE	Sedi- ment.	
AP	Turb'y.	HE CANAL THE CONTROL OF CONTROL O
OF	1899 Exami- nation.	July 7 ** 28 Aug. 4 ** 28 ** 18 ** 18 ** 28 ** 28 ** 28 ** 28 ** 28 ** 28 ** 28 ** 28 ** 28 ** 18 ** 18 ** 28 ** 18 ** 28 ** 18 ** 28 ** 28
DATE OF	1899 Collec- Frion.	533 July 6 July 533 (27 %) 534 (27 %) 535 Aug. 27 % 536 Aug. 3 Aug. 538 (17 %) 539 (27 %) 540 (27 %) 541 (28 %) 541 (28 %) 542 (27 %) 543 (27 %) 544 (28 %) 545 (27 %) 546 (27 %) 546 (27 %) 547 (28 %) 548 (28 %) 549 Nov. 2 Nov. 550 (28 %) 540 (28 %) 541 (28 %) 541 (28 %) 542 (28 %) 543 (28 %) 544 (28 %) 545 (28 %) 546 (28 %) 546 (28 %) 547 (28 %) 548 (28 %) 549 (28 %) 540 (28 %)
No.	Seriai	11.000 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0

M., Medium. VVS., Very, Very Slight. S., Slight. VS., Very Slight. APPEARANCE-H., Heavy. VH., Very Heavy. VVH., Very, Very Heavy.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Adolph Gehrmann, D. B. Bishee. Municipal Laboratory, Chicago, Ill.

SOURCE OF WATER-MISSOURI RIVER, FORT BELLEFONTAINE, WEST ALTON, MO.

No. of Bac.	Per	Centi- meter.	24.650	771,750	38,250	30.750	36.250	17,000	52,000	56,750	61,750	60,750	18,500	31,500	18,500	38.500	37.250	25.000		47.750	47,500
ce or Coll.		Pres.	+	+	+	+	+	+	+	+	+	+	-	+	+	-+	-+	-+	- :	+	
		z jo luoj,		34.	.30	33.	%	53	34.	150	33	17.	3	138	19.	33	0	.33	.6I	.0I	17.
		TempT W lo	99	65	53	88	88	89	653	33	.08	-8	<u>-</u> -	33	200	19.	=	12.	12	=	10.
10 J	113 113		13.3	13.4	11.6	£.:	ã.6 6	∞ ∞	£-	6.7	6.1	5.5	ت	∞. **	4.6	4.5	17	10	7	4.9	ъ.
RO-NAS	ธอ	Mitrat	8.	01.	04.	1.00	.70	8.	:	:	8.		•	8.	-		0.	04.	.40		.30
GEN A	89	MICHE				.003			:	:	:	:	:	۰	0.		:		:		800.
OGEN.		solve solve		હાં	ঞ	4 2.34	€3	_				_	_		_	_	_				
ORGANIC NITROGEN.	-	RIOT'	1.88	80	72	84	88	38													
	Am.	p,pud -ang	·	-		.98	-	_												88:	
EN AS NIA.		Dis- solvid	80.	2	80.	<u>3</u>	. I5	01:	. I3	8.	91.	.15	80	=	=	80.	88	=	e(;	80	80.
NITROGEN AS	Albuminoid	Toty	1.20	1.10	.93	1.10	20.1	8	.45	.45	.46	.41	* * * * * * * * * *	.33	.31	.24		.32	66.	.37	0g.
Z		Freeki	.01		.03	80.	.03	:	:	:		910.	•	:	:	:	•	910.		ව ව	8.
EN.	pd pd	BySus	99.9					•	12.1											9	6.2
OXYGEN CONSUMED.		By Dis	3.3	က	_	3.4		9		က	જ	જ	6 2.2	-	જ	જ	65		63	5 9.5	
	!	Total	33						. 14.		-		_		_		_	_			
	1	орцо		5	-			-	-		_	_								-	
E ON ATION.	-	sng	. 2873	•	-	. 2514.	_	_	_	_	-		:		_					6230	-
RESIDUE ON EVAPORATION		sl(I evfos	224	٠	_	3. 199	_	-0			_	_	•			_	-	_	_	313.	
——————————————————————————————————————		B10T	3097.	-	324	27.15	1886	1468	115	125	Ξ	3,	:	818	-	E	£	686	676	952	78
	Odor		None	9.9	9.9	;	91	9.9	9.3	9 4	9.9	9.9	7.7	9.9	9.9	1,1	7.9	9.9	ud d	id d	
		Color.	.05	ક.	B.		:	:	:	•			:			:		:	:		:
APPEARANCE		Sed!-	VVH.	TI.	1.1.H.	.Y.H.	V.V.H.	LY.H.	V.V.H.	W.H.	L'YH.	VV.H.	VVH.	VH.	VH.	V.H.	VH.	V.H.	V.H.	V.H.	V.II.
A.		Turb'y.	VV.H.	VVH.	VVII.	L'VII.	H.	T. H.	VVII.	LVH.	V.H.	VVII.	T.Y.H.	V V.H.	L'VIII.	VVIII.	VVII.	VVH.	V.V. 11.	VVH.	VVIII.
DATE OF		Exami- nation.	July 28			. 100	25		3	99	-d -d									33 : 24	Dec. 1
DAT	1899	Collec- tion.	July 27	Aug. 3	01 .,	:	4 .	4 4	Sept.	4	d -0		Oct.		4 0		NOV.	2 2	11	4	1
.o.X.	451	45%	1	1	455	456	457	25.5	459	460	461	46%	463	164	465	466	467	468	469		

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Silght. VVII.. Very, Very Heavy. VH., Very Heavy. APPEARANCE-H.. Heavy.

TABLE 191.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, 100 VARDS FROM ILLINOIS SHORE, JEFFERSON BARRACKS, MO.

Report of Adolph Gehrmann,
D. B. Bisbee,
Municipal Laboratory, Chicago, Ill.

No. of	per	Cubic Centi- meter.	49,000	10,250	80,230 80,950	27.750		25,250	28,000	100,000	52,000		25,750	43,750	35,250	29,000	38,000	27,250	15,750	25,500	34,000	15,750
.iloo)](Presend Abs. of		1 -	1-1	- 1	:	1	+	+	+	:	+	-+-	+		+	-	+	+	-	+
		odm9T A 10		:				:		:	:		:	:	:		:	•		:	:	:
	Temperat ,TetaV lo		38	80.00	R	29	29.	30.	30.	30.	33	19.	17.	16.	ട ജ	19.	19.	6	6	10.	12.	·6
		BlaH BW	23.6	19.8	7.01	14.4	13.9	8.4	7.5	6.3	6.8	6.7		4.1	ယ က်	3.2	8.3	4	5.4	6.5	6.6	5.6
-0% N A 8	86	Mitrate		38	00.4	38	02.	38.	.40	.36	:	0.	:	•	:	:	0.	:	.40	55	98:	:
NITRO-	86	Nitrite		98	•			800.		:	:	:	:	:	:	•	900.				800.	•
IIC JEN.	pa	-sng	° .	٠.	18	03				.40									•		<u>8</u>	
ORGANIC NITROGEN.		Dis-		_	3.8				-	.40		_			Ľ				:		02.	
OZ	1	[gioT	39.6	<u> ২</u>	38	2.3	1.48	1.5	∞ .	<u>∞</u> .	<u>~</u>	8 6.	<i>ਰ</i> ;	<u>×</u> .	<u>۲</u> .	1.1	<u>×</u> .	∞. -	:	<u>86.</u>	1.8	<u>ক</u>
	1 Am.	-su8	1.38	33.2	3.6	8.	.54	.42	. 33	.21	.17	.34	8.	83.	.15	.21	.21	8.	.16	.27	.24	.17
NITROGEN AS AMMONIA.	Albuminoid	p'vios	.18	91.	16	.13	.16	.18	.14	83	: :33	.24	83.	83.		,25	.23	22.	98.	<u>8</u>	.29	.21
NITROGAMM	Albu	Total	1.56	1.18	200	.8:	.70	39.	.47	.43	.40	.58	.49	.48	.48	.46	.44	.44	.42	.52	.53	88.
		FreeAr inom	88.	98	3.6	8	경.	30.	1	.024	:	:	:	:	:	:	:	.03	₹O:		8.	.13
S.D.	DQ T.	Bysus	9.1		9.03 50.03								4.		4.3		3.3		2.9		3.8	2.6
OXTGEN	·p	By Dia	က	7 -		9	30	-	∞	-									9.1			
OXYGEN N	-	[BIOT	₩.	4	86.		_	4	7		00	∞	C3	9	9	_	<i>c3</i>	_		<u>03</u>	4	63
.91	nin	СРЈО			o 70 O 70																	
ON.	pa	puəd sns	2368.	1826.	1773	2446.	1215.	674.	679.	345.	2003 3003	151.	121.	•		113.	88	112.	114.	8	97.	61.
RESIDUE ON EVAPORATION		Solved	188.	191.	211.	199.	178.	186.	215.	196.	208.	198.	182.	:	211.	204.	202	211.	179.	153.	168.	172.
RESEVAF	-1	Total	2556.	2017.	1985	2645.	1393.	860	894	541.	410.	349.	303.	:	:	317.	330	323.	293.	213.	265.	233.
	Odor		None	; ;	: 1	9.9	9.0	9.9	99	3,	9 9	"	99	3,	,	;	99	9.9	,,	99	99	,,
		Color.	.05	કે ક	3.8	.e.		:	:	:	:	:	:	:	:	•	:	:	:	:	:	
APPEARANCE	-	Sedi- ment.	VVH.	V V H.	VV.H	VVH.	VVH.	VH.	Ë	VH.	VH.	M.	M.	M.	M.	M.	H.	H.	M.	ιά	M.	MH.
AP		Turb'y.	VVH.	V V H.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VVH.	VH.
DATE OF	1000	Exami- nation.	July 17	99	Ang.	14	** 18	99	Sept.	6 3,	"	9.9	9.9	Oct.	9.9	9 9	99	Nov.	9,9	,,	11	Dec.
DAT	10001	Collec- tion.	July 14	5 5			17	25	31	Sept. 8		22		Oct. 6		61 ,,		Nov. 3	111	16	23	08 ,,
No.	[[Seria	850	836	917	923	919	912	_	_	-	923	924	930	936	242	846	941	955	1481	1482	1488

M., Medium.

VS., Very Slight. VVS., Very, Very Slight.

S., Slight.

VH., Very Heavy. VVH., Very, Very Heavy.

APPEARANCE—H., Heavy.

TABLE 192.

Report of ADOLPH GEHRMANN, STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, EAST OF MIDSTREAM, JEPFERSON BARRACKS, MO.

	111.
	Chicago,
KISBEE,	Laboratory,
D. 15.	Municipal I

No. of Bac.	Cubic Centi- meter.	23.500 46,000 112,000 112,000 113,000 113,000 113,500
	Presen 10 .sdA	++ + + + + + + + + + + + + + + + +
ature ., C.	regmeT rik to	
	Temper otrW lo	88888888888888888888888888888888888888
	Helgh Wate	44600000000000000000000000000000000000
NITRO- GEN AS	Nitrates	S. S. S. S. S. S. S. S. S. S. S. S. S. S
	Nitrites	0.000000000000000000000000000000000000
TIC GEN.	-sng	8.8.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2
ORGANIC NITROGEN.	Dis-	2.08 11.08
	Total.	
Ø	A sus	4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4. 4
NITROGEN AS AMMONIA.	Pioniminoid Pivios Pivios	
NITRO	Totyl	88 48 48 48 48 48 48 48 48 48 48 48 48 4
	FreeAm- monla.	89.99.89.89.89.99.99.99.99.99.99.99.99.9
ED.	Hysuspd Matter.	21.00.00.00.00.00.00.00.00.00.00.00.00.00
OXYGEN CONSUMED.	By Dis-	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
CON	Total.	888887-1183881-100037557-
.9n	Chlorin	### 0 0 0 0 14 0 0 0 14 14 0 0 0 0 0 0 0 0
ON ION.	papuad -sng	2833 2893 2893 2893 280 280 280 280 280 280 280 280 280 280
RESIDUE ON EVAPORATION.	Dis-	200. 200. 200. 200. 200. 200. 200. 200.
RE: Eva	Total.	22.63 22.63 23.63
	Odor	None
	Color.	8.2
APPEARANCE.	Sedi. ment.	V V V V V V V V V V V V V V V V V V V
A	Turb'y.	V V V H I V V V H I V V V V H I V V V V
DATE OF	1899 Exami- nation.	Aug. 44 Sept. 95 Sept. 97 Sept. 77 Nov. 27 Dec. 24 Dec. 25
DAT	1899 Collec- tion.	Aug. 3 171 171 171 171 171 172 173 174 175 175 175 175 175 175 175 175 175 175
.oN	Serial	2

M., Medium.

VVS., Very, Very Slight.

VS., Very Slight.

S., Slight.

VVH., Very, Very Heavy.

VH., Very Heavy.

APPEARANCE-H., Heavy.

TABLE 193.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-MISSISSIPPI RIVER, MIDSTREAM, JEFFERSON BARRACKS, MO.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

No. of Bac.	per Cubic Centi- meter.	26.000 27.000	
	Presenc	+++++++++++++++++++++++++++++++++++++++	1
ture C.	rempers,		lium.
	rempers etaW lo	888888888888888888888888888888888888888	M., Medium
lo t	HelglaH etaV/	8355448000000000000000000000000000000000	
RO- N AB	Nitrates	\$8.58.88.6 S S S S S S S S S S S S S S S S S S S	Slight
NITRO	Nitrites	000 000 000 000 000 000 000 000 000 00	, Very, Very
EN.	pepued -sng	6.21.21.1.1.1.1.1.2.2.2.2.2.2.2.2.2.2.2.	ery,
ORGANIC NITROGEN.	Dis- solved.	888.9888888888888888888888888888888888	-
OZ	Total.	8.6.2. 8.6.2.	V VS.
	e sus	6:00 6:00 6:00 6:00 6:00 6:00 6:00 6:00	bt.
NITROGEN AS AMMONIA.	Piori Dis- Bla- Bla- Bla- Bla-	7.14.2.1.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2.	y Slight.
NITRO	T'10T	5-21 5-21 5-21 5-21 5-22 5-22 5-22 5-22	S., Very
	FreeAm- monia.	88.838.828.83 9.938 9.938 9.938.83 9.938.83 9.938.83 9.938.83 9.938.83 9.938.83 9.93	VS.
N SD.	Bysuspd Matter.	######################################	Slight.
OXYGEN	solved.	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	
OXYGEN CONSUMED	Total.	250.8 25	Š
,en	Срјоци	8.0.5.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.	leavy.
ON ION.	pepued ens-	2672. 2265. 2265. 2265. 2265. 260. 260. 260. 262. 262. 262. 275. 275. 275. 275. 275. 275. 275. 27	Very I
RESIDUE ON EVAPORATION	Dis- solved.	199. 174. 174. 174. 174. 174. 174. 174. 174	VVH., Very, Very
RE Eva	Total.	2871. 1937.	VVH.,
	Odor	None	vy.
	Color.	<i></i>	VH., Very Heavy.
APPEARANCE.	Sedi- ment.	V V H. V V V H. V V V H. V V V H. V V H. V H. V H. V H. V H. V H. V H. V H. V H. V H. H. V H. H. H. H. H. H. H. H. H. H.	VH.,
AP	Turb'y.	V V H H H H H H H H H H H H H H H H H H	APPEARANCE-H., Heavy.
S OF	1899 Exami- nation.	25	ANCE-H
DATE OF	1899 Collec- tion.	Aug. 37 Aug.	APPEAR
.oV	Serial	8833 8833 8859 8859 8951 9917 9917 9921 9931 9934 9939 9939 1480	

TABLE 194.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, Ill.

SOURCE OF WATER-MISSISSIPPI RIVER, WEST OF MIDSTREAM, JEFFERSON BARRACKS, MO.

1		1	0	9		9	2	2	2	• (2	2	2	2	<u>ي</u>	35	8	9	9	2	:
No. of Bac.	Cubic	Centi- meter.	53,000	55,25				121,350			31,6	19,2	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	76,6	32,5	48,55	19,37	23,7	40,78	26,0	
ce or Colf.			1	1	+	+	+	+	1	-	+	+	+	+-	+	+-	+	+	+	+	:
	Temperature of Air, C.			:			:	:	:	:	:	:	:	:	:		:	:		:	:
ature er, C.		Temy W 10	33	ි. ලි	8	8	30.	200		<u>.</u>	17.	9	3	<u>.</u>	6	တံ	6	10.	120	Gi	:
t of	d ₂									6.7					23 ·	- -		6.5			
RO-NAS	89	Mitrat	9.	.49	9.1	8	3	.36		क्ष		:	:	:	0.	:		. S	02.	:	:
NITRO GEN.	89	Mitrit			Ť.	_		:			.03 660			Ľ	.01 <u>2</u>	•	.007		:	:	:
HC GEN.		sng	2.38	es.	οį		•			·	Ċ		_				•	_			:
ORGANIC NITROGEN.		slU svlos	48 .20	Ĭ.											_	_		8 5			:
		r bnq Tota	CS	63	दश	C3	_	_	_			_		_	_	_	-	_			:
02	d Am.	-sns	.74	_	_			_			_					_	_	_	_		•
NITROGEN AS AMMONIA.	Albuminoid	Dis-	.16	.13	.18	.16	.18	.14	.14	.16	.16	.21	02.	.16	.16	. 19	.16	.26	.13	.12	
NITRO	Alb	['30T	8.	.93	1.05	1.8	.55	.45	<u>S</u>		4.4.	.45	.41	.45							:
		FreeAi	8.	.04	ਲ ਹ	30.	•	.024		•	•	:	:		:	8	.018	910.	30.	ð.	:
EN MED.	pd	BySusp	33	8 21.2	5	25	13.	3 6.9	6:	∞ ∞ ∞	φ.		3 6.1	٠.	4.4	3.4.6	÷	∞ ∞ ∞	4	5 4.4	:
OXYGEN		By Dis	4	0 3.8	4.	4	4	Θ.	4	~. 4.	 	6 4.8	큣.	4	00 	+		6 12.8			:
J	*1	Total	83	3	31	83	=	13.	_	13		=	10.	10	∞ —	<u>ه</u>		_		=	:
.9í	LĮI	СРЈО								9.6										13.4	:
ON TION.	pa	epued sns	2537.	2276.	1924.	1291.	916.	658.	719.	563	33.4		411.	381.	332.	362.	255	122.	332.	273	
RESIDUE ON EVAPORATION		Dis	225.	204	188	196	208	233	237.	245.	339		270.	249.	200	259.	251.	152.	2:3	264.	:
RE Ev		Total	27.62.	2480.	2112.	1487.	1124.	890	926	808	693.	:	681.	630.	594.	621.	509.	274.	575.	537.	
	Odor		None	9 9	9.0	9 9	. 91	9.9	9 9	9 4	9 9	9 9	9.9	9 9	9.9	9.9	9.9	,,	9.9	9 9	
		Color.	.69.	48.									•				•				
APPEARANCE		Sedi- ment.	VVH.	**	**	**	* *	**	:	4 4	* *	**	V1I.	:	:	:	**	N.	V111.	:	
AI		Turb'y.	VVH.	90	99	9.	**	:	:	**	:	9.9	**	* *	:	*	9.9	.,	:	:	
0 PF	000	Exami- nation.	A 112. 4	14	3	: :	Sept. 1		121	:	29	Oct.	4 0	9 9	9.9	Nov.	0.9	7 7	9 9	Dec.	:
DATE OF		Coilec- tion.	1112. 3	11	17	25	31	Sent. x		3	3	Oct. 6		19		NOV. 3		91	53		
No.	[1	Sorta	914	3	213	900	200	23	911	0.35	125	_		686	944	956	050	1478	1479	1485	

M., Medium.

VS., Very Slight. VVS., Very, Very Slight.

S., Slight.

VII., Very Heavy. VVII., Very, Very Heavy.

APPEARANCE-11. Heavy.

TABLE 195.

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION. SOURCE OF WATER-MISSISSIPPI RIVER, 100 YARDS FROM MISSOURI SHORE, Jefferson Barracks, Mo.

Report of Adolph Gehrmann, D. B. Bisbee, Municipal Laboratory, Chicago, III.

No. of Bac.	per Cubic Centi- meter.	10.500 10.5000 10.500 10.500 10.500 10.500 10.500 10.500 10.500 10.500 10.5000 10.5
	Presenc	+ +1+ 1+1++1++++++
	riedmer The lo	
r, C.	rempera etaW lo	8888888888888888
	Helght 918W	80004440001-000010400034100010 0001040410001-00014000341000
NITRO- GEN AS	Nitrates	(a) 1
N.r.	Nitrites	000000000000000000000000000000000000000
TC GEN.	-sng	6.50 1.50 1.00 1.00 1.00 1.00 1.00 1.00 1
ORGANIC NITROGEN.	Dis- solved.	8888858588888888888
ōZ —	Total.	89.4.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.
	E P,pud	
NITROGEN AS	Pior Pier Dis-	resissieresistesissessesis:
NITRO	T'10T	898.48.56.44.69.44.44.48.88.89.88
	FreeAm- monfa.	88888888
EN.	Bysuspd	######################################
OXYGEN CONSUMED.	By Dis-	αο + ωο + + + η υ + + η υ + + ω + η υ ο φ ιυ μ. μ. ω ω ο α + ω - η υ υ η μ. υ η μ.
OS	Total.	8898988758311100re9e0110 4 6 4648968878311100re9e0110
,9ı	Chlorin	4466644546800011187486888
ON TON.	pepuad -sng	2892 2892 2892 2892 2892 2892 2893 2893
RESIDUE ON EVAPORATION	DJs-	868.83.83.83.83.83.83.83.83.83.83.83.83.83
REEVA	'l'otal,	98861. 9744. 9746. 9746. 9776. 1177. 1177. 108.
	Odor	None contraction of the contract
	Color.	<i>ස</i> සිසිසි
APPEARANCE.	Sedi. ment.	VVH.
AF	Turb'y.	H.::::::::::::::::::::::::::::::::::::
OF	1899 Exami- nation.	Aug. 44 Aug. 48 Sept. 1 Sept. 1 Sept. 23 Oct. 7 Nov. 4 Dec. 1 Dec. 1
DATE OF	1899 Collec- E tion.	Aug. 37 Aug. 3
No.	Serial	840 865 865 865 865 865 865 865 865 865 865

M., Medium. VVS., Very, Very Slight. VS., Very Slight. S., Slight. VH., Very Heavy. VVH., Very. Very Heavy. APPEARANCE—H., Heavy.

TABLE 196

STREAMS EXAMINATION—SANITARY DISTRICT OF CHICAGO.

SANITARY WATER ANALYSIS-PARTS PER MILLION.

SOURCE OF WATER-ST. LOUIS, MO., TAP WATER.

Report of Adolph Gehrmann.
D. B. Bisber.
Municipal Laboratory, Chicago, Ill.

No. of Isac.	per Cubic Centi- meter.	25.000 25.000	
	Presenc	+ + + + + + + + + + +	
	Tempera		Medium.
eture r, C.	ragmaT otaM lo		M., Me
	Helgbl Mate		
RO-	Vitrates	5.08885.00	VVS., Very, Very Slight
NITRO- GEN A	Mitrites	700.00	Very
GEN.	-sns	888: .02.98.98.98	/ery,
ORGANIC NITROGEN.	Dis-	88848886848888888864 48848888888888888 4884888888888888	VS., 1
-	Total.		7
on.	A -sug	29888 88 8 5 5 88208	ght.
NITROGEN AS AMMONIA.	Tol'l Albuminoid	####################################	ry Sllg
NITRO	Tot.1	8 9 9 8 8 2 7 8 7 7 7 6 8 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	VS., Very Sllght.
	Freedm- monia.	8208 8	
SN (ED.	BySuspd	0.01 0.00 : : : : : : : : : : : : : : : : :	Slight.
OXYGEN	By Dis-	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	S. S.
0 ည	Total.	10 10 00 10 00 00 00 00 00 00 00 00 00	
.et	Chlorfr	4 r r r r r r r r r r r r r r r r r r r	Heavy
ON TON.	papuad -suS	######################################	Very
RESIDUE ON EVAPORATION	Dis- solved.	8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	Very,
RE Eva	Total.	255.00 25	VVIII Very, Very
	Odor	None None	75.
	or.	8	Hea
# W	Color	Æ	Very
APPEARANCE	Sedi- ment.		VH., Very Heavy.
APE	Turb'y.	/V/H,	eavy.
			f., H
OF	1899 1899 Collec- Examl- tion. nation.	3 Aug. 44 1177 118 285 28 282 286 0 115 287 115 288 11	APPEARANCE-H., Heavy.
DATE OF	1899 Collec- F	: 25.85 = 25.85 = 25.85 = 1 = 25.85 = 1	PPEARL
1017			A
No.	Serlal	######################################	







